



SERVICE CATALOGUE

with Recommended Patient Price (RPP)

CONTENTS

SECTION A GENERAL INFORMATION

	<i>Page</i>
Introduction	A-1
Useful Contacts	A-2
Service Hours	A-2
Test Request & Collection of Specimen	A-3
Guidelines for Specimen Collection & Handling	A-4
Result Turn-Around Time	A-5
Charging & Billing Policy	A-6

SECTION B PROFILE TESTS

General Screening Profiles

General Profile Summary 2025	B-2
Tag-on Tests for Screening Profile	B-3
General Biochemistry Screen	B-4
Foreign Worker Screen – Male	B-4
Foreign Worker Screen – Female	B-4
International Student Screen	B-5
Pre-Marital Screen Male / Female	B-5

Cardiovascular & Coagulation

Anti-Phospholipid Profile	B-6
Anti-Phospholipid Syndrome Studies	
Atherogenic Risk Profile – A	B-6
Atherogenic Risk Profile – B	B-6
Atherogenic Risk Profile – C	B-6
Atherogenic Risk Profile – D	B-7
Atherogenic Risk Profile – F	B-7
Cardiac Markers – A	B-7
Cardiovascular Risk Markers	B-7
Lipids Profile	B-7
Lipoprotein Electrophoresis	B-7

Endocrinology & Metabolic

Adrenal Evaluation	B-8
Ageing Evaluation – A	B-8
Ageing Evaluation – B	B-8
Ageing Evaluation – C	B-8

Aldosteronism Studies	B-9
Basal Metabolic Profile	B-9
Bone Markers	B-9
Catecholamine Fractions	B-9
Diabetic Diagnostic – A	B-9
Diabetic Diagnostic – B	B-9
Diabetic Function Studies	B-9
Diabetic Care Profile	B-10
Parathyroid Evaluation	B-10
Pituitary Evaluation	B-10
Thyroid Autoimmune Studies	B-10
Thyroid Function Tests – A	B-11
Thyroid Function Tests – B	B-11
Comprehensive Rejuvenation Profile	B-11

Gynaecology & Reproductive

Amenorrhoea Studies – A	B-12
Amenorrhoea Studies – B	B-12
Androgen Hormone Evaluation	B-12
Antenatal Screen – B	B-12
Antenatal Screen – C	B-12
Antenatal Screen – D	B-13
Antenatal Screen – E	B-13
Antenatal Screen – F	B-13
Antenatal Screen – G	B-13
Antenatal Screen – J	B-13
Cervical Cancer Risk - A	B-13
Cervical Cancer Risk - B	B-13
Hirsutism Studies	B-14
Fertility Hormonal Studies – Female A	B-14
Fertility Hormonal Studies – Female B	B-14
Fertility Hormonal Studies – Female C	B-14
Fertility Hormonal Studies – Female D	B-14
Fertility Hormonal Studies – Male	B-14
Menopausal Hormone Studies – A	B-14
Menopausal Hormone Studies – B	B-14
Ovarian Function Evaluation	B-15
Menstrual Cycle Studies	B-15
Prenatal & Hereditary Genetic Assay: Chromosomal Microarray Analysis (CMA) ~ CytoOneArray	B-16
Foetal Screen – 1 st Trimester Test	B-16
Foetal Screen – 2 nd Trimester Triple	B-16
Foetal Screen – 2 nd Trimester Quadruple	B-16
Pre-Eclampsia Risk at 1 st Trimester	B-16
Pre-Eclampsia Risk & Foetal Screen at 1 st Trimester	B-16
Pre-Eclampsia Risk at 2 nd & Late Trimester	B-16
Genetic Hereditary Risk Profile – A	B-16

Page

Haematology & Coagulation

Anaemia Studies – A	B-17
Anaemia Studies – B	B-17
Anaemia Studies – C	B-17
Iron Status Evaluation	B-17
Full Blood Count	B-17
Full Blood Count with CRP	B-17
Full Blood Picture with CRP	B-18
Full Blood Picture with ESR	B-18
Thalassaemia Studies – A	B-18
Thalassaemia Studies – B	B-18
Thalassaemia DNA Analysis (Alpha & Beta)	B-18
Thalassaemia DNA Analysis (Alpha, 7 mutations)	B-18

Thalassaemia DNA Analysis (Beta, 22 mutations)	B-18
Coagulation Studies	B-18
Thrombophilia Screen	B-18

Hepatobiliary & Gastrointestinal

Autoimmune Hepatitis Serology	B-19
Colitis Studies	B-19
Diarrhea Studies	B-19
Dyspepsia Studies	B-19
Hepatitis ABC Screen	B-19
Hepatitis B Carrier Studies – A	B-19
Hepatitis B Carrier Studies – B	B-19
Hepatitis B Screening	B-20
Hepatitis B Treatment Monitoring	B-20
Hepatitis C Treatment Monitoring	B-20
Hepatitis Immune Status Profile	B-20
Liver Function Test	B-20
Malabsorption Studies	B-20
Protein Electrophoresis- Serum/Urine	B-21
Viral Hepatitis Diagnostic	B-21

Infectious Diseases

Dengue Antigen & FBC	B-22
Dengue Duo Tests	B-22
Dengue IgG/IgM & FBC	B-22
Febrile Studies – A	B-22
Febrile Studies – B	B-22
HIV Monitoring	B-22
STD Studies – A	B-22
STD Studies – B	B-23
STD Studies – C	B-23
STD Studies – D	B-23
STD Studies – E	B-23
STD Studies – F	B-23
STD Studies – G	B-23
STD PCR Profile – A	B-23
STD PCR Profile – B	B-24
STD PCR Profile – C	B-24
STD PCR 7 Pathogens – A	B-24
STD PCR 7 Pathogens – B	B-24
TORCH Infections	B-24
TOUCH Screen	B-24

Paediatric

Neonatal Jaundice Studies - A	B-25
Neonatal Jaundice Studies – B	B-25
Neonatal Screen – A	B-25
Neonatal Screen – B	B-25
Neonatal Screen – C	B-25
Neonatal Health Screen	B-25

Renal & Urinary

Dialysis Indices Monitor – 1	B-26
Dialysis Indices Monitor – 2	B-26
Dialysis Indices Monitor – 3	B-26
Dialysis Indices Monitor – 4	B-26
Dialysis Virology Screen – A	B-26
Glomerulonephritis Serology	B-27
Renal Function Tests	B-27

Renal Function Studies	B-27
Estimated Glomerular Filtration Rate (with Cystatin-C)	B-27

Rheumatology, Immunology & Allergy

Arthritis Studies	B-28
Arthritis Screen	B-28
Extractable Nuclear Antigens Identification (6 subtypes)	B-28
Immunoglobulins Evaluation	B-28
Immunofixation Electrophoresis (Serum)	B-29
Immunofixation Electrophoresis (Urine)	B-29
Lupus Screening	B-29
Lupus Erythematosus Studies	B-29
Monoclonal Gammopathy Studies	B-29
Rheumatoid Screen	B-29
Total IgE, Eosinophil Cationic Protein, Phadiatop	B-30
Mix Allergens Panels	B-30
Specific IgE Individual Allergen Test	B-31
Allergy and Food Intoleranace Packages	B-32

Tumour Markers

Tumour Markers for Female (General)	B-33
Tumour Markers for Male (General)	B-33
Tumour Markers – Breast	B-33
Tumour Markers – Cervical	B-33
Cervical Cancer Risk	B-33
Tumour Markers – Colorectal	B-33
Tumour Markers – Liver	B-33
Tumour Markers – Lung (Small Cells)	B-34
Tumour Markers – Lung (Non-Small Cells)	B-34
Tumour Markers for Lung	B-34
Tumour Markers – Neuroblastoma	B-34
Tumour Markers – Nasopharyngeal	B-34
Tumour Markers – Ovary	B-34
Risk of Ovarian Malignancy Algorithm	B-34
Tumour Markers – Prostate	B-34
Tumour Markers - Prostate Health Index	B-34
Tumour Markers – Stomach & Pancreas	B-35
Tumour Markers – Testis	B-35
Tumour Markers – Thyroid	B-35

SECTION C
SINGLE TESTS
 (By Alphabetical Index)

	<i>Page</i>
Numeric (1, 2, etc.)	C-1
A	C-1
B	C-3
C	C-3
C Culture & Sensitivity Tests	C-5
D	C-6
E	C-6
F	C-7
G	C-7
H	C-8
I	C-9
J	C-10
K	C-10
L	C-10
M	C-10
N	C-11
O	C-11
P	C-11
R	C-13
S	C-13
T	C-14
U	C-15
V	C-15
W	C-16
Z	C-16
Toxicology	
Abuse Drugs Tests	C-17
Abuse Drugs Confirmation Assays	C-18
Therapeutic Drug Assays	C-19
Chemical Hazards	C-20
Trace & Toxic Elements	C-21
Cytopathology & Histopathology	C-22

Section A

GENERAL INFORMATION

INTRODUCTION

With the recent development of evidence-based medicine, individualised or targeted therapy, and the advancement of biomedical sciences and technology, more efficient, sensitive and specific diagnostic tests are highly needed to help clinician to make accurate and efficient medical decisions. For these reasons, *Gnosis Laboratories* was established in Malaysia in year 2002, as part of the state-of-the-art medical laboratory by a network of medical laboratory professionals from Malaysia and Taiwan.

History

Lezen Reference Laboratories founded in year 1979 in Taipei city by a team of medical laboratory professionals. Beside its headquarters in Taipei, Lezen have established regional laboratories in Kaoshiung and Taichung, and with numerous adjunct or satellite labs throughout Taiwan. In year 2002, *Lezen* expanded its network of medical laboratory services in Kuala Lumpur, Malaysia, with a new collaborated laboratory called *Gnosis Laboratories*, with the vision to expand its network and operation into Southeast Asia region.

Expertise

Our group of laboratories consist of over 100 staffs in Malaysia with half are qualified Medical Lab Technologists & Scientists; some are doctorate and master degree holders in various disciplines. We also have a team of information technology specialists to develop and maintain our laboratory information management system and the real-time online test result access facility. We also established strategic collaboration in specific areas with world's leading medical and research laboratories.

Advance

As a reference and research-lead medical laboratory, we focus on cutting-edge research and development of new assays. This includes refinement of existing diagnostic tests to produce assays with greater sensitivity, specificity, efficiency and clinical value for reliable and cost-effective clinical investigation. We pioneered nuclear medicine and radioimmunoassay technology in Taiwan. In year 1999, we have developed the molecular diagnostics services, which applied the latest technology available in molecular biology, such as polymerase chain reaction and genomic sequencing technology into clinical application. Thus, with these comprehensive clinical tests menu, we can support a wide range of clinical fields, namely Obstetrics & Gynaecology, Cardiology, Oncology, Haematology, Gastroenterology, Nephrology, Paediatrics, Endocrinology, Infectious Disease, Allergy & Immunology, Rheumatology, Neurology, Toxicology, Genetics and etc.

Quality

Accurate and time efficient lab diagnostic results to support the medical providers in decision-making are our main concern. Therefore, all our activities are subjected to good laboratory practice and stringent quality assurance programs. The laboratories in Taiwan are accredited by Taiwan Accreditation Foundation (TAF) according to ISO 15189 standards. Likewise, *Gnosis Laboratories* in Malaysia recognises its responsibility to provide quality services to better satisfy the needs of our customers and to improve the management of our services, thus we have developed a management system that complies to the requirements of MS ISO 15189 standard, and we are in the process to seek accreditation for a wide range of our testing activities.

Affordable

As an Asian pioneer medical laboratory, through years of experiences, we are able to develop cost-effective diagnostic tests with affordable price for the Asian market. These included those highly advanced and specialised assays, without compromising on our quality of test and service.

Partnership

We strongly believed in working together and partnership with healthcare providers & institutions, various exteem diagnostic products manufacturers, not just meeting needs but to work alongside with them to achieve their missions, create greater values and bring out the best in their medical practise, professionalism and entrepreneurship.

Our Mission

Our mission is to provide comprehensive and advanced medical laboratory and diagnostic services and solutions with an unmatched level of reliability, service and support, to the healthcare providers and institutions, for the best of the patient care.

Yours sincerely,



Jo-Kuan Yee

B. Biomedical Sc (Hons) (UKM), MIBMS (UK)

Laboratory Director

USEFUL CONTACT

Website: gnosis-healthcare.com

PENINSULAR

Subang Jaya (HQ & Central Lab):

1-1 & 1-2, Jalan USJ 21/11, 47630 Subang Jaya, Selangor, Malaysia.

Tel: +603-5885 8501 / 8502 / 8503 Fax: +603-5885 8510

e-mail: hqlab@gnosis-healthcare.com / business.national@gnosis-healthcare.com / jyemay@gnosis-healthcare.com

Ipoh: Hospital Seri Botani, Level 1, Pathology Lab, No.3, Dataran Botani 2, Bandar Seri Botani, 31350 Ipoh.

Tel: +605-226 2768

e-mail: ipohlab@gnosis-healthcare.com

Penang: 70-2-56A, D'Plazza Mall, Jalan Mahsuri, 11900 Bayan Baru, Pulau Pinang.

Tel: +604 -6428771 / +6010-2797500

e-mail: penanglab@gnosis-healthcare.com

Kulim: 9, Jalan Lunas, Taman Badlishah, Kelang Lama, 09000 Kulim, Kedah.

Tel: +604-4902315

Kulimlab@gnosis-healthcare.com

Johor Bharu: 10-A, Jalan Indah 1, Taman Bukit Indah, 81200 Skudai, Johor Bharu, Johor.

Tel: +607-2445177 Fax: +607-2445077

e-mail: jblab@gnosis-healthcare.com

Melaka: 9-1, Jalan KL 3/15, Taman Kota Laksamana, Seksyen 3, 75200 Melaka.

Tel: +606-2811288 Fax: +606-2811288

e-mail: melakalab@gnosis-healthcare.com

Taiping: B04 No. 62, Tingkat 1, Jalan Taming Sari, 34000 Taiping, Perak

Tel: 016-621 2477, e-mail: gnosis.taiping@gmail.com

Klang: 38A, First Floor, Jalan Batu Unjur 1, Taman Bayu Perdana, 41200 Klang, Selangor

Tel: 03-3311 5817, e-mail: klanglab@gnosis-healthcare.com

Seremban: 28-1, Jalan Rasah Prima 2, Pusat Komersial Rasah Prima, 70200 Seremban, Negeri Sembilan

Tel: 06-6316 788, e-mail: serembanlab@gnosis-healthcare.com

Kuantan: A9-A11, Mezzanine Floor, Jalan Dato Wong Ah Jang, 25100 Pahang

Tel: 011-1653 7500, e-mail: kuantanlab@gnosis-healthcare.com

SABAH

Kota Kinabalu: Lot A-1-1, First Floor, Block A, 88 MarketPlace, Jalan Pintas, 88300 Kota Kinabalu, Sabah

Tel: +6088-212 525

e-mail: kklab@gnosis-healthcare.com

SARAWAK

Sibu (Regional Lab): 14, Jalan Pedada, 96000 Sibu, Sarawak.

Tel: +6084-332600

e-mail: sibuadmin@gnosis-healthcare.com

Kuching: 33, Sublot 7, Lot 1108, Block 10, KCLD Kueh Hock Kui Comm Ctr, Jalan Tun Ahmad Zaidi Adruce, 93250 Kuching, Sarawak.

Tel: +6082-245996

e-mail: kuchingadmin@gnosis-healthcare.com

Miri: Lot 292, Ground Floor, Jalan Bendahara, Beautiful Jade Centre, Sarawak, 98000 Miri

Tel: +6085-439 068

e-mail: bintululab@gnosis-healthcare.com

Bintulu: 166, Parkcity Commerce Square, Jln Tun Ahmad Zaidi, 97000 Bintulu, Sarawak.

Tel: +6086-317400

e-mail: bintululab@gnosis-healthcare.com

SERVICE NOTES

SERVICE HOURS

Day	Routine Hours	*On-Call Hours
Monday to Friday	9:00 a.m. – 5:30 p.m.	5:30 p.m. – 7:00 p.m.
Saturday	9:00 a.m. – 1:00 p.m.	1:00 p.m. – 3:00 p.m.
Sunday	-	9:00 a.m. - 3:00 p.m.
Public Holiday	Close	To be notified

Overnight on-call or off days on-call can be honoured by special arrangement.

* Surcharge is applicable on such services.

Above service hours are applicable to HQ only. Please check with particular branch for the exact hour and working day.

TEST REQUEST & COLLECTION OF SPECIMEN

- Our courier will make the routine specimen collection **twice a day** (morning or afternoon) for normal working day. Please kindly inform us if you need different arrangement for routine specimen collection.
- Urgent Requests-** We will respond immediately to urgent requests during our routine hours. Urgent specimens should be packaged in our red coloured marked as Urgent packaging specimen bags. It is **ESSENTIAL** to tick the urgent box on the request form and that contact details are provided for results to be telephoned, fax or email when required. For urgent specimen collection, please phone our service line as stated above.
- During **On-Call Hours**, our laboratory will only respond to urgent tests. An extra **RM 4.00 Surcharge** for urgent tests will be charged per case send within On-Call Hours. If Sunday collection for non-urgent or routine tests, pre-arrangement is needed, please contact your service branch or service representative for such arrangement.
- Submission of requests for laboratory testing shall be accompanied by a GNOSIS request form. The **Request Form** must be filled-in clearly (type or hand printed). **Name, Sex, Age, IC number, Doctor/Clinic's stamp & Signature and date and time of Specimen Collection** are essential, **Time** of collection will be important for time sensitive parameters. **Clinically relevant information and drugs treated** would be most helpful in the interpretation of results, and for the lab to access the necessity for follow-up testing or alternative tests which are more indicative for a particular condition. Include the anatomic site o origin of specimen, where relevant.
- Gnosis **Tests Codes** should be clearly written on the Request Form to indicate the tests ordered. All phoned in verbal requests for additional test requests to be added to a written order should be followed by written confirmation as soon as possible.
- All specimen containers must be **labelled** with **patient's name** and a unique numerical identifier and accompanied with the Request Form. The specimen container must be tightly capped and put into the Biological Specimen Bag and zipped it. The Request Form shall be place into the "kangaroo" pocket.
- Use the **appropriate containers** for different types of tests and provide enough amount of sample. Specimen send in with incorrect container or improper/no labelling will be rejected. Refer to Gnosis Service Guide Guideline for further detail.
- If immediate delivery of specimen to laboratory is unavailable, please keep the specimens in the **suitable temperature**. Refer to Guideline for Specimen Collection & Handling and the Test List for suitable specimen storage conditions.
- If a test is ordered that can be performed by several **methodologies** and the method is not specified on the Test Request Form, the test will be performed by method which generally has the greatest clinical utility or availability.
- An **additional** or **follow-up test** can be ordered on a previously submitted specimen, provided there is sufficient specimen volume and the specimen still useable. In some cases, the laboratory personals may consult the referral physician and suggest a further investigational/differential/confirmatory test. In any case, written verification must be provided by fill-in the Test Request Form again with appropriate note and fax or courier to the laboratory.
- The laboratory will accept requests for **test cancellation** received prior to test set-up at no charge. Cancellation requests received after test set-up cannot be honoured; the test will be performed, a report will be issued, and the client will be charged appropriately.

GUIDELINES FOR SPECIMEN COLLECTION & HANDLING

The quality of laboratory results depends greatly on the proper collection and handling of specimen.

Collection of Blood Specimen

1. Red cells or white cells **haemolysis** in blood specimen may cause clinical invalid result. Always apply proper phlebotomy technique. Avoid water, alcohol or other agents to come in contact with the sampling needle or blood specimen. Do not apply sudden force in blood sampling or transferring; avoid transfer blood from syringe to blood tube through needle. It is advisable to use needle with gauge size 22G or larger (21G) to avoid haemolysis.
2. Avoid apply **prolong tourniquet**, prolonged venous stasis may cause increase in certain parameters.
3. Effect of **posture, medication, intravenous infusion** and **examination** of the prostate may alter or interfere with certain assays.
4. Blood specimen for electrolytes assay, especially potassium, blood for cytogenetics, leukocytes antigen, lymphocytic markers, cryoglobins and blood culture **should NOT keep refrigerated**. It should be delivered to lab as soon as possible.
5. Most hormones and enzymes will decrease in activity through time or in warm temperature. It should be **keep refrigerated** if immediate delivery to lab is unavailable.
6. An overnight (10 to 12 hour) fast is required for most **fasting** blood specimens.

Type of Vacuum Blood Collection Tubes (as supplied by our labs, specification may change)

Cap Colour	Type & Additives	Vol.	Note
Yellow	Plain SST (with Serum Separation Gel & Clot Activator)	5 ml	No need mixing, avoid shaking. Mainly for most routine biochemistry assays, hormones, tumour markers, serology assays, virology assays etc. Preferably using this tube rather the total plain tube, for speedy processing, unless specifically indicated.
Red	Plain (without Gel)	7 ml	No need mixing, avoid shaking.
Lavender	EDTA (K2EDTA)	3 ml	Gently Mix well. For most haematology assays, blood lead assay. If ESR, need min 1.5 ml.
Grey	Fluoride (KF+Na2EDTA)	2 ml	Mix well. For glucose test. Min 1 ml, too little blood may cause blood haemolysis. Too much blood may reduce glycolytic inhibition.
Green	Heparin (Na/Li Heparin)	2 ml	Mix well
Light Blue	Citrate (Buffered Citrate 3.2%)	2 ml	Fill to the marked line/specified volume and mix well. Ensure citrate to blood ratio is 1:9. Not to draw first blood for coagulation tests, either use the first draw for other tube or discard the 1 st tube.
Royal Blue w Red Label	Trace Element (TET) Plain	7 ml	No need mixing. For trace element or heavy metal assays on serum or urine.
Royal Blue w Lavender Label	Trace Element (TET) EDTA	7 ml	Mix well. For trace element or heavy metal assays on whole blood or red blood cells.
Royal Blue w Royal Blue Label	Trace Element (TET) Heparin	7 ml	Mix Well. For trace element specially blood Aluminium assay.

For vacuum blood collection system, the blood will fill to the stated volume automatically.

Collection of Other Specimen

Sterile Universal Container:

For Urine, Stool/Faeces, Sputum, Body Fluid, Effusion, Aspirate, Seminal Fluid, Scrapping, Small Tissue and etc.

Seminal fluid must keep in room temperature, avoid light and dispatch to laboratory within 1 hour of collection.

Stool or Urine for detection of life parasites should keep in room temperature, if for culture should keep refrigerated to avoid overgrowth of normal flora if immediate delivery to lab is unavailable.

Specimen for routine Urinalysis, Biochemistry assays should keep refrigerated if immediate delivery to lab is unavailable.

Body fluids for culture should keep in room temperature.

Glass Slide: For fixed smear. e.g. PAP smear, direct smears for bacteria, parasite and etc.

Swab, Plain: For direct or fixed smear, e.g. Gram's stain, AFB stain. For nucleic acid (PCR) detection. Keep in 2 - 8°C and transport to lab as soon as possible.

Swab, in Microbial Transport Medium:

For transport of microbiological specimens for culture & sensitivity tests from wound, pubs and etc. And also for wet mount preparation for the detection of Monilia & Trichomonas. Keep and transport in **Room Temperature** to lab as soon as possible.

For bacterial nucleic acid (PCR) detection, keep in 2 - 8°C and transport to lab as soon as possible.

Swab, in Viral Transport Medium:

For transport of viral specimens for nucleic acid (PCR) detection or viral culture. Keep in 2 – 8°C and transport to lab as soon as possible. Recommended to use flocced tip made with synthetic fibre, do not use cotton fibre swab.

Blood Culture Bottle (Aerobic or Anaerobic):

For either Aerobic or Anaerobic bacterial culture & sensitivity tests. Specimens keep in room temperature, do not refrigerate.

24 hr Urine Container:

For collection of 24-hour urine. **Acid Preservative (normally in 25ml 6M HCL acid or 10g Boric Acid)** has to be added for specific assay **and/or keep refrigerated during collection period**. Patient should be informed of any additive added in the container. Proper instruction should be understanding by the patient. At 0-hr, the bladder is emptied completely; discard the urine. Collect all urine passed until the 24-hr. At 24-hr, the bladder is emptied completely by urinating *into the collection container*.

Mid-Stream Urine (MSU):

Clean the genital area with water. Discard the first void urine and only collect clean voided mid-stream urine into container. This is important to obtain a clinical relevant result for Urinalysis and Culture & Sensitivity test. Please keep specimen in refrigerator if immediate delivery to laboratory is unavailable.

First Morning Urine (FMU) or First Void Urine (FVU):

First or second urine in the early morning or first void/stream urine is important for detection of certain urine's parameters, as the concentration is higher in the early morning urine or in the first void/stream urine. Please keep specimen in refrigerator if immediate delivery to laboratory is unavailable.

Special Collection Tube/Kit:

Specific collection tube or kit is required for certain test. E.g. liquid-based PAP test, Fibrin Degradation Product (FDP), Urea Breath Test, Hair Heavy Metal assays, Urine Metabolic/Amino Acids assays etc.

All the above specimen containers, needles, swabs, spatulas, brushes, slides, slide holders, fixatives (for cytology), request forms, specimen bags and service catalogue are supplied free of charge if use it to collect specimen for testing in our labs. Please order with us using the **Supplies Request Form** before they are required. May fax or hand to our dispatch personnel to deliver to our labs.

RESULT TURN-AROUND-TIME

1. The **Turn-Around-Time (TAT)** in **normal working day(s)** stated in the list for Profile Tests (Section B) or Single Tests (Section C) is for general reference only, and it mainly apply to HQ Lab. For branch labs, it may require an extra day for transportation. If any particular profile/test is time critical, please notify the particular lab/branch for attention.

2. Generally, the TAT is as follow:

	Type of test	TAT (working day)
Urgent	FBC, spot/rapid test for infection (Dengue, ASOT & WWF), common biochemistry (Liver Function Tests, Renal Function Tests, Glucose), Urine FEME, etc	Same day* or 1 day * Surcharge may apply
Routine	Common test for biochemistry, hormones, serology & tumour markers; Culture, routine coagulation tests, cytology & histology, etc.	1 to 5 days
Esoteric	Special hormones/proteins & serology, molecular diagnostic, AFB culture, etc.	7 days or more

3. The **Laboratory Report** will be printed and send to you by our courier after all the results are available. Upon request, **Interim Report** for routine test may be delivers before esoteric test result is available. Final report will be delivers once all the results are available. Two copies are supplied, one for your record and one for the patient. *Please inform us if you need different arrangement for results reporting.*
4. All indicated urgent tests will be **email, fax or phone** as requested to attending doctor immediately after the results are available and the printed report will be sent to client on the next routine dispatch trip available. Any critical results will be conveyed to the ordering clinician as soon as possible.
5. **Urgent Tests & Surcharge:** Special courier trip for urgent pick-up of specimen and result inform back to client (via email, fax or phone) within shortest possible time can be arranged with surcharge applicable as blow:

Tests/ Profiles	TAT*	Surcharge
Full Blood Count, Full Blood Count w CRP, Dengue Fever Tests (DEN1, DEN2, DEN3), Urine FEME	3 hours	RM4.00
Full Blood Picture w ESR, Febrile Studies (FEBA, FEBB)	4 hours	RM4.00

**Above TAT are indicate for HQ only, for branches, please arrange with particular branch that serve your area.*

CHARGING & BILLING POLICY

1. All Laboratory Fees listed in this catalogue are in **Malaysia Ringgit (RM)**. Whenever possible, prior notice will be given for fees revision. However, the company reserved the rights to adjust any fees with immediate effect or notify after the changes especially on non-routine test that referred out to other lab. The fees stated are not inclusive of any statutory or governmental taxes or charges, if such charges applied, it shall be added on in the billing.
2. The fees indicated as **Doctor's Fee** are fee chargeable to clinic or hospital, only inclusive for test analysis and report, specimen collection and delivery of report within local area and relevant specimen collection container (some are charge separately). It does not include specimen collection from patient or phlebotomy service for patient, out-station specimen collection & report delivery and interpretation of report to the patient. Any such service required shall be charged additionally.
3. The fees indicated as **Recommended Patient Price (RPP)** or **Patient's Fee** are recommended retail prices that our clients may charge on their patients. However, those are just recommended or indicative prices and our clients have full discretion on the actual price to charge on their patients, unless notified otherwise or impose by any statutory regulations.
4. Credit term is **30 days** from the date of invoice. The policy will be strictly enforced. It will be much appreciated if invoices are settled timely. Interests will be applied on overdue payment.
5. Please check your invoices upon receipt and notify us of any **discrepancy** within 10 days. Your statement will be re-process immediately or a separate credit/debit note will be given.
6. The payment should be made by crossed cheque or electronic transfer payable to name as indicate in the invoice:
GNOSIS LABORATORIES (M) SDN BHD for Peninsular Malaysia.
GNOSIS LABORATORIES (SABAH) SDN BHD for Sabah, Malaysia.
GNOSIS LABORATORIES (S) SDN BHD for Sarawak, Malaysia.
7. Our service representative / marketing staffs and couriers are authorised to collect crossed cheque payment on behalf of the company.

Please kindly contact our Service Representative / Manager for any inquiries regarding pricing or any other client's service matters. For invoices, statement and payment can be directed to our account staffs on our service line within normal office hour.

For further guide details, please refer to **Gnosis Service Guide or contact our clinician service representatives (medical sales representatives) or laboratory managers.*

Section B

PROFILE TESTS

General Profiles Summary 2025 (Promotion effective 1 Jan 2025 till further notice)

Test Name	Code	GEA	GE3	GEB	GEF	GEM M / F	GE72	GE75 M / F	GQY3	GHY3	GQY5	GQY8	GHY8
Full Blood Count		√	√	√	√	√	√	√	√	√	√	√	√
Blood Group	ABORH								A	A	A	A	A
Liver Functions													
Bilirubins		√	√	√	√	√	√	√	√	√	√	√	√
Total Protein		√	√	√	√	√	√	√	√	√	√	√	√
Albumin		√	√	√	√	√	√	√	√	√	√	√	√
Globulin		√	√	√	√	√	√	√	√	√	√	√	√
ALP		√	√	√	√	√	√	√	√	√	√	√	√
AST (SGOT)		√	√	√	√	√	√	√	√	√	√	√	√
ALT (SGPT)		√	√	√	√	√	√	√	√	√	√	√	√
γ-GT		√	√	√	√	√	√	√	√	√	√	√	√
Renal Functions, Diabetes & Metabolic													
Urea		√	√	√	√	√	√	√	√	√	√	√	√
Creatinine		√	√	√	√	√	√	√	√	√	√	√	√
Uric Acid		√	√	√	√	√	√	√	√	√	√	√	√
Calcium		√	√	√	√	√	√	√	√	√	√	√	√
Phosphate		√	√	√	√	√	√	√	√	√	√	√	√
Magnesium		√	√	√	√	√	√	√	√	√	√	√	√
Electrolytes		√	√	√	√	√	√	√	√	√	√	√	√
MicroAlb /Cre	UMAL			√	√			√		√			√
Serum Iron	SI			√	√	√	√	√	AS	AS	AS	AS	AS
Iron Status (Iron, Transferrin, %)	IRST												
Glucose		√	√	√	√	√	√	√	√	√	√	√	√
HbA1c	A1C		√	√	√	√	√	√		√			√
Lipids, Inflammation & CVD Risks													
Total Chol.		√	√	√	√	√	√	√	√	√	√	√	√
HDL-Cholesterol		√	√	√	√	√	√	√	√	√	√	√	√
LDL-Chol. (Direct if TG high)		√	√	√	√	√	√	√	√	√	√	√	√
Triglycerides		√	√	√	√	√	√	√	√	√	√	√	√
Total/HDL-Chol		√	√	√	√	√	√	√	√	√	√	√	√
hs-CRP		√	√	√	√	√	√	√	√	√	√	√	√
LDH	LDH								R	R	R	R	R
Apo-B & Apo-A1	APOP												
Homocysteine	HCY												
Hormones, Infections, Tumour Markers & Others													
TSH	TSH				√	√	√	√					
FT4	FT4												
FT3	FT3												
HBs Ag & Ab	HBS				√	√	√	√					
Anti-HAV	HAV						√	√					
HIV Ab+Ag	HIV						√	√					
√ TPAb (reflex RPR)	TPAB1							√					
√ RPR Screen (reflex TPAb)	RPR						√		V	V	V	V	V
H. pylori IgG	HPYG												
RF	RF				√	√	√	√	RSV	RSV	RSV	RSV	RSV
AFP	AFP					√	√	√					
CEA	CEA					√		√					
PSA (for Male)	PSA					√ M		√ M					
CA125 (Female)	125					√ F		√ F					
CA15-3	153												
CA 19-9	199												
EBV VCA IgA	EBVA												
Urine FEME	UFEM	√	√	√	√	√	√	√	√	√	√	√	√
Option Test #	A (ABS): ABO & Iron, R (RFLD): RF & LDH, S (RFSI): RF & Iron, V (VDRF): RPR & RF								3	3	5	8	8
Specimen for All		If profile with both ABO & HbA1c tests, need extra											
Promo Patient Price RM		56	68	86	130	184	172	228	110	142	156	220	256

Tag-On Tests for General Screening Profiles *(Promotional effective 1st Jan 2025 until further notice)*

Code	Test Name	RPP	Code	Test Name	RPP
Haematology & Anaemia					
IRST	Iron Status (Serum Iron, Transferrin, % Saturation)	30	ABORH	ABO & Rh Blood Group	12
FER	Ferritin	30	ABSI	Blood Group & Serum Iron	20
SI	Iron, Serum	10	ESR	Erythrocyte Sedimentation Rate	10
BFC	Blood Film Comment	6	HBE	Hb Electrophoresis	158
Renal Functions, Liver, Cardiovascular, Inflammation & Others					
A1C	HbA1c	26	UMAL	Microalbumin & Cret. Ratio	20
HCRP	C-Reactive Protein, high-sensitive (hsCRP)	20	CYSC	Cystatin C & eGFR cr-cys	90
HCY	Homocysteine (separate serum a.s.a.p.)	36	LDH	Lactate Dehydrogenase (separate serum a.s.a.p.)	10
APOP	Apo-A1, Apo-B & Ratio	36	TNIH	hs-Troponin I (w risk stratification, non-urgent only)	80
LPA	Lipoprotein (a)	96	SOD	Superoxide Dismutase (SOD)	90
Infectious Diseases					
HIV	HIV 1 & 2 Ab+Ag	20	H1G	HSV Type 1 IgG	36
TPAB1	Syphilis TP Ab (reflex RPR)	20	H2G	HSV Type 2 IgG	36
RPR	Syphilis RPR (VDRL) Titter	10	CLG	Chlamydia IgG	36
HBS	HBs Ag & HBs Ab	20	HPYG	H. pylori IgG	30
EAB	Anti-HBe	46	HPYA	H. pylori Abs (with Current Infection Marker)	60
EAG	HBe Ag	46	UBT4	14C Urea Breath Test	146
HAV	HAV Ab (Anti-HAV)	20	RG	Rubella IgG	30
HCV	HCV Ab (Anti-HCV)	40	VDRF	Syphilis RPR & RF	20
SC2GS	SARS-CoV IgG (anti-S1 RBD)	140			
Immunology & Serology					
RF	Rheumatoid Factor	12	ANAQ	Anti-Nuclear Antibody (Quantitative)	68
ACCP	Anti-CCP	80	ADNA	Anti-dsDNA (Quantitative)	98
ATPO	Anti-Thyroid Peroxidase	68	RFLD	RF & LDH	20
ATAB	Anti-Thyroglobulin Ab	68	RFSI	RF & Serum Iron	20
Hormones & Others					
TSH	Thyroid Stimulating Hormone (TSH)	20	TES	Testosterone, total	34
FT4	Free T4	20	SHBG	Sex Hormone Binding Globulin	60
FT3	Free T3	20	HOMA	Insulin with HOMA-IR	116
E2	Estradiol (E2)	30	FOL	Folic Acid (Folate)	50
FSH	Follicle Stimulating Hormone (FSH)	30	VB12	Vitamin B12 (Cobalamin)	58
LH	Luteinizing Hormone (LH)	30	VDI	Vitamin D (25-OH), Total	70
PRL	Prolactin	30	HCG	β-hCG	58
PRG	Progesterone (P4)	30	ALDS	Aldosterone, Renin Concentration & Ratio	490
Tumour Markers					
AFP	Alpha-Fetoprotein	20	PSA	Prostate Specific Ag	20
CEA	Carcino-Embryonic Ag	20	PSAF	PSA, Free	70
FOB	Faecal Occult Blood	32	125	CA 125	20
PAP	PAP Smear	26	153	CA 15-3	20
PALP	Liquid Based PAP Test	54	199	CA 19-9	20
HPVD	HPV Genotyping	166	724	CA 72-4	80
HPVS	HPV DNA Screen (16, 18 & HR types)	138	SCC	Squamous Cell Ca Ag	110
CXC	HPV Genotyping & Liquid PAP	196	211	CYFRA 21-1	110
CXCQ	HPV DNA Screen & Liquid PAP	166	NSE	Neuron-Specific Enolase	80
EBVA	EBV VCA IgA	30			
EBA	EBV EA+NA1 IgA	120			

Code	Test Description	Specimen Type	TAT	RPP (RM)
GBS	General Chemistry Screen	5 ml Plain 2 ml Fluoride	1	46
	Liver Functions	13. Uric Acid		
	1. Bilirubin, Total	14. Sodium		
	2. Bilirubin, Direct & Indirect (if TBil elevated)	15. Potassium		
	3. Total Protein	16. Chloride		
	4. Albumin	17. Calcium		
	5. Globulin	18. Phosphorus		
	6. A/G Ratio	Lipids & Cardiovascular Risk		
	7. Alkaline Phosphates (ALP)	19. Cholesterol, Total		
	8. AST (SGOT)	20. HDL-Cholesterol		
	9. ALT (SGPT)	21. LDL-Cholesterol (direct method)		
	10. γ-Glutamyl Transpeptidase (GGT)	22. Triglycerides		
	Renal & Metabolic	23. Total / HDL-Cholesterol Ratio		
	11. Urea	Diabetes		
	12. Creatinine	24. Glucose		
FWM	Foreign Worker Screen – Male	5 ml Plain	2	66
	1. HIV 1&2 Ab+Ag	3 ml EDTA		
	2. HBs Ag	20 ml MSU		
	3. Syphilis RPR			
	4. Malaria Parasite			
	5. Morphine / Heroin Derivatives, by EIA			
	6. Cannabinoids, by EIA			
	7. Urine FEME			
	8. Blood Group			
FWF	Foreign Worker Screen – Female	5 ml Plain	2	70
	1. HIV 1&2 Ab+Ag	3 ml EDTA		
	2. HBs Ag	20 ml MSU		
	3. Syphilis RPR			
	4. Malaria Parasite			
	5. Morphine / Heroin Derivatives, by EIA			
	6. Cannabinoids, by EIA			
	7. Urine FEME			
	8. Blood Group			
	9. Urine Pregnancy Test			
FWM3	Foreign Worker Screen – Male (New)	5 ml Plain	2	116
	1. FWM	3 ml EDTA		
	2. Urine Amphetamine (Metamphetamine)	20 ml MSU		
	3. Anti-HCV			
	4. Filariasis			
FWF3	Foreign Worker Screen – Female (New)	5 ml Plain	2	120
	1. FWF	3 ml EDTA		
	2. Urine Amphetamine (Metamphetamine)	20 ml MSU		
	3. Anti-HCV			
	4. Filariasis			

Code	Test Description	Specimen Type	TAT	RPP (RM)
ISSB	International Student Screen	5 ml Plain	2	130
	1. HIV 1&2 Ab+Ag Screen	3 ml EDTA		
	2. HBs Ag	20 ml MSU		
	3. Anti-HBs			
	4. HCV Ab			
	5. Syphilis TP-Ab (reflex RPR if +ve)			
	6. Malaria Parasite			
	7. Amphetamine, by EIA			
	8. Morphine / Heroin Derivatives, by EIA			
	9. Cannabinoids, by EIA			
	10. Urine FEME			
PMSM	Pre-Marital Screen (Male)	5 ml Plain	4	200
PMSF	Pre-Marital Screen (Female)*	2x 3 ml EDTA		220
	1. Full Blood Count & Film Comment			
	2. ABO & Rh			
	3. Iron Status (Iron, Transferrin, TIBC & % Transferrin Saturation)			
	4. Hb Electrophoresis			
	5. TP Ab (reflex RPR if reactive)			
	6. HBs Ag & Anti-HBs			
	7. HCV Ab			
	8. HIV 1 & 2 Ag+Ab Screen			
	* 9. Rubella IgG (for Female only)			

Cardiovascular

Code	Test Description	Specimen Type	TAT	RPP (RM)
APLS	Antiphospholipid Syndrome Studies 1. Anti Beta-2 Glycoprotein IgG 2. Anti-Cardiolipin IgG 3. Anti-Phospholipid IgG	5 ml Plain	7-14	560
APLSS	Anti-Phospholipid Syndrome Studies 1. Anti-Beta-2 Glycoprotein IgG 2. Anti-Cardiolipin IgG 3. Anti-Phospholipid IgG 4. Lupus Anticoagulant (by LA-PTT & dRVVT)	5 ml Plain 2 x 2* ml Citrate <i>*as specified on tube</i> (freeze plasma) Stop heparin therapy one day prior to test	7-14	990
ATRA	Atherogenic Risk Profile – A 1. Total Cholesterol 2. HDL-Cholesterol 3. LDL-Cholesterol (direct method) 4. Triglycerides 5. Total/HDL-Cholesterol Ratio 6. High sensitive CRP (hs-CRP) 7. Homocysteine	5 ml Plain (refrigerated) (separate serum a.s.a.p.)	3	88
ATRB	Atherogenic Risk Profile – B 1. Urea 2. Uric Acid 3. Glucose 4. Total Cholesterol 5. HDL-Cholesterol 6. LDL-Cholesterol (direct method) 7. Triglycerides 8. Total/HDL-Cholesterol Ratio 9. High sensitive CRP (hs-CRP) 10. Homocysteine 11. Apolipoprotein-A1 12. Apolipoprotein-B 13. Apo-B/A1 Ratio	5 ml Plain 2 ml Fluoride (refrigerated) (separate serum a.s.a.p.)	5-7	180
ATRC	Atherogenic Risk Profile – C 1. Apolipoprotein-A1 2. Apolipoprotein-B 3. Apo-B/A1 Ratio 4. High sensitive CRP (hs-CRP) 5. Homocysteine 6. Lipoprotein (a)	5 ml Plain (refrigerated) (Separate serum a.s.a.p.)	5-7	220

Code	Test Description	Specimen Type	TAT	RPP (RM)
ATRD	Atherogenic Risk Profile – D 1. Total Cholesterol 2. HDL-Cholesterol 3. LDL-Cholesterol (direct method) 4. Triglycerides 5. Total/HDL-Cholesterol Ratio 6. High sensitive CRP (hs-CRP)	5 ml Plain	2	44
ATRF	Atherogenic Risk Profile – F 1. Total Cholesterol 2. HDL-Cholesterol 3. LDL-Cholesterol (direct method) 4. Triglycerides 5. Total/HDL-Cholesterol Ratio 6. High sensitive CRP (hs-CRP) 7. Troponin I, High Sensitive Quantitative (hsTnI)	5 ml Plain	3	140
CDMA	Cardiac Markers - A 1. AST (SGOT) 2. LDH 3. CK 4. High-Sensitive Troponin I (cTnI), CMIA	5 ml Plain	1	160
CDMC	Same as CDMA - No.4. High-Sensitive Troponin I (cTnI), FIA			
CDMB	Cardiac Markers - A 1. AST (SGOT) 2. LDH 3. CK 4. Troponin I (cTnI)		1	140
CRM	Cardiovascular Risk Markers 1. High sensitive CRP 2. Troponin I, High Sensitive Quantitative (hsTnI), CMIA 3. NT pro-BNP	5 ml Plain	5	450
LIP	Lipids Profile 1. Total Cholesterol 2. HDL-Cholesterol 3. LDL-Cholesterol (direct method) 4. Triglycerides 5. Total / HDL-Cholesterol Ratio	5 ml Plain	1	28
LEP	Lipoprotein Electrophoresis 1. Alpha Lipoproteins 2. Pre-Beta Lipoproteins 3. Beta Lipoproteins 4. Chylomicrons	5 ml Plain	5	140

Endocrinology & Metabolic

Code	Test Description	Specimen Type	TAT	RPP (RM)
ADRE	Adrenal Evaluation 1. Aldosterone 2. Cortisol 3. ACTH	5 ml Plain 3 ml EDTA (refrigerate plasma)	5	400
AGEA	Ageing Evaluation - A 1. IGF-1 2. DHEA-S 3. Cortisol 4. Estradiol (E2) 5. Total Testosterone 6. Insulin 7. TSH 8. FT4 9. FT3	5 ml Plain	5-7	660
AGEB	Ageing Evaluation - B 1. IGF-1 2. DHEA-S 3. Cortisol 4. Estradiol (E2) 5. Total Testosterone 6. SHBG 7. Free Androgen Index 8. Insulin 9. TSH 10. FT4 11. FT3	5 ml Plain	5-7	720
AGEC	Ageing Evaluation - C 1. IGF-1 2. DHEA-S 3. Cortisol 4. Estradiol (E2) 5. Total Testosterone 6. SHBG 7. Free Androgen Index 8. Insulin 9. TSH 10. FT4 11. FT3 12. ACTH 13. Prolactin 14. Beta-HCG	2 x 5 ml Plain 1 x 3ml EDTA (plasma)	5-7	980

Code	Test Description	Specimen Type	TAT	RPP (RM)
ALDS	Aldosteronism Studies 1. Aldosterone 2. Renin Concentration, Total 3. Aldosterone / Renin Ratio	5 ml Plain 3 ml EDTA (refrigerate plasma) (Specify patient's posture. Normal sodium intake & stop medication for 2 days)	7	620
BAM	Basal Metabolic Profile 1. Sodium 2. Potassium 3. Chloride 4. Calcium 5. Urea 6. Creatinine 7. Glucose	5 ml Plain 2 ml Fluoride (fasting / non-fasting)	1	34
BOM	Bone Markers 1. C-Telopeptide Crosslinks (CTX) 2. Osteocalcin	2x 3 ml EDTA Plasma (fasting, morning sample)	5-7	220
CAT	Catecholamine Fractions (by HPLC) 1. Epinephrine 2. Norepinephrine 3. Dopamine	24 hr Urine <i>in Acid preservative and Refrigeration</i> If possible, discontinue all medication prior to specimen collection	7-14	300
DDA	Diabetes Diagnostic - A 1. HbA1c 2. Glucose, fasting 3. Glucose, 2 hr postprandial/Glucose load	3 ml EDTA 2x 2 ml Fluoride - Fasting & 2 hr	2	48
DDB	Diabetes Diagnostic - B 1. HbA1c 2. Glucose 3. C-Peptide	3 ml EDTA 2 ml Fluoride 5 ml Plain	7	180
DDC	Diabetes Diagnostic - C 1. HbA1c 2. Glucose	3 ml EDTA 2 ml Fluoride	2	42
DFS	Diabetic Function Studies 1. Sodium 2. Potassium 3. Chloride 4. Calcium 5. Urea 6. Creatinine 7. Glucose 8. HbA1c 9. C-Peptide	3 ml EDTA 2 ml Fluoride 5 ml Plain	7	180

Code	Test Description	Specimen Type	TAT	RPP (RM)
DCP	Diabetic Care Profile	3 ml EDTA	2	78
	1. Glucose	2 ml Fluoride		
	2. HbA1c	5 ml Plain		
	3. Cholesterol, Total	20 ml Urine		
	4. HDL-Cholesterol			
	5. LDL-Cholesterol (direct method)			
	6. Triglycerides			
	7. Total / HDL-Cholesterol Ratio			
	8. C-Reactive Protein, high-sensitive			
	9. Urea			
	10. Creatinine			
	11. Uric Acid			
	12. Sodium			
	13. Potassium			
	14. Chloride			
	15. Urine Microalbumin, Creatinine & Ratio			
PRT	Parathyroid Evaluation	5 ml Plain	7	150
	1. Calcium			
	2. Phosphate			
	3. Urea			
	4. Creatinine			
	5. Alkaline Phosphatase			
	6. PTH-Intact			
PIT	Pituitary Evaluation	5 ml Plain	5-10	300
	1. FSH	(note time of collection and		
	2. LH	fasting/stimulation/suppression for GH)		
	3. Prolactin			
	4. TSH			
	5. Growth Hormone			
TASA	Thyroid Autoimmune Studies – A	5 ml Plain	7	250
	1. TSH (ultra-sensitive)			
	2. Free T4			
	3. Free T3			
	4. Anti-Thyroid Peroxidase (Anti-TPO)			
	5. Anti-Thyroglobulin Ab (ATAB)			
TASB	Thyroid Autoimmune Studies – B	5 ml Plain	7-10	380
	1. TSH (ultra-sensitive)			
	2. Free T4			
	3. Free T3			
	4. Anti-Thyroid Peroxidase (Anti-TPO)			
	5. Anti-Thyroglobulin Ab (ATAB)			
	6. TSH Receptor Ab			

Code	Test Description	Specimen Type	TAT	RPP (RM)
TFA	Thyroid Function Tests - A 1. TSH (ultra-sensitive) 2. Free T4	5 ml Plain	2	52
TFB	Thyroid Function Tests - B 1. TSH (ultra-sensitive) 2. Free T4 3. Free T3	5 ml Plain	2	84
AGECRF (Female)	Comprehensive Rejuvenation Profile Female / Male 1. Full Blood Count, Blood Film Comment	3 ml EDTA 2 x 5 ml Plain	7-10	1160
AGECRM (Male)	2. Lipids Profile, Liver Function, Renal Functions, 3. Calcium, Phosphate, Magnesium 4. Iron Status (Total Iron, % Transferrin & TIBC) 5. Hs-CRP 6. Homocysteine 7. Glucose, HbA1C 8. Insulin 9. Urine Microalbumin / Creatinine 10. Vitamin D 11. TSH 12. FT4 13. FT3 14. IGF-1 15. SHBG 16. Testosterone & Free Testosterone Index 17. Estradiol (E2) 18. FSH 19. LH 20. Progesterone 21. Prolactin 22. DHEA-S 23. Cortisol 24. Hep A Ab 25. HBs Ag & Ab 26. HIV 1 & 2 Ag+Ab 27. TP Ab (reflex RPR if +ve) 28. H. Pylori Ab 29. AFP 30. CEA 31. CA19-9 32. CA125 (for Female) 33. CA15-3 (for Female) 34. PSA (for Male) 35. Urine FEME	2 ml Fluoride 20 ml MSU		

Gynaecology & Reproductive

Code	Test Description	Specimen Type	TAT	RPP (RM)
AME	Amenorrhoea Studies - A 1. TSH (ultra-sensitive) 2. Prolactin 3. Follicle Stimulating Hormone (FSH) 4. Luteinising Hormone (LH) 5. Testosterone, Total	5 ml Plain	5	166
AMEB	Amenorrhoea Studies - B 1. TSH (ultra-sensitive) 2. Prolactin 3. Follicle Stimulating Hormone (FSH) 4. Luteinising Hormone (LH) 5. Estradiol (E2)	5 ml Plain	5	166
AHE	Androgen Hormons Evaluation 1. Androstenedione (ASD) 2. Testosterone, Total 3. Testosterone, Free 4. Dihydrotestosterone (DHT) 5. Sex Hormone Binding Globulin (SHBG)	5 ml Plain	10-14	940
ATB	Antenatal Screen – B 1. Full Blood Count 2. Blood Group 3. Syphilis TP-Ab (reflex RPR titre if TP-Ab +) 4. HBs Ag 5. Anti-HBs 6. HIV 1&2 Ab+Ag Screen	3 ml EDTA 5 ml Plain	2	60
ATC	Antenatal Screen – C 1. Full Blood Count 2. Blood Group 3. Syphilis TP-Ab (reflex RPR titre if TP-Ab +) 4. HBs Ag 5. Anti-HBs 6. HIV 1&2 Ab+Ag Screen 7. HCV Ab 8. Rubella IgG	3 ml EDTA 5 ml Plain	2	110

Code	Test Description	Specimen Type	TAT	RPP (RM)
ATD	Antenatal Screen – D 1. Full Blood Count 2. Blood Group 3. Syphilis TP-Ab (reflex RPR titre if TP-Ab +) 4. HBs Ag 5. Anti-HBs 6. HIV 1&2 Ab+Ag Screen 7. Rubella IgG	3 ml EDTA 5 ml Plain	2	80
ATE	Antenatal Screen – E 1. Full Blood Count 2. Blood Group 3. Syphilis TP-Ab (reflex RPR titre if TP-Ab +) 4. HIV 1&2 Ab+Ag Screen	3 ml EDTA 5 ml Plain	2	48
ATF	Antenatal Screen – F 1. Full Blood Count 2. Blood Group 3. Syphilis TP-Ab (reflex RPR titre if TP-Ab +) 4. HBs Ag 5. Anti-HBs	3 ml EDTA 5 ml Plain	2	48
ATG	Antenatal Screen – G 1. Full Blood Count 2. Blood Group 3. Glucose 4. Syphilis TP-Ab (reflex RPR titre if TP-Ab +) 5. HBs Ag 6. Anti-HBs 7. HIV 1&2 Ab+Ag Screen 8. Rubella IgG	3 ml EDTA 5 ml Plain 3 ml Fluoride	2	86
ATJ	Antenatal Screen – J 1. Full Blood Count 2. Blood Group 3. TSH 4. Syphilis TP-Ab (reflex RPR titre if TP-Ab +) 5. HBs Ag 6. Anti-HBs 7. HIV 1&2 Ab+Ag Screen 8. Rubella IgG	3 ml EDTA 5 ml Plain	2	100
CXC	Cervical Cancer Risk - A 1. HPV DNA Genotyping (All types) 2. Liquid-based PAP Test	Cervical brush in Collection vial	7-10	280

Code	Test Description	Specimen Type	TAT	RPP (RM)
CXCQ	Cervical Cancer Risk - B 1. HPV DNA Screen – 16, 18 & other 13 high risk types 2. Liquid-based PAP Test	Cervical brush in Collection vial	5-7	200
HIR	Hirsutism Studies 1. DHEA-S 2. Testosterone, Total (consider adding 17-OH-Progestrone)	5 ml Plain	5	120
IFFA	Fertility Hormonal Studies – Female A 1. Prolactin 2. FSH 3. LH 4. Progesterone (P4) 5. Estradiol (E2)	5 ml Plain	3	176
IFFB	Fertility Hormonal Studies – Female B 1. FSH 2. Progesterone (P4) 3. Prolactin	5 ml Plain	3	104
IFFC	Fertility Hormonal Studies – Female C 1. FSH 2. LH 3. Prolactin	5 ml Plain	3	104
IFFD	Fertility Hormonal Studies – Female D 1. FSH 2. LH 3. Progesterone (P4) 4. Estradiol (E2) 5. Anti-Mullerian Hormone (AMH)	5 ml Plain	7	440
IFM	Fertility Hormonal Studies – Male 1. Prolactin 2. FSH 3. LH 4. Total Testosterone	5 ml Plain	3	140
MEHA	Menopausal Hormone Studies – A 1. FSH 2. LH 3. Estradiol (E ₂) 4. Prolactin 5. TSH (ultra-sensitive)	5 ml Plain	3	166

Code	Test Description	Specimen Type	TAT	RPP (RM)
MEHB	Menopausal Hormone Studies – B 1. FSH 2. LH 3. Estradiol (E ₂)	5 ml Plain	3	104
OFE	Ovarian Function Evaluation 1. FSH 2. LH 3. LH/FSH Ratio	5 ml Plain	5	72
MCS	Menstrual Cycle Studies 1. FSH 2. LH 3. Estradiol (E ₂) 4. Progesterone (P ₄)	5 ml Plain	5	140

Prenatal & Hereditary Genetic Assay:		<i>*Use Prenatal & Genetic Test Requisition Form</i>		
CYTOOA	Chromosomal Microarray Analysis (CMA) ~ CytoOneArray (Prenatal / Postnatal) Detect 327 chromosomal abnormalities	Prenatal: 15ml Amniotic Fluid Postnatal: 2x 3 ml EDTA Blood /Cord Blood	7-14	5600
FE1P	Foetal Screen- 1st Trimester Test 1. PAPP-A 2. βhCG, Free 3. NT measurement (by physician) 4. Risk Profiling	5 ml plain Taken between 11-13 wks of gestation	12-14	480
FE4P	Foetal Screen- 2nd Trimester Quadruple Test 1. AFP 2. βhCG, Total 3. uE3 4. Inhibin A 5. Risk Profiling	5 ml Plain Taken between 15-20 wks of gestation	12-14	590

Code	Test Description	Specimen Type	TAT	RPP (RM)
FEPRE	Pre-Eclampsia Risk at 1st Trimester 1. PAPP-A 2. PIGF 3. Risk profiling for Pre-eclampsia & IUGR	5 ml Plain Taken between 11-13 wks of gestation	12-14	700
FE1PR	Pre-Eclampsia Risk & Foetal Screen for 1st Trimester 1. β hCG, free 2. PAPP-A 3. PIGF 4. Risk Profiling for Trisomy 21, 18, 13, Pre-Eclampsia & IUGR	5 ml Plain Taken between 11-13 wks of gestation	12-14	900
FEPR2	Pre-Eclampsia Risk at 2nd & Late Trimester 1. sFlt-1 2. PAPP-A 3. Risk Profiling for Pre-Eclampsia	5 ml Plain Taken between 15-20 wks of gestation	12-14	1200
GHRA	Genetic Hereditary Risk Profile - A 1. Fragile X Syndrome (FXS) Genetic Assay 2. Spinal Muscular Atrophy (SMA) Genetic Assay 3. Huntington's Disease Genetic Assay 4. Methylenetetrahydrofolate reductase (MTHFR) Genetic Assay	2x 3 ml EDTA Blood	7-14	2400

Haematology & Coagulation

Code	Test Description	Specimen Type	TAT	RPP (RM)
AEA	Anaemia Studies – A	3 ml EDTA	3	90
	1. Full Blood Picture	5 ml Plain		
	2. Serum Iron			
	3. Transferrin			
	4. TIBC			
	5. % Transferrin Saturation			
	6. Ferritin			
AEB	Anaemia Studies – B	3 ml EDTA	5-7	180
	1. Full Blood Picture	5 ml Plain		
	2. Serum Iron			
	3. Transferrin			
	4. TIBC			
	5. % Transferrin Saturation			
	6. Ferritin			
	7. Vitamin B12			
AEC	Anaemia Studies – C	3 ml EDTA	5-7	120
	1. Full Blood Picture	5 ml Plain		
	2. Vitamin B12			
	3. Folic Acid			
IRO	Iron Status Evaluation	5 ml Plain	5	80
	1. Serum Iron			
	2. TIBC			
	3. Transferrin			
	4. % Transferrin Saturation			
FBC	Full Blood Count	3 ml EDTA	1	26
	1. Haemoglobin (Hb)			
	2. Red Blood Count (RBC)			
	3. Haematocrit (Hct/PCV)			
	4. Mean Corpuscular Volume (MCV)			
	5. Mean Corpuscular Hb (MCH)			
	6. Mean Corpuscular Hb Conc. (MCHC)			
	7. Red Cell Distribution Width (RDW)			
	8. White Blood Cell Count (WBC)			
	9. White Blood Cell Differential Count (5 parts)			
FBCC	Full Blood Count with CRP (titter)	3 ml EDTA	1	34
	1. Full Blood Count	2 ml Plain		
	2. High-Sensitive C-Reactive Protein			

Code	Test Description	Specimen Type	TAT	RPP (RM)
FBPC	Full Blood Picture with CRP (titter) 1. Full Blood Count 2. High-Sensitive C-Reactive Protein 3. Blood Film Comment*	3 ml EDTA 2 ml Plain	1	42
FBP	Full Blood Picture with ESR 1. Full Blood Count 2. Erythrocyte Sedimentation Rate (ESR) 3. Blood Film Comment*	3 ml EDTA (min. 2 ml)	1	36
BFCH	<i>* If Blood film refer to & comment by Haematologist</i>		3-5	+120
THLA	Thalassaemia Studies – A 1. Full Blood Picture 2. Serum Iron 3. Transferrin 4. TIBC 5. % Iron Saturation 6. Haemoglobin Electrophoresis 7. Comment and Interpretation by Consultant Haematologist	2x 3 ml EDTA 5 ml Plain	5	188
THLB	Thalassaemia Studies – B 1. Full Blood Picture 2. Serum Iron 3. Transferrin 4. TIBC 5. % Iron Saturation 6. Ferritin 7. Haemoglobin Electrophoresis 8. Comment and Interpretation by Haematologist	2x 3 ml EDTA 5 ml Plain	5	218
THLD	Thalassaemia DNA Analysis (both Alpha & Beta) (for couples of suspected carrier if MCV <80 & MCH <25) 1. Hb Electrophoresis 2. Serum Ferritin 3. Genetic detection of abnormalities for both alpha & beta-thalassaemia	2x 3 ml EDTA (Cord Blood for neonate) 5 ml Plain <i>*Use Prenatal & Genetic Test Requisition Form</i>	7-14	1800
THLDA	Thalassaemia DNA Analysis, Alpha (7 mutations)	2x 3 ml EDTA (Cord Blood for neonate)	7-14	900
THLDB	Thalassaemia DNA Analysis, Beta (22 mutations)	2x 3 ml EDTA (Cord Blood for neonate)	7-14	1300
COA	Coagulation Studies 1. Platelet Count 2. Prothrombin Time (PT) 3. Activated Partial Thromboplastin Time (APTT)	2x 2 ml Citrate (1:9) 3 ml EDTA	2	100
TBPS	Thrombophilia Screen 1. Anti-Thrombin III 2. Protein C 3. Protein S	2x 2 ml Citrate (1:9) (double spin, freeze plasma)	14-21	520

Hepatobiliary & Gastrointestinal

Code	Test Description	Specimen Type	TAT	RPP (RM)
ALDA	Autoimmune Liver-Disease (ALD) Antibody Panel Included Antibodies: AMA-M2 (Anti-Mitochondrial Antibody) LKM-1 (Liver Kidney Microsomal type 1) LC-1 (Liver Cytosol type 1) SLA/LP (Soluble Liver Antigen/Liver Pancreas)	5 ml Plain	14-16	1180
COLS	Colitis Studies 1. Stool Culture 2. <i>C. difficile</i> Culture & Toxins 3. ANCA & pattern 4. Faecal Occult Blood	Fresh Stool 5 ml Plain	5-10	520
DIAR	Diarrhoea Studies 1. Stool Culture 2. Faecal Ova & Parasite 3. Rotavirus Ag	Fresh Stool	3-5	180
DYSP	Dyspepsia Studies 1. Full Blood Count 2. <i>H. pylori</i> IgG 3. Amylase 4. Liver Function Tests (may consider adding Gastrin)	3 ml EDTA 5 ml Plain	3	120
HABC	Hepatitis ABC Screen 1. Anti-HAV (HAV IgG) 2. HBs Ag 3. Anti-HBs 2. Anti-HCV	5 ml Plain	2	130
HBC	Hepatitis B Carrier Studies – A 1. AST 2. ALT 3. AFP 4. HBe Ag 5. Anti-HBe	5 ml Plain	4	110
HBCB	Hepatitis B Carrier Studies – B 1. AST 2. ALT 3. AFP 4. HBs Ag, Quantitative 5. HBe Ag 6. Anti-HBe	5 ml Plain	5	220

Code	Test Description	Specimen Type	TAT	RPP (RM)
HBS	Hepatitis B Screening 1. HBs Ag 2. Anti-HBs	5 ml Plain	2	26
HBT	Hepatitis B Treatment Monitoring 1. AST 2. ALT 3. AFP 4. HBe Ag 5. Anti-HBe 6. HBV DNA Viral Load	5 ml Plain 3 ml EDTA (Plasma)	7-10	510
HCT	Hepatitis C Treatment Monitoring 1. AST 2. ALT 3. AFP 4. HCV-RNA Viral Load	5 ml Plain 3 ml EDTA (Plasma)	7-14	630
HIS	Hepatitis Immune Status Profile 1. HBs Ag 2. Anti-HBs 3. Anti-HAV (HAV IgG)	5 ml Plain	2	70
LFT	Liver Function Test 1. Bilirubin, Total 2. Bilirubin, Direct & Indirect (if T.Bil elevated) 3. Total Protein 4. Albumin 5. Globulin 6. A/G Ratio 7. Alkaline Phosphates (ALP) 8. AST (SGOT) 9. ALT (SGPT) 10. γ-Glutamyl Transpeptidase (GGT)	5 ml Plain	1	34
MALS	Malabsorption Studies 1. Full Blood Count 2. Ferritin 3. Folic Acid, Serum 4. Vitamin B12 5. Calcium 6. Liver Function Tests	3 ml EDTA 5 ml Plain	7	240

Code	Test Description	Specimen Type	TAT	RPP (RM)
PEP	Protein Electrophoresis, Serum	5 ml Plain	5	140
UPEP	Protein Electrophoresis, Urine	20 ml FMU	5	200
	1. Total Protein			
	2. Albumin			
	3. Alpha 1 Globulin			
	4. Alpha 2 Globulin			
	5. Beta Globulin			
	6. Gamma Globulin			
	(consider Immunofixation Ep if paraprotein is suspected)			
VHD	Viral Hepatitis Diagnostic	5 ml Plain	2-5	240
	1. Liver Function Tests			
	2. HAV IgM			
	3. HBs Ag			
	4. HBc IgM (If HBs Ag Positive)			
	5. Anti-HCV			

Infectious Diseases

Code	Test Description	Specimen Type	TAT	RPP (RM)
DEN1	Dengue Antigen & FBC (recommend if fever within 1-5 days) 1. Full Blood Count 2. Dengue NS1 Antigen	3 ml EDTA 5 ml Plain	1	120
DEN2	Dengue Duo Tests (recommend if fever within 1-14 days) 1. Full Blood Count 2. Dengue NS1 Antigen 3. Dengue IgG & IgM	3 ml EDTA 5 ml Plain	1	180
DEN3	Dengue IgG/IgM & FBC (recommend if fever within 5-14 days) 1. Full Blood Count 2. Dengue IgG & IgM, rapid test	3 ml EDTA 5 ml Plain	1	90
FEBA	Febrile Studies - A 1. Full Blood Count & Blood Film 2. Hs-CRP 3. Malarial Parasites 4. Widal Test (Typhoid) 5. Weil Felix Test (Typhus) 6. Monospot (Infectious Mononucleosis) 7. Urine FEME	3 ml EDTA 5 ml Plain 20 ml Urine	1	84
FEBB	Febrile Studies – B 1. Full Blood Count & Blood Film 2. Hs-CRP 2. Malarial Parasites 3. Widal Test (Typhoid) 4. Weil Felix Test (Typhus) 5. Monospot (Infectious Mononucleosis) 6. Urine FEME 7. Dengue IgG & IgM, rapid	3 ml EDTA 5 ml Plain 20 ml Urine	1	150
HIVM	HIV Monitoring 1. CD4 & CD8 Count 2. HIV RNA Viral Load (by RT-PCR)	3x 3 ml EDTA Call Lab	7-14	1000
STDA	STD Studies –A (asymptomatic screen) 1. Syphilis TP-Ab (reflex RPR if Pos) 2. HIV 1&2 Ab+Ag Screen 3. HSV type 1 IgG 4. HSV type 2 IgG 5. <i>Chlamydia trachomatis</i> DNA 6. <i>Neisseria gonorrhoeae</i> DNA	5 ml Plain Cervical Swab (Female) Urethral Swab (Male)	2-5	350

Code	Test Description	Specimen Type	TAT	RPP (RM)
STDB	STD Studies - B <i>(with lesions, lymph nodes enlargement)</i> 1. Syphilis TP-Ab (reflex RPR if Pos) 2. <i>Treponema pallidum</i> DNA 3. Herpes simplex virus DNA type 1 & 2	5 ml Plain Swab/ Scraping from lesions	2-5	380
STDC	STD Studies - C <i>(with discharge, dysuria)</i> 1. Gram's Stain for Gonococcus/Candida 2. Wet Mount for Trichomonas 3. Swab C & S, Genital 4. <i>Chlamydia trachomatis</i> DNA 5. <i>Neisseria gonorrhoeae</i> DNA	2x Genital swab in medium 1x Genital swab, plain (Female) 1x Urethral Swab (Male)	2-5	320
STDD	STD Studies – D (Female Only) <i>(suspect pelvic inflammatory disease)</i> 1. C-Reactive Protein, quantitative 2. <i>Chlamydia trachomatis</i> DNA 3. <i>Neisseria gonorrhoeae</i> DNA 4. Swab C & S, Cervical	5 ml Plain 1x Cervical swab in medium 1x Cervical swab, plain	1-5	300
STDE	STD Studies – E (common screening) 1. Syphilis TP-Ab (reflex RPR if Pos) 3. HIV 1&2 Ab+Ag Screen 4. Hepatitis Bs Ag 5. Hepatitis Bs Ab	5 ml Plain	2	66
STDG	STD Studies - G 1. Syphilis TP-Ab (reflex RPR if Pos) 3. HIV 1&2 Ab+Ag Screen 4. HSV type 1 IgG 5. HSV type 2 IgG 6. <i>Chlamydia</i> IgG	5 ml Plain	2-5	200
STPA	STD PCR Profile – A 1. Human papillomavirus (HPV) DNA Screen (type 16, 18 & HR) 2. <i>Chlamydia trachomatis</i> DNA 3. <i>Neisseria gonorrhoeae</i> DNA 4. Herpes simplex type 1 & 2 DNA	Genital Swab (Female) Urethral Swab (Male)	7-14	460

Code	Test Description	Specimen Type	TAT	RPP (RM)
STPB	STD PCR Profile – B 1. Herpes simplex type 1 & 2 DNA 2. <i>Chlamydia trachomatis</i> DNA 3. <i>Neisseria gonorrhoeae</i> DNA 4. <i>Treponema pallidum</i> DNA with 5. Syphilis TP-Ab (reflex RPR if Pos)	5 ml Plain Genital Swab (Female) Urethral Swab (Male)	5-7	480
STPC	STD PCR Profile – C 1. <i>Chlamydia trachomatis</i> DNA 2. <i>Neisseria gonorrhoeae</i> DNA 3. <i>Trichomonas vaginalis</i> DNA 4. <i>Mycoplasma hominus</i> DNA 5. <i>Mycoplasma genitalium</i> DNA 6. <i>Ureaplasma urealyticum</i> DNA 7. <i>Ureaplasma parvum</i> DNA	Genital Swab (Female) Urethral Swab (Male)	7	340
STPD	STD PCR Profile – D 1. Human papillomavirus (HPV) DNA Screen (type 16, 18 & HR) 2. <i>Chlamydia trachomatis</i> DNA 3. <i>Neisseria gonorrhoeae</i> DNA 4. Herpes simplex type 1 & 2 DNA 5. <i>Trichomonas vaginalis</i> DNA 6. <i>Candida albicans</i> DNA 7. <i>Treponema pallidum</i> DNA with 8. Syphilis TP-Ab (reflex RPR if +ve)	5 ml Plain Genital Swab (Female) Urethral Swab (Male)	7	660
TOR	TORCH Infection 1. Toxoplasma IgM 2. Rubella IgM 3. CMV IgM 4. HSV type 1 IgM 5. HSV type 1 IgM 6. TP-Ab (reflex RPR if Pos)	5 ml Plain	3-5	420
TOS	TORCH Screen 1. Toxoplasma IgG 2. Rubella IgG 3. CMV IgG 4. HSV type 1 IgG 5. HSV type 2 IgG 6. TP-Ab (reflex RPR if Pos)	5 ml Plain	3	220

<i>Paediatric</i>				
Code	Test Description	Specimen Type	TAT	RPP (RM)
NNJA	Neonatal Jaundice Studies A 1. Bilirubin, total 2. Bilirubin, direct & indirect	1 ml Plain (paediatric vial) (avoid light)	1	24
NNJB	Neonatal Jaundice Studies B 1. Bilirubin, total 2. Bilirubin, direct & indirect 3. Full Blood Count & Film Comment 4. Coomb's Test, direct	1 ml Plain (paediatric vial) (avoid light) 1 ml EDTA (paediatric vial)	1	70
NNA	Neonatal Screen A 1. ABO & Rh Blood Group 2. G6PD Screen 3. TSH	3 ml EDTA 5 ml Plain (Neonate's blood or Cord blood)	1	56
NNB	Neonatal Screen B 1. ABO & Rh Blood Group 2. G6PD Screen 3. TSH 4. Syphilis TP-Ab (reflex RPR if Pos)	3 ml EDTA 5 ml Plain (Neonate's blood or Cord blood)	1	66
NNC	Neonatal Screen C 1. ABO & Rh Blood Group 2. G6PD Screen 3. TSH 4. Neonatal IgE (Low range) for Allergy Risk	3 ml EDTA 5 ml Plain (Neonate's blood or Cord blood)	5	150
NHS	Neonatal Health Screen 1. ABO & Rh Blood Group 2. G6PD (Quantitative) 3. TSH 4. Neonatal IgE (Low range) for Allergy Risk 5. Amino Acid Analysis (22 types) for Inborn Error of Metabolism	3 ml EDTA 5 ml Plain (Neonate's blood or Cord blood)	7-10	640

Renal & Urinary

Code	Test Description	Specimen Type	TAT	RPP (RM)
DIM1	Dialysis Indices Monitor – 1	3 ml EDTA	2	52
	1. Full Blood Count	5 ml Plain (Pre-Dialysis)		
	2. hs-CRP	3 ml Plain (Post-Dialysis)		
	3. Liver Function Test	3 ml Fluoride		
	4. Renal Function Test (Creatinine Enzymatic method) with Pre & Post Urea			
	5. Calcium w corrected Ca			
	6. Phosphorus			
	7. Glucose			
DIM2	Dialysis Indices Monitor - 2	3 ml EDTA	2	170
	1. Full Blood Count	5 ml Plain (Pre-Dialysis)		
	2. hs-CRP	3 ml Plain (Post-Dialysis)		
	3. Liver Function Test	3 ml Fluoride		
	4. Renal Function Test (Creatinine Enzymatic method) with Pre & Post Urea			
	5. Calcium w corrected Ca			
	6. Phosphorus			
	7. Glucose			
	8. Iron Status (Iron, Transferrin, % Saturation & TIBC)			
	10. Ferritin			
	11. Treponema pallidum Ab (TP-Ab) (reflex RPR titter)			
	12. Hep. B Ag & Ab			
	13. Hep. C Ab			
	14. HIV 1&2 Ag+Ab			
DIM3	Dialysis Indices Monitor - 3	3 ml EDTA X 2	5	270
	As DIM2 plus	5 ml Plain (Pre-Dialysis)		
	15. I-PTH	3 ml Plain (Post-Dialysis)		
		3 ml Fluoride		
DIM4	Dialysis Indices Monitor - 4	Same as above	5	300
	As DIM3 plus			
	16. HbA1c			
	Tag-on for DIM Profiles			
ABORH	ABO & RH Group		1	8
IRST	Iron Status (Iron, Transferrin , % & TIBC)		2	32
FER	Ferritin		3	36
A1C	HbA1c		2	30
GALB	Glycated Albumin		3	56
IPTH	i-PTH		5	100
EPO	Erythropoietin		7	320
ALU	Serum Aluminum	3 ml Trace Element Tube Heparin	7-10	180
DIVA	Dialysis Virology Screen - A	5 ml Plain	2	100
	1. Hepatitis Bs Ag & Ab			
	3. HCV Ab			
	4. HIV 1&2 Ab+Ag Screen			
	5. Treponema pallidum Ab (TP-Ab) (reflex RPR titter)			

Code	Test Description	Specimen Type	TAT	RPP (RM)
GNS	Glomerulonephritis Serology 1. Anti-Neutrophil Cytoplasmic Ab (ANCA) with pattern 2. Glomerular Basement Membrane Ab (Anti-GBM) 3. Anti-Nuclear Ab (ANA) with pattern 4. Immunoglobulin A (IgA) 6. Complement C3 7. ASOT, Quantitative	5 ml Plain	7-14	640
RFT	Renal Function Test 1. Urea 2. Creatinine w eGFR 3. Uric Acid 4. Sodium 5. Potassium 6. Chloride	5 ml Plain	1	32
RFS	Renal Function Studies 1. Urea 2. Creatinine w eGFR 3. Uric Acid 4. Sodium 5. Potassium 6. Chloride 7. Albumin 8. Phosphorus 9. Calcium with corrected Ca 10. Urine FEME	5 ml Plain 20 ml MSU	1	44
GFRCY	Estimated Glomerular Filtration Rate (with Cystatin C) 1. Serum Creatinine 2. Serum Cystatin C 3. eGFR cr-cys (with CKD-EPI 2012 calculation)	5 ml Plain (Provide Age, Race & Gender)	5	160

Rheumatology, Immunology & Allergy

Code	Test Description	Specimen Type	TAT	RPP (RM)
ARSA	Arthritis Screen - A 1. Full Blood Count 2. ESR 3. C-Reactive Protein, high sensitive 4. Rheumatoid Factor 5. Uric Acid 6. Anti-Nuclear Abs (CLIA Quantitative) 7. ASOT (latex agglutination)	3 ml EDTA 5 ml Plain	3	ARSA: 140
ARSAF	Note: Substitute No.6 ANAQ with ANA by IFA		10	ARSAF: 170
ARSB	Arthritis Studies - B 1. Full Blood Count 2. ESR 3. C-Reactive Protein, high sensitive 4. Calcium 5. Phosphorous 6. Uric Acid 7. Creatine Kinase (CK) 8. Rheumatoid Factor 9. Anti-CCP 10. Anti-Nuclear Abs (CLIA Quantitative) 11. ASOT (latex agglutination) 12. TP-Ab (reflex RPR if Pos)	3 ml EDTA 5 ml Plain	3	ARSB: 230
ARSBF	Note: Substitute No.10 ANAQ with ANA by IFA		10	ARSBF: 260
(Recommend adding Synovial Fluid FEME/C&S for Gout/Infection, HLA-B27 for Ankylosing Spondylitis and Reactive Arthritis)				
IGEV	Immunoglobulins Evaluation 1. IgA 2. IgG 3. IgM 4. IgD 5. IgE (may consider Immunofixation Electrophoresis)	5 ml Plain	7	330

Code	Test Description	Specimen Type	TAT	RPP (RM)
IFE	Immunofixation Electrophoresis (Serum)	5 ml Plain	10-14	360
IFEU	Immunofixation Electrophoresis (Urine/CSF)	50 ml FMU / 24hr Urine or 10 ml CSF	10-14	440
	1. IgA			
	2. IgG			
	3. IgM			
	4. Kappa Light Chain			
	5. Lambda Light Chain			
	6. Interpretation			
LUS	Lupus Screening	5 ml Plain	7	190
	1. Full Blood Count	3 ml EDTA		
	2. ESR			
	3. C-Reactive Protein, high sensitive			
	4. Anti-Nuclear Abs (CLIA Quantitative)			
	5. Anti-dsDNA			
LUSF	Note: Substitute No.6 ANAQ with ANA by IFA			LUSF: 220
LES	Lupus Erythematous Studies	5 ml Plain	7	370
	1. Full Blood Count	20 ml MSU		
	2. ESR	3 ml EDTA		
	3. C-Reactive Protein, high sensitive			
	4. Albumin			
	5. Urea			
	6. Creatinine			
	7. Creatine Kinase (CK)			
	8. Coombs Test, Direct			
	10. Anti-Nuclear Abs (CLIA Quantitative)			
	11. Anti-dsDNA			
	12. Complement C3			
	13. Complement C4			
	14. Urine FEME			
	(may consider adding Anti-ENA)			
LESF	Note: Substitute No.6 ANAQ with ANA by IFA			LESF: 400
MCG	Monoclonal Gammopathy Studies	5 ml Plain	5	190
	1. Serum Protein Electrophoresis	20 ml FMU		
	2. Bence-Jones Protein (agglutination)			
	3. Beta-2 Microglobulin, Serum			
	(may consider Serum or Urine Immunofixation Electrophoresis IFE)			
RHES	Rheumatoid Screen	5 ml Plain	3-5	160
	1. Rheumatoid Factor, Quantitative			
	2. Anti-CCP			

Code	Test Description	Specimen Type	TAT	RPP (RM)
IGE	Total IgE For allergy screening, but low sensitivity and specificity for atopic allergy.	5 ml Plain	5	76
IGEC	Total IgE Low Range (Neonatal) Low range detection, for allergy risk in neonate.	3 ml EDTA Cord Blood (Plasma)	5	110
ECP	Eosinophil Cationic Protein For asthma monitoring	5 ml Plain	5	260
PHAD	Phadiatop (Screening) To differentiate between atopic and non-atopic patients. Results indicate probability for atopy. A negative result indicates that the symptoms are not caused by common environmental allergens, and the physician may explore other possibilities. Sensitivity: 93%, Specificity: 89%	5 ml Plain	5-7	180
MIX PENAL				
fx2	fx2 Sea Food Mix Fish, shrimp, blue mussel, tuna, salmon	5 ml Plain	7	180
fx3	fx3 Grain Mix Wheat, Oat, Maize, Sesame seed, Buckwheat	5 ml Plain	7	180
fx5	fx5 Common Food Mix Egg white, milk, fish, wheat, peanut, soya bean	5 ml Plain	7	180
fx7	fx7 Vegetable Mix Tomato, Yeast, Garlic, Onion, Celery	5 ml Plain	7	180
fx30	fx30 Fruit Mix Kiwi, Mango, Banana, Avocado, Papaya	5 ml Plain	7	280
hx2	hx2 House Dust Mix House dust (Hollister-Stier Labs), <i>D. pteronyssinus</i> , <i>D. farina</i> , <i>Batella germanica</i>	5 ml Plain	7	180
ex2	ex2 Animal Epidermal Mix Cat dander, Dog dander, Guinea pig epithelium, Rat, Mouse	5 ml Plain	7	220
mx2	mx2 Microorganisms Mix <i>Penicillium notatum</i> , <i>Cladosporium herbarum</i> , <i>Aspergillus fumigates</i> , <i>Candida albicans</i> , <i>Alternaria tenuis</i> , <i>Helminthosporium halodes</i>	5 ml Plain	7	180

Code	Test Description	Specimen Type	TAT	RPP (RM)
Specific IgE for Individual Allergen:				
Single Allergen – Select code & allergen below		5 ml Plain	7-14*	90
Select from any allergen below. *TAT may be delayed depends on availability of reagent stock. Call Lab for availability on the list below or any other allergens.				
Code	Allergens	Code	Allergens	
MITES & HOUSE DUST		MICROORGANISMS		
d1	<i>D. pteronyssinus</i> (House dust mite)	m1	<i>Penicillium notatum</i>	
d2	<i>D. farinae</i> (House dust mite)	m2	<i>Cladosporium herbarum</i>	
d201	<i>Blomia tropicalis</i> (Storage mite)	m3	<i>Aspergillus fumigatus</i>	
h1	House Dust (Greer Labs)	m5	<i>Candida albicans</i>	
		m6	<i>Alternaria tenuis</i>	
INSECTS		POLLENS		
i6	Cockroach	w1	Common ragweed	
EPIDERMAL & ANIMAL PROTEINS		g2	Bermuda grass	
e1	Cat dander			
e5	Dog dander			
e85	Chicken feathers			
FOODS				
f1	Egg White	f27	Beef	
f2	Cow Milk	f33	Orange	
f3	Fish (Cod)	f45	Yeast	
f4	Wheat	f47	Garlic	
f7	Oat	f48	Onion	
f8	Maize, corn	f49	Apple	
f10	Sesame seed	f75	Egg yolk	
f11	Buckwheat	f83	Chicken	
f13	Peanut	f84	Kiwi	
f14	Soybean	f85	Celery	
f17	Hazel nut	f87	Melon	
f18	Brazil nut	f92	Banana	
f20	Almond	f93	Cacao (Cocoa)	
f23	Crab	f259	Grape	
f24	Shrimp	f284	Turkey meat	
f25	Tomato			
f26	Pork			

Allergy and Food Intolerance Packages

v	AE64 (IgE)	AE107 (IgE) including AE64 (except in*)	AG08 (IgG)	AG60 (IgG)	AG90 (IgG)	
1	Total IgE	Alder	Gluten	Almond	Cow's Milk	Asparagus
2	House Dust	Birch	Soy bean	Hazelnut	Goat's Milk	Bamboo Shoots
3	D. Pteronyssinus	Oak	Cow milk	Pine nut	Cheddar Cheese	Broccoli
4	D. Farinae	Japanese hop	Peanut	Sesame	Yogurt	Cabbage
5	Cat dander	Mugwort	Egg	Cacao	Egg White	Carrot
6	Dog dander	Cladosporium Herbarum	Beef	Sweet chestnut	Egg Yolk	Cauliflower
7	Cockroach	Crab	Cod Fish	Pea	Abalone	Celery
8	Rye	Mackerel	Shrimp	Soy Bean	Cod	Cucumber
9	Common Ragweed	Sweet Vernal/ Orchard grass/ Common reed/ Bent grass		Peanut	Crab	Eggplant
10	Timothy Grass	Bermuda grass		Walnut	Clam	Green onion
11	Alternaria Alternata	Willow		Pistachio nut	Cuttlefish	Leek
12	Aspergillus fumigatus	Cotton wood		Sunflower	Eel	Lettuce
13	P. Notatum	White pine		Corn	Kelp	Onion
14	Candida Albicans	Japanese cedar		Rye	Oyster	Peas
15	Olive	Ox-eye daisy		Buckwheat	Salmon	Peppers
16	Bee Venom	Dandelion		Barley	Sharks Fin	Potato
17	Horse	Plantain		Rice	Shrimp	Spinach
18	Mouse / Rat	Russian Thistle		Wheat	Tuna	String Bean
19	Rabbit	Golden rod		Eel	Buckwheat	Sweet Potato
20	Wool	Common pig weed		Codfish	Gluten/Wheat Bran	Taro
21	Acarus Siro	Latex		Mackerel	Corn	Tomato
22	T. Putrescentiae	Wasp venom		Salmon	Rice	Mushroom
23	Acacia	Guniea pig		Tuna	Oat	Chilli Pepper
24	White ash	Hamster		Plaice	Wheat	Garlic
25	CCD	Horse		Anchovy	Adlay	Ginger
26	Yeast's baker	Tuna/ Salmon		Trout	Kidney Bean	Pepper
27	Almond	Plaice/ Anchovy/ Alaska pollock		Oyster	Mung Bean	Yeast
28	Hazelnut	Lobster/ Pacific squid		Scallop	Soya Bean	Honey
29	Coconut *	Blue mussel / Oyster/ Clam/ Scallop		Clam	Sunflower seed	Vanilla
30	Egg white	Silkworm Pupa		Blue mussel	Almond	Coffee
31	Egg yolk *	Buckwheat		Loster	Walnut	Tea
32	Milk	Citrus mix		Pacific squid	Sesame Seed	
33	Cheese	Sweet chestnut		Crab	Cocoa bean	
34	Peanut	Almond/ Pine nut/ Sunflower		Shrimp	Peanut	
35	Soybean	Cocoa		Edd yolk	Apple	
36	Wheat			Edd white	Banana	
37	Shrimp			Milk	Grape	
38	Lobster			Lamb Meat	Grapefruit	
39	Codfish			Beef	Guava	
40	Barley			Chicken	Kiwi Fruit	
41	Rice			Pork	Lemon	
42	Pea *			Carrot	Longans	
43	Sesame			Celery	Mango	
44	Apple			Mushroom	Olive	
45	Mango / Banana			Potato	Orange	
46	Peach			Cucumber	Papaya	
47	Strawberry			Garlic	Peach	
48	Tomato			Onion	Pear	
49	Corn			Tomato	Pineapple	
50	Carrot			Eggplant	Cherry	
51	Potato			Apple	Strawberry	
52	Garlic / Onion			Banana	Watermelon	
53	Pumpkin *			Coconut	Coconut	
54	Eggplant *			Grape	Avocado	
55	Cucumber			Peach	Beef	
56	Orange *			Strawberry	Chicken	
57	Pork			Orange	Lamb	
58	Beef			Mango	Pork	
59	Chicken			Kiwi	Duck	
60	Lamb meat			Yeast, baker's		

Profile Code	RPP (RM)
AE64	540
AE107	640
AG08	180
AG90	860

Profile Combination		RPP (RM)
AEG72	AE64+AG08	580
AEG115	AE107+AG08	680
AEG154	AE64+AG90	1300
AEG197	AE107+AG90	1400

Sample Requirement:

2 months min of age
3 ml plain tube

TAT: 3-5 working days

<i>Tumour Markers</i>				
Code	Test Description	Specimen Type	TAT	RPP (RM)
TMF	Tumour Markers For Female (General) 1. AFP 2. CEA 3. CA 125 4. CA 15-3 5. CA 19-9 6. CA 72-4 7. EBV VCA IgA	5 ml Plain	5	420
TMM	Tumour Markers For Male (General) 1. AFP 2. CEA 3. PSA 4. CA19-9 5. CA 72-4 6. EBV VCA IgA	5 ml Plain	5	360
TMBR	Tumour Markers - Breast 1. CA 15-3 2. CEA 3. Tissue Polypeptide Ag (TPA)	5 ml Plain	10	240
TMCX	Tumour Markers - Cervical 1. SCC Ag 2. CEA (consider CXC Cervical Cancer Risk profile for screening)	5 ml Plain	5	180
CXC	Cervical Cancer Risk - A 1. HPV DNA genotyping (All genotypes) 2. Liquid-based PAP Test	Cervical brush in Collection vial	7-10	280
CXCQ	Cervical Cancer Risk - B 1. HPV DNA Screen – 16, 18 & other high risk types 2. Liquid-based PAP Test	Cervical brush in Collection vial	5-7	200
TMCP	Tumour Markers - Colorectal 1. CEA 2. CA 19-9 3. CA 72-4 4. Faecal Occult Blood (human-Hb specific)	5 ml Plain Fresh Stool	5	260
TMLI	Tumour Markers - Liver 1. AFP 2. CEA	5 ml Plain	2	76

Code	Test Description	Specimen Type	TAT	RPP (RM)
TMLS	Tumour Markers - Lung (Small Cell) 1. NSE 2. CEA 3. Calcitonin	5 ml Plain	10	460
TMLN	Tumour Markers - Lung (Non-Small Cell) & Oesophagus 1. CYFRA 21-1 2. SCC Ag	5 ml Plain	5	260
TML	Tumour Marker For Lung 1. CYFRA 21-1 2. NSE 3. SCC Ag	5 ml Plain	5	340
TMNE	Tumour Markers - Neuroblastoma 1. Catecholamine Fractions 2. Homovanillic Acid (HVA) 3. Vanillylmandelic Acid (VMA) 4. Neuron-Specific Enolase (NSE)	24 hr Urine <i>in Acid preservative and Refrigeration</i> 5 ml Plain	7-14	840
TMNP	Tumour Markers - Nasopharyngeal 1. EBV VCA IgA 2. EBV EB+NA1 IgA	5 ml Plain	5-7	160
TMOV	Tumour Markers - Ovary 1. CA 125 2. CEA 3. CA 72-4	5 ml Plain	7	220
ROMA	Risk of Ovarian Malignancy Algorithm (ROMA) 1. Human Epididymis Protein 4 (HE4) 2. CA 125 3. Risk Calculation	5 ml Plain	7-10	500
TMPR	Tumour Markers - Prostate 1. PSA, total 2. PSA, free 3. Total/Free PSA Ratio 4. Acid Phosphatase, Prostatic	5 ml Plain	5	170
PHI	Prostate Health Index 1. PSA, total 2. PSA, free 3. % Free PSA 4. [-2]proPSA Isoform (p2PSA) & PHI Risk	5 ml Plain	5	580

Code	Test Description	Specimen Type	TAT	RPP (RM)
TMST	Tumour Markers – Stomach & Pancreas 1. CA 72-4 2. CEA 3. CA 19-9	5 ml Plain	5	220
TMTE	Tumour Markers - Testis 1. AFP 2. β -hCG	5 ml Plain	5	80
TMTH	Tumour Markers - Thyroid 1. Thyroglobulin 2. Calcitonin 3. CEA	5 ml Plain	10	440

Section C

SINGLE TESTS

Single Tests – Alphabetical Index

Code	Test Description	Specimen Type	TAT	RPP (RM)
17OH	17-Hydroxycorticosteroids (17-OHCS)	24hr Urine, in Acid preservative or Refrigeration	10	260
17AO	17-Hydroxyprogesterone (17-OHP)	5 ml Plain, Separate serum & freeze. Specify the age, sex, and phase of the cycle. For women, the sample must be taken at the start of the follicular phase.	10	260
17KS	17-Ketosteroids (17-KS)	24hr Urine, in Acid preservative or Refrigeration	10	260
5HI	5-Hydroxyindoleacetic Acid (5-HIAA)	24hr Urine, in Acid preservative or Refrigeration	17	260
A				
ACH	Acetylcholine Receptor Ab (AchR Ab)	5 ml Plain, Separate serum and freeze	7-14	400
	Acid Fast Bacilli (AFB) Culture & Sensitivity (refer to Culture)			
AFBC	Acid Fast Bacilli (AFB) Concentrated (Membrane-base Smear)	Sputum/ Pleural Fluid/ FMU/ CSF/ Swab Significant higher sensitivity than direct smear.	2	100
AFBS	Acid Fast Bacilli (AFB) ZN Stain (Direct Smear)	Sputum/ Pleural Fluid/ FMU/ CSF/ Swab Collect fresh sputum recommended on three consecutive days or early morning urine or pleural fluid, CSF, swab.	2	28
ACPP	Acid Phosphatase, Prostatic	5 ml Plain	7	80
ACP	Acid Phosphatase, Total	5 ml Plain	5	36
APTT	Activated Partial Thrombin Time (APTT)	2 ml Citrate If possible, please specify if the patient is on an anticoagulation treatment as well as the clinical context.	2	50
D208	Additive Drugs (188 types) Qualitative Screening in Food/Medication (by HPLC)	Call Lab	14	Call
ADA	Adenosin Deaminase (ADA), pleural fluid	Pleural Fluid	10-14	400
ADENO	Adenovirus DNA (by PCR)	Discharge/ Aspirate/ Swab	7	350
ACTH	Adrenocorticotrophic Hormone (ACTH)	3 ml EDTA (Plasma) Not to be performed on haemolysed sample. Sample on a cooled tube at 4°C, centrifuge a.s.a.p and freeze the plasma.	7	140
ALT	Alanine Aminotransferase (ALT/SGPT)	5 ml Plain	1	10
ALB	Albumin	5 ml Plain	1	10
FALD	Aldosterone, Urine	24hr Urine, in Acid preservative or Refrigeration	7	400
ALD	Aldosterone, Serum	5 ml Plain Do not refer haemolysed or lipaemic samples. Specify upright or reclined (Sampling in upright position after 1 hr walking; sampling in reclined position after 1 hr in supine position).	7	300
ALKP	Alkaline Phosphatase (ALP)	5 ml Plain	1	10
ALPB	Alkaline Phosphatase - Bone (BAP)	5 ml Plain, Separate serum and freeze	5-7	180
ALPE	Alkaline Phosphatase Isoenzymes Electrophoresis - Liver, Bone, Intestine, Fast Liver	5 ml Plain	7-14	200
	Allergy Tests (refer to Allergy Section in Profile Tests)			
A1A	Alpha-1 Antitrypsin	5 ml Plain	7	110
A1AS	Alpha-1 Antitrypsin, Stool	Fresh Stool	15	240
AFP	Alpha-Fetoprotein (AFP)	5 ml Plain	2	36

Code	Test Description	Specimen Type	TAT	RPP (RM)
AMMO	Ammonia	3 ml EDTA (Plasma)	7	90
AMOE	Amoebic Antibody (<i>E histolytica</i>)	5 ml Plain	7	160
AMY	Amylase	5 ml Plain	2	20
UAM	Amylase (Diastase), Urine	20 ml Urine	7	30
ASD	Androstenedione (ASD)	5 ml Plain, <i>Separate serum and freeze</i>	15	280
Antibody Screen (refer to Coombs' Test, Indirect)				
ABMZ	Anti-Basement Membrane Zone Abs (Anti-BMZ)	5 ml Pain	15	140
ABGPG	Anti Beta-2 Glycoprotein-I IgG	5 ml Plain	7-10	240
ABGPM	Anti Beta-2 Glycoprotein-I IgM	5 ml Plain	7-10	240
ACAG	Anti-Cardiolipin IgG (ACA IgG) (a type of Antiphospholipid Ab)	5 ml Plain	8	160
ACAM	Anti-Cardiolipin IgM (ACA IgM) (a type of Antiphospholipid Ab)	5 ml Plain	8	160
ACCP	Anti-Cyclic Citrullinated Peptide Ab (Anti-CCP)	5 ml Plain <i>Interference possible in patients treated with biotin (vitamin B7, B8 or H) or taking any food supplement containing biotin. Essential to stop treatment 8 days before taking the sample.</i>	5	160
ADNA	Anti-double stranded DNA (Anti-dsDNA)	5 ml Plain	7	140
ENAT	Anti-Extractable Nuclear Antigen (Anti-ENA), Total (Quatitative)	5 ml Plain	7	180
ENA6	Anti-ENA, 6 types: RNP Ab, Sm Ab, SS-A Ab, SS-B Ab, Scl-70 Ab & Jo-1 Ab	5 ml Plain	7-10	540
RNP SM SSA SSB SCL70 JO1	Single Antibody of Anti-ENA: - RNP Ab - Sm Ab - SS-A Ab - SS-B Ab - Scl-70 Ab - Jo-1 Ab	5 ml Plain	7-10	280 each
GPCA	Anti-Gastric Parietal Cell Abs (APCA)	5 ml Plain	7-14	140
GBM	Anti-Glomerular Basement Membrane Abs (Anti-GBM)	5 ml Plain	7-14	200
INTAB	Anti-Intercellular Substance Abs	5 ml Plain	7-14	160
Anti-Microsomal Ab (AMA) (see Anti-TPO)				
AMIA	Anti-Mitochondrial Ab	5 ml Plain	12	140
AMH	Anti-Mullerian Hormone (AMH)	5 ml Plain	5-7	340
ANCA	Anti-Neutrophil Cytoplasmic Ab (ANCA) w pattern (by IFA method)	5 ml Plain	12	200
ANAQ	Anti-Nuclear Abs (Qunatitative)	5 ml Plain	5	78
ANA	Anti-Nuclear Abs (ANA) w pattern (by IFA)	5 ml Plain	5	100
APSG	Anti-Phosphatidylserine IgG	5 ml Plain	7-10	290
APLA	Anti-Platelet Antibody	5 ml Plain	7-10	520
ASMA	Anti-Smooth Muscle Ab (ASMA)	5 ml Plain	12	140

Code	Test Description	Specimen Type	TAT	RPP (RM)
ASA	Anti-Sperm Ab, Circulating	5 ml Plain	7-10	260
ASOT	Anti-Streptolysin O Titre (ASOT), Spot test	5 ml Plain	2	44
AT3	Anti-Thrombin III	2x 2 ml Citrate <i>Always specific the treatment and clinical context of the test prescription: Unfractionated heparin may induce a weak decrease in antitrombin level.</i>	15-20	160
ATAB	Anti-Thyroglobulin Ab (ATA)	5 ml Plain <i>Interference possible in patients treated with biotin (vitamin B7, B8 or H) or taking any food supplement containing biotin. Essential to stop treatment 8 days before taking the sample.</i>	5	90
ATPO	Anti-Thyroid Peroxidase (Anti-TPO)	5 ml Plain <i>Interference possible in patients treated with biotin (vitamin B7, B8 or H) or taking any food supplement containing biotin. Essential to stop treatment 8 days before taking the sample.</i>	5	90
APOA	Apolipoprotein A1 (Apo-A1)	5 ml Plain	3	60
APOB	Apolipoprotein B (Apo-B)	5 ml Plain	3	60
APOP	Apo- A1, Apo-B & Ratio	5 ml Plain <i>Fasting sample.</i>	3	100
APOEG	Apolipoprotein E genotyping (by PCR)	3 ml EDTA	7-14	560
AST	Aspartate Aminotransferase (AST/SGOT)	5 ml Plain	1	10
	Aspirate Culture & Sensitivity (refer to Culture)			
B				
BJP	Bence-Jones Protein (by heat test) (consider Immunofixation Ep)	20 ml FMU, <i>Refrigeration, without preservative and acid</i>	5	56
B2GG	Beta-2 Glycoprotein 1 IgG	5 ml Plain	7	240
B2GM	Beta-2 Glycoprotein 1 IgM	5 ml Plain	10	240
B2M	Beta-2 Microglobulin, Serum	5 ml Plain, <i>Always indicated the beginning or end of dialysis</i>	5	100
UB2M	Beta-2-Microglobulin, Urine	20 ml FMU	5	100
HCG	Beta-Human Chorionic Gonanotropin, Total (β-hCG)	5 ml Plain	2	58
HCGF	β-hCG, Free	5 ml Pain	7	120
BID	Bilirubin, Direct/Conjugated *	5 ml Plain, <i>Wrap or Store away from light.</i>	1	10
BIT	Bilirubin, Total *	5 ml Plain, <i>Wrap or Store away from light.</i>	1	10
	Blood Culture & Sensitivity (refer to Culture)			
	Blood Film Comment (refer Full Blood Picture)			
ABORH	Blood Grouping & Rhesus (ABO & Rh)	3 ml EDTA	1	16
BRCA	BRCA 1 & 2 Gene Mutation Analysis Genetic risk marker for breast & ovarian cancer	2x 3 ml EDTA <i>Attach the current clinical and therapeutic data</i>	14	4000
BNPF	B-type Natriuretic Peptide (NT pro-BNP)	3 ml EDTA (Plasma)	5	360

Code	Test Description	Specimen Type	TAT	RPP (RM)
C				
125	CA 125 (Ovarian cancer)	5 ml Plain	2	80
153	CA 15-3 (Breast cancer)	5 ml Plain	2	80
199	CA 19-9 (Pancreatic, gastric cancer)	5 ml Plain <i>Interference possible in patients treated with biotin (vitamin B7, B8 or H) or taking any food supplement containing biotin. Essential to stop treatment 8 days before taking the sample.</i>	2	80
724	CA 72-4 (TAG-72) (Gastrointestinal, Breast cancer)	5 ml Plain <i>Interference possible in patients treated with biotin (vitamin B7, B8 or H) or taking any food supplement containing biotin. Essential to stop treatment 8 days before taking the sample.</i>	5	140
CAC	Calcitonin	5 ml Plain <i>Collect sample away from meal in order to avoid hypergastrinemy. In case of proton pumps treatment, recommend to collect sample after intake. Or please collect sample 2 weeks after treatment stopped. Separate serum & freeze.</i>	6-8	380
CA	Calcium	5 ml Plain	1	10
CAF	Calcium, Free (Ionized)	5 ml Heparin (whole blood)	7-14	170
UCA	Calcium, Urine	24 hr Urine, in Acid preservative and Refrigeration	5	20
CAD	Candida albicans DNA (by PCR)	Genital Swab / FVU	7	160
CEA	Carcino-Embryonic Antigen (CEA)	5 ml Plain <i>Interference possible in patients treated with biotin (vitamin B7, B8 or H) or taking any food supplement containing biotin. Essential to stop treatment 8 days before taking the sample.</i>	2	50
	Carcinoid Syndrome (refer to 5-HIAA)			
CARNI	Carnitine, Free & Total	5 ml Plain	14-20	760
CD48	CD4 / CD8 Count	3 ml EDTA Call Lab	5	540
CER	Ceruloplasmin	5 ml Plain	5	120
CHIM	Chikungunya IgM	5 ml Plain	2	120
CLPM	Chlamydia pneumoniae IgM	5 ml Plain	7-10	160
	Chlamydia Ag (refer to Chlamydia DNA PCR)			
CLA	Chlamydia trachomatis IgA	5 ml Plain	5	140
CLG	Chlamydia trachomatis IgG	5 ml Plain	5	90
CLM	Chlamydia trachomatis IgM	5 ml Plain	5	140
CTNG	Chlamydia trachomatis & Neisseria gonorrhoeae DNA (by PCR)	Genital Swab / FVU	5	250
	Chloride (Cl) (refer to Electrolytes)			
	Chloride, Urine (refer to Electrolytes)			
CHO	Cholesterol, Total	5 ml Plain	1	10
CGA	Chromogranin A (CgA)	5 ml Plain <i>Centrifuge without delay, separate the supernatant and freeze. PPIs should be stopped 2-3 weeks before test.</i>	7-10	560

Code	Test Description	Specimen Type	TAT	RPP (RM)
CYTOOA	Chromosomal Microarray Analysis (CMA) ~ CytoOneArray (Prenatal / Postnatal) Detect 327 chromosomal abnormalities	Prenatal: 15ml Amniotic Fluid /CVS Postnatal: 2x 3 ml EDTA Blood / Cord Blood	7-14	5600
CHAA	Chromosome Analysis (Karyotype), Amniotic Fluid	15 ml Amniotic Fluid (<i>room temperature</i>)	7	2600
CDPCR	<i>Clostridium difficile</i> DNA (by PCR)	Fresh Stool in steril container	3	700
HCO2	CO2 Total / Bicarbonate	5 ml Plain	5	20
CF9	Coagulation Factor IX	2x 2 ml Citrate	14	300
CF8	Coagulation Factor VIII	2x 2 ml Citrate	14	300
COLA	Cold Agglutinin	5 ml Plain, <i>Keep warm before serum separation.</i>	5	90
C3	Complement 3 (C3)	5 ml Plain	5	60
C4	Complement 4 (C4)	5 ml Plain	5	60
COOD	Coombs' test, Direct	3 ml EDTA	5	40
COOI	Coombs' test, Indirect	3 ml EDTA	5	60
Coronavirus Infectious Disease 2019 (COVID-19) (refer to SARS-CoV-2)				
UFC	Cortisol, Free, Urine	24 hr Urine, <i>in Acid preservative and Refrigeration</i>	10	100
CORSA	Cortisol, Free, Saliva	3 ml Saliva	7	76
CORT	Cortisol, Serum	5 ml Plain, <i>Sample collect between 7am and 10am, or according to the instruction from prescribing party. Interference possible in patients treated with biotin (vitamin B7, B8 or H) or taking any food supplement containing biotin. Essential to stop treatment 8 days before taking the sample.</i>	5	70
CPE	C-Peptide	5 ml Plain	8	170
HCRP	C-Reactive Protein, high-sensitive (hsCRP)	5 ml Plain	2	44
CK	Creatine Kinase (CK)	5 ml Plain	1	20
CKE	Creatine Kinase (CK) Isoenzyme Electrophoresis (CK-Total, MM, MB, BB)	5 ml Plain	7	160
CRE	Creatinine	5 ml Plain	1	10
CCT	Creatinine Clearance Test & Rate	5 ml Plain & 24hr Urine	2	40
UCRE	Creatinine, Urine	20 ml Urine	2	14
UCRE24	Creatinine, 24 hrs Urine	24 hrs Urine, <i>in refrigeration</i>	2	18
CYFG	Cryofibrinogen (Qualitative)	2x 2 ml Citrate (fasting) <i>Keep warm 37°C before plasma separation.</i>	5-10	80
CYGLO	Cryoglobulins (Qualitative)	5 ml Plain (fasting) <i>Keep warm 37°C before serum separation.</i>	10-14	80
CRYA	<i>Cryptococcus</i> Antigen	1 ml CSF / 5 ml Plain	3-5	200
CSF Culture & Sensitivity (refer to Culture)				
CFEM	CSF FEME - Cell count, Gram's stain, India ink stain, glucose & protein	Min 2 ml CSF	2	80
CTX	C-Telopeptide Crosslinks (CTx) (beta-CrossLaps)	3 ml EDTA Plasma (Fasting Morning) <i>Interference possible in patients treated with biotin (vitamin B7, B8 or H) or taking any food supplement containing biotin. Essential to stop treatment 8 days before taking the sample.</i>	5-7	136

Code	Test Description	Specimen Type	TAT	RPP (RM)
211	CYFRA 21-1 (Cytokeratin 19) (non-small cell lung cancer)	5 ml Plain <i>Interference possible in patients treated with biotin (vitamin B7, B8 or H) or taking any food supplement containing biotin. Essential to stop treatment 8 days before taking the sample.</i>	5	150
CYSC	Cystatin C	5 ml Plain	5	180
CMVD	Cytomegalovirus (CMV) DNA Detection	Swab/ FVU /CSF/ Serum	7	340
CMVL	Cytomegalovirus (CMV) DNA Viral Load	5 ml Plain		560
CMVG	Cytomegalovirus (CMV) IgG	5 ml Plain	3-5	80
CMVM	Cytomegalovirus (CMV) IgM	5 ml Plain	3-5	110

Culture & Sensitivity ~ Culture, Microscopy & Sensitivity tests

CAFB	AFB/Mycobacterium Culture (include Identification & 1st line Drug Sensitivity tests if positive) (may consider Mycobacterium DNA PCR assay)	Sputum/ FMU/ CSF/ Swab	6-8 Wks	500
CASP	C&S Aspirate	Aspirate in sterile-capped syringe/tube/ bottle (specify source of specimen)	3-5	70
CASC	~ C&S Aspirate w Chemistry (Protein, Glu & LDH)	As above	3-5	130
CASJ	~ C&S Aspirate w Chemistry (Protein, Glu & RF)	As above	3-5	130
CBAE	~ C&S Blood, Aerobic (w/o supply of Broth)	Blood/ Pus/ Fluid in blood culture broth (aerobic)	10-12	130
CBAN	~ C&S Blood, Anaerobic (w/o supply of Broth)	Blood/ Pus/ Fluid in blood culture broth (anaerobic)	10-12	130
CCSF	~ C&S CSF	CSF in sterile-capped syringe/tube	4-7	60
CCSC	~ C&S CSF w Chemistry (Pro, Glu)	As above	4-7	120
CSTO	~ C&S Stool for <i>Salmonella</i>, <i>Shigella</i> & <i>Vibrio cholera</i> only	Fresh Stool/Rectal Swab	3-5	64
CSPU	~ C&S Sputum (non-Mycobacterium only)	Sputum (wash mouth, avoid saliva)	3-5	60
CSWA	~ C&S Swab	Swab in transport medium (specify source of specimen)	3-5	50
CURI	~ C&S Urine w Chemistry	20 ml MSU	3-5	50

Culture Only (without Sensitivity tests)

CSSH	~ Stool for <i>Salmonella</i> & <i>Shigella</i> only	Fresh Stool/Rectal Swab	3-5	44
CSAU	~ Stool for <i>Staphylococcus aureus</i> only	Fresh Stool/Rectal Swab	Call	44
CSVC	~ Stool for <i>Vibrio cholera</i> only	Fresh Stool/Rectal Swab	3-5	44
	~ Stool for <i>Clostridium difficile</i> Culture & Toxins (refer <i>Clostridium difficile</i> PCR test)			
CFUN	~ Fungus/ Yeast w Microscopy	Relevant specimen, Skin Scraps	14-21	60

Code	Test Description	Specimen Type	TAT	RPP (RM)
D				
DDI	D-Dimer (Quantitative) (may also consider FDP)	2 ml Citrate (1:9)	5	180
DHEAS	Dehydroepiandrosterone Sulfate (DHEA-S)	5 ml Plain <i>Interference possible in patients treated with biotin (vitamin B7, B8 or H) or taking any food supplement containing biotin. Essential to stop treatment 8 days before taking the sample.</i>	7	90
DENGM	Dengue Serology (IgG & IgM), Rapid Test	5 ml Plain	1	80
DENAG	Dengue NS1 Antigen (intended for 1st -5th days)	5ml plain	1	100
DHT	Dihydrotestosterone (DHT)	5 ml Plain <i>Separate serum & freeze.</i>	7-10	340
	Dilantin (refer to Phenytoin in Toxicology section)			
DNAF	DNA Profiling for Paternity (for non-legal proceeding only)	3 ml EDTA <i>Specimen each from child, mother & alleged father. Use Specific Form</i>	14	4600 Call
DNA1	~ Add each additional child			1600
	Down Syndrome Screen (refer to Foetal Screen in Profile Test)			
E				
ELT	Electrolytes (Na, K & Cl), Serum	5 ml Plain	1	24
UFLT	Electrolytes (Na, K & Cl), Urine	24 hr Urine, <i>in Refrigeration without acid preservative</i>	1	50
ELAL	Endotoxin, LAL Assay	Water/ Dialysate	7	560
	Entamoeba histolytica Ab (refer to Amoebic Ab)			
ECP	Eosinophil Cationic Protein (ECP)	5 ml Plain	5	260
EBVD	Epstein-Barr virus (EBV) DNA Detection	Serum/ Swab/ Fluid	7	300
EBA	EBV EA+NA1 IgA (for NPC)	5 ml Pain	5	150
EBEG	EBV Early Ag (EA) IgG (for IM/NPC)	5 ml Plain	5	180
EBVA	EBV Viral Capsid Ag (VCA) IgA (for NPC)	5 ml Plain	3	100
EBVG	EBV Viral Capsid Ag (VCA) IgG (for IM)	5 ml Plain	8	110
EBVM	EBV Viral Capsid Ag (VCA) IgM (for IM)	5 ml Plain	8	150
ESR	Erythrocyte Sedimentation Rate (ESR)	3 ml EDTA	1	10
EPO	Erythropoietin (EPO)	5 ml Plain, <i>Sample in morning (variations during day). Separate serum & freeze.</i>	7	320
E2	Estradiol (E2) (Ovarian Function)	5 ml Plain	3	44
E3U	Estriol, Free (uE3) (Pregnancy Monitoring)	5 ml Plain	7	110
F				
FOB	Faecal Occult Blood (human-Hb specific)	Fresh Stool	2	40
	Faeces Culture & Sensitivity (refer to Culture)			

Code	Test Description	Specimen Type	TAT	RPP (RM)
FFEM	Faeces FEME- <i>for Amoeba/Ova/Cyst</i>	Fresh Stool	2	20
FER	Ferritin	5 ml Plain	5	50
FHB	Foetal Haemoglobin (APT Test)	Stool/ Vomitus	5	70
	Foetal Abnormalities Risk Screening 1st & 2nd Trimester (refer to Profile Test Page 18)			
FDP	Fibrin Degradation Products (FDP) (may also consider D-Dimer)	2x 2 ml Citrate	5	200
FIB	Fibrinogen	2x 2 ml Citrate	5	70
FOLR	Folic Acid (Folate), RBC	3 ml EDTA, <i>wrap and avoid light</i>	Research, call	
FOL	Folic Acid (Folate), Serum	5 ml Plain, <i>Fasting sample.</i>	5	60
FSH	Follicle Stimulating Hormone (FSH)	5 ml Plain	3	44
FXSG	Fragile X Syndrome (FXS) Genetic Assay	3 ml EDTA		800
FRUC	Fructosamine	5 ml Plain	Research, call	
FBC	Full Blood Count- Hb, RBC indices, PLT, TWDC	3 ml EDTA	1	26
FBCC	Full Blood Count with CRP (titter)	3 ml EDTA	1	34
FBP	Full Blood Picture with ESR - FBC, Blood Film Comment & ESR	3 ml EDTA	1	36
FBPC	Full Blood Picture with CRP - FBC, Blood Film Comment & CRP	3 ml EDTA 5 ml Plain	1	42
FBPH	Full Blood Picture with Blood Film Comment by Haematologist	3 ml EDTA	3	156
	Fungus Culture (refer to Culture)			
FUNG	Fungus Microscopic	Skin Scraps	2	24

G

G6PD	G6PD Screening	Blotted blood on paper or 3 ml EDTA	2	18
G6PQ	G6PD, Quantitative	3 ml EDTA	3-5	70
GGT	gamma-Glutamyl Transferase (GGT)	5 ml Plain	1	10
GAS	Gastrin	5 ml Plain, <i>Fasting minimum 10 – 12 hours. After removing the coagulated mass, quickly centrifuge the sample to separate the serum & freeze.</i>	7	260
GLU	Glucose, Fasting	2 ml Fluoride, <i>Fasting</i>	1	10
GTT2	Glucose Tolerance Test (2 Points) (w/o Glucose supply)	2x 2 ml Fluoride <i>Fasting & 2 hours Post-Glucose</i>	1	20
GTT3	Glucose Tolerance Test (3 Points) (w/o Glucose supply)	3x 2 ml Fluoride <i>Fasting, 1 & 2 hours Post-Glucose</i>	1	30
75GL	~ 75g Glucose Load supply			10 / pack
GLU2	Glucose, 2 hrs Post-Glucose Load	2x 2 ml Fluoride, <i>Fasting & 2 hours Post-Glucose</i>	1	10
GAD	Glutamic Acid Decarboxylase AutoAbs (GAD AutoAbs)	5 ml Plain	15-20	400
	Glutathione Peroxidase			Research, Call

Code	Test Description	Specimen Type	TAT	RPP (RM)
GALB	Glycated Albumin (GA) (diabetic monitoring)	5 ml Plain	3-5	80
	Glycated Haemoglobin (refer to HbA1c)			
GRAM	Gram's Stain Examination	Swab / Smear / Urine	1	20
HGH	Growth Hormone (hGH)	5 ml Plain, <i>Separate serum & freeze.</i> <i>Note time of collection & fasting / stimulation / suppression.</i>	7	220
H	Haemoglobin (Hb) (refer to Full Blood Count)			
A1C	Haemoglobin A1c (HbA1c)	3 ml EDTA	1	40
HBE	Haemoglobin Electrophoresis + FBC + Comment and Interpretation by Haematologist	3 ml EDTA	7	168
HAPG	Haptoglobulin	5 ml Plain	5	110
HDEP	HDL, LDL, VLDL Electrophoresis	5 ml Plain	5	120
HDLC	HDL-Cholesterol (Direct Method)	5 ml Plain	1	10
HPYG	<i>H. pylori</i> IgG (semi-quantitative)	5 ml Plain	2	76
HPYA	<i>H. pylori</i> Abs (with Current Infection Marker)	5 ml Plain	2	90
UBT4	<i>H. pylori</i> ¹⁴C-Urea Breath Test	Breath in card <i>Resting patient, fasting for a minimum of 6 hours (no drink, no food, no smoke). No antibiotic treatment for 4 weeks. No Proton-pump inhibitor treatment for 2 weeks. No antacids and gastro-intestinal protectants 24 hours before the test.</i>	3	160
HAV	Hepatitis A IgG (Anti-HAV)	5 ml Plain	1	44
AVM	Hepatitis A IgM (HAV IgM)	5 ml Plain	5	70
BCG	Hepatitis B core Abs (Anti-HBc)	5 ml Plain	5	70
BCM	Hepatitis B core IgM (HBc IgM)	5 ml Plain	5	110
BCAG	Hepatitis B core-related Antigen (HBcr Ag), Quantitative	5 ml Plain	7-10	640
EAB	Hepatitis B envelope Ab (Anti-HBe)	5 ml Plain	3	64
EAG	Hepatitis B envelope Ag (Hbe-Ag)	5 ml Plain	3	64
SAB	Hepatitis B surface Ab (Anti-HBs)	5 ml Plain	1	18
SAG	Hepatitis B surface Ag (HBs-Ag)	5 ml Plain	1	18
SAGQ	Hepatitis B surface Ag, Quantitative	5 ml Plain	5	120
HBVPC	HBV Pre-core (PreC) Mutant	5 ml EDTA (Plasma), <i>Separate plasma & freeze</i>	7-14	600
HBVPS	HBV Pre-surface (PreS) Mutant	5 ml EDTA (Plasma), <i>Separate plasma & freeze</i>	7-14	600
HBVY	HBV Drug Resistance Analysis	5 ml EDTA (Plasma), <i>Separate plasma & freeze.</i> <i>Please specify the date and result of the last viral load. The minimum viral load is 100 IU.</i>	7-14	760
HBVG	HBV-DNA Genotyping	5 ml EDTA (Plasma), <i>Separate plasma & freeze.</i> <i>Please specify the date and result of the last viral load. The minimum viral load is 100 IU.</i>	7-14	700
HBVL	HBV-DNA Viral load (by qPCR)	5 ml EDTA (Plasma), <i>Separate plasma & refrigerate</i>	7-10	400
HCV	Hepatitis C Ab (Anti-HCV)	5 ml Plain	2	90
HCVG	HCV-RNA Genotyping	3 ml EDTA (Plasma), <i>Separate plasma & freeze.</i> <i>Please specify the date and result of the last viral load. The minimum viral load is 500 IU.</i>	7-14	700
HCVR	HCV-RNA Viral load (qPCR)	3 ml EDTA (Plasma), <i>Separate plasma & refrigerate</i>	7-10	580

Code	Test Description	Specimen Type	TAT	RPP (RM)
H1G	Herpes Simplex Virus (HSV) Type 1 IgG	5 ml Plain	5	70
H2G	HSV Type 2 IgG	5 ml Plain	5	70
H12M	HSV Type 1 & 2 IgM	5 ml Plain	5	240
HSVD	HSV DNA type 1 & 2 DNA	Swab/ Urine/ CSF	7-14	300
HCY	Homocysteine, total (tHcy)	3 ml Plain, <i>Fasting, refrigerated, separate serum a.s.a.p.</i>	3	90
HVA	Homovanillic Acid (HVA)	24 hr Urine, <i>in Acid preservative and Refrigeration</i>	5-7	360
HVAR	Homovanillic Acid/ Creatinine Ratio	20 ml Urine	5-7	360
HIV	Human Immunodeficiency Virus (HIV) 1&2 Ab+Ag Screen (HIV 1&2 Ab+p24 Ag)	5 ml Plain <i>Interference possible in patients treated with biotin (vitamin B7, B8 or H) or taking any food supplement containing biotin. Essential to stop treatment 8 days before taking the sample.</i>	2	30
HIVCT	HIV 1 & 2 Confirmation Test (ICA) - identify both type 1 & 2	5 ml Plain	7	440
HIVL	HIV-RNA Viral load (RT-PCR)	2x 3 ml EDTA (Plasma), <i>Separate plasma & refrigerate</i>	7	600
HLA12	Human Leukocyte Ag (HLA) Typing (by NGS) Total 12 loci – HLA-A, B, C, DRB1, DQB1, DPB1, DPA1, DQA1, DRB3/4/5 & G	3 ml EDTA	20-30	6000
HLAA	HLA Typing – HLA-A	3 ml EDTA	14-18	700
HLAB	HLA Typing – HLA-B	3 ml EDTA	14-18	700
HLAC	HLA Typing – HLA-C	3 ml EDTA	14-18	700
HLARB	HLA Typing – HLA-DRB1	3 ml EDTA	14-18	700
HLAQB	HLA Typing – HLA-DQB1	3 ml EDTA	14-18	700
HLAPB	HLA Typing – HLA-DPB1	3 ml EDTA	14-18	700
B27	Human Leukocyte Ag (HLA) -B27 Genes	3 ml EDTA	14-18	660
B1502	Human Leukocyte Ag (HLA) -B1502 Genes	3 ml EDTA	14-18	800
B5801	Human Leukocyte Ag (HLA) -B5801 Genes	3 ml EDTA	14-18	1100
HPVD	Human Papilloma Virus (HPV) DNA Genotyping - detect & report 32 major high & low risk genotypes	Cervical Brush in collection kit	7-10	240
HPVS	Human Papilloma Virus (HPV) DNA Screen – detect 16, 18 and 13 other high-risk genotypes	Cervical Brush in collection kit	7	160
CXC	HPV DNA Genotyping (32 types) & Liquid Based PAP	As above	7-10	280
CXCQ	HPV DNA Screen & Liquid Based PAP	As above	7	200
HDGA	Huntington's Disease Genetic Assay	3 ml EDTA	7-14	700
I				
IFE	Immunofixation Electrophoresis (IFE), Serum	5 ml Plain	14	360
IFEU	Immunofixation Electrophoresis (IFE), Urine/CSF	50 ml FMU or 24hr Urine/10 ml CSF	14	440
IGA	Immunoglobulin A (IgA)	5 ml Plain	5	76
IGD	Immunoglobulin D (IgD)	5 ml Plain	8	90

Code	Test Description	Specimen Type	TAT	RPP (RM)
IGE	Immunoglobulin E (IgE), Total	5 ml Plain	5	76
IGEC	Immunoglobulin E (IgE), Low Range (for Newborn)	3 ml EDTA Cord Blood (Plasma)	5	110
IgE for specific allergen (refer to Allergy Section in Profile)				
IGG	Immunoglobulin G (IgG)	5 ml Plain	5	76
IGGS	Immunoglobulin G Subclass 1 - 4	5 ml Plain	10	1200
IgG for specific allergen (refer to Allergy Section in Profile)				
IGM	Immunoglobulin M (IgM)	5 ml Plain	5	76
INAB	Influenza Virus type A & B Antigen - Rapid	Nasopharyngeal Swab	1	180
INHA	Inhibin A	5 ml Plain	7	400
INS	Insulin	5 ml Plain, Fasting sample. Separate serum & freeze. For HOMA index calculation, confirm the fasting blood sugar level performed on the same day as the sample collection for insulin. Interference possible in patients treated with biotin (vitamin B7, B8 or H) or taking any food supplement containing biotin. Essential to stop treatment 8 days before taking the sample.	5	130
IGF1	Insulin-like Growth Factor-1 (IGF-1), Total (Somatomedin-C)	5 ml Plain (fasting), Separate serum & freeze.	7	240
SI	Iron, Serum	5 ml Plain, Fasting sample.	1	12
IA2	Islet Antigen-2 (IA2) Antibodies	5 ml Plain, Separate serum & freeze.	30	250
J				
Joint Fluid (refer to Synovial fluid)				
K				
UKET	Ketones (Acetons) (Semi-quantitative)	20 ml Urine	2	10
L				
LCA	Lactate (Lactic Acid), plasma	3 ml Fluoride, fasting, avoid tourniquet	5	90
LDH	Lactate Dehydrogenase (LDH)	5 ml Plain	1	16
LDHE	LDH Isoenzyme Electrophoresis	5 ml Plain	5	130
LDL1	LDL-Cholesterol (Direct Method)	5 ml Plain	1	12
LEG	<i>Legionella</i> Antibodies	5 ml Plain	7-10	140
LEGUG	<i>Legionella</i> Urinary Antigen	20 ml Urine	5	280
	Leptin	5 ml Plain	Research, call	
LPSM	<i>Leptospira</i> IgM	5 ml Plain	3	280
LIPA	Lipase	5 ml Plain	5	140
LPA	Lipoprotein (a), Lp(a)	5 ml Plain	5	100

Code	Test Description	Specimen Type	TAT	RPP (RM)
LEP	Lipoprotein Electrophoresis	5 ml Plain	5	140
	Liver-Kidney Microsomal 1 Ab (refer to Anti-LKM1)			
LUPA	Lupus Anticoagulant (by PTT-LA & dRVVT)	2x 2* ml Citrate *as specified on tube, Stop heparin therapy one day before test. (freeze Plasma)	5-7	450
LH	Luteinising Hormone (LH)	5 ml Plain	3	44

M

MGS	Magnesium, Serum	5 ml Plain	2	14
MALA	Malaria Parasite	3 ml EDTA	2	16
MEAG	Measles IgG	5 ml Plain	7	100
MEAM	Measles IgM	5 ml Plain	7	130
MTHFR	Methylenetetrahydrofolate reductase (MTHFR) Genetic Assay	3 ml EDTA	7-14	440
UMAL	Microalbumin, Creatinine & Ratio	20 ml Urine (prefer FMU)	2	50
MFP	Microfilaria Parasite	3 ml EDTA	2	20
MON	Monospot (Infectious Mononucleosis) (may also consider EBV)	5 ml Plain	1	30
MUMG	Mumps IgG	5 ml Plain	8	100
MUMM	Mumps IgM	5 ml Plain	8	120
	<i>Mycobacterium tuberculosis</i> Ab (refer to PCR test)			
MTD	<i>Mycobacterium tuberculosis</i> (MTB) DNA (by PCR) Detect MTB & differentiate with other Mycobacteria spp.	Sputum/ Bronchial Wash/ Fresh Tissue/ CSF	5-7	400
TBIGRA	TB- Interferon Gamma Release Assay (TB-IGRA) (QuantIFERON)	5 ml Heparin Blood, reach lab within 2hrs In room temperation, Call Lab	7	700
MGEND	<i>Mycoplasma genitalium</i> DNA Detection	Swab	7	220
MHOMD	<i>Mycoplasma hominis</i> DNA Detection (may consider profile STPC)	Swab	7	220
MPG	<i>Mycoplasma pneumoniae</i> IgG	5 ml Plain	5	110
MPM	<i>Mycoplasma pneumoniae</i> IgM	5 ml Plain	7	150
MYG	Myoglobin, serum	5 ml Plain	5	140
MYGU	Myoglobin, urine	Urine	5	140

N

	<i>Neisseria gonorrhoeae</i> DNA (by PCR) (refer to CTNG Chlamydia & N. gonorrhoeae DNA)			
NSE	Neuron-Specific Enolase (NSE) (small cell lung, neuroblastoma, kidney ca)	5 ml Plain Interference possible in patients treated with biotin (vitamin B7, B8 or H) or taking any food supplement containing biotin. Essential to stop treatment 8 days before taking the sample.	5	150

Code	Test Description	Specimen Type	TAT	RPP (RM)
O				
OSS	Osmolality, Serum	5 ml Plain	5	60
OSU	Osmolality, Urine	20 ml Urine	5	60
OTC	Osteocalcin	3 ml EDTA (Plasma) fasting, morning	7	130
P				
PAP	PAP Smear	Fixed smear	4-6	26
PALP	PAP Test, Liquid based preparation	Cervical brush in Collection kit	4-6	64
PTHl	Parathyroid Hormone, Intact (PTH-l)	3 ml EDTA, Fasting morning Separate serum/plasma a.s.a.p. & refrigerated.	5-7	120
PB19D	Parvovirus B19 DNA (by PCR)	5 ml Plain Blood / CSF	7	240
	Pemphigoid Ab (refer to Anti-Basement Membrane Ab)			
	Pemphigus Ab (refer to Anti-Intercellular Substance Ab)			
PHOS	Phosphorus, Serum	5 ml Plain	1	10
UP	Phosphorus, Urine	24 hr Urine, in Acid preservative and Refrigeration	5	20
	Platelet Antibodies (refer to Anti-Platelets Ab)			
	Platelet Count (refer to Full Blood Count)			
	Porphyrin (refer to Coproporphyrin)			
	Potassium (K) (refer to Electrolytes)			
PALB	Prealbumin	5 ml Plain	5	90
	Pre-Eclampsia Risk (refer to Profile Test)			
PCT	Procalcitonin (PCT)	5 ml Plain	7	320
P1NP	Procollagen 1 N-Propeptide (P1NP)	5 ml EDTA (Plasma)	Research, call	
PRG	Progesterone (P4)	5 ml Plain	5	44
PRL	Prolactin	5 ml Plain	3	44
PACND	Propionibacterium acnes DNA (by PCR)	Swab	7-14	240
PGE2	Prostaglandin E2 (PGE2)	24hr Urine, in Refrigeration without acid preservative Stop anti-inflammatory medication for 2 days	7-14	120
PSA	Prostatic Specific Antigen (PSA)	5 ml Plain	2	50
PSAF	Prostate Specific Ag, Free	5 ml Plain	5	70
PHI	Prostate Health Index (PSA, Fpsa, p2PSA)	5 ml Plain	5	580
PROC	Protein C	2x 2 ml Citrate PC is vitamin K dependant and testing must be performed well after any AVK treatment (1 month). Do not assay in the event of ongoing treatment with a direct-acting oral anticoagulant. Separate plasma & freeze.	15-20	200
PROS	Protein S	2x 2 ml Citrate PS is vitamin K dependant and testing must be performed well after any AVK treatment (1 month). Do not assay in the event of ongoing treatment with a direct-acting oral anticoagulant. Separate plasma & freeze.	10-12	200
PEP	Protein Electrophoresis, Serum	5 ml Plain	5	140

Code	Test Description	Specimen Type	TAT	RPP (RM)
UPEP	Protein Electrophoresis, Urine	20 ml FMU	8-10	200
PRO	Protein, Serum	5 ml Plain	1	10
UPRO	Protein, Urine	24 hr Urine, in Refrigeration without acid preservative	7	20
PT	Prothrombin Time (PT/INR)	2 ml Citrate <i>Do not draw blood from a heparinized line. Avoid contaminating the sample with tissue thromboplastin or heparin. Venipuncture must be performed with no trauma. If blood is drawn from an indwelling catheter, flush with 5 mL of saline and discard the first 5 mL of blood collected. If blood is drawn with a butterfly device, draw a discard tube first to remove air from tubing.</i>	2	50

R

REN	Renin Activity	3 ml EDTA (Plasma), Separate plasma and refrigerate a.s.a.p. Specify patient's posture, upright or recumbent. Normal sodium intake & stop medication for 2 days	7	400
RENC	Renin Concentration, Total	3 ml EDTA (Plasma), Separate plasma and refrigerate a.s.a.p. Specify patient's posture, upright or recumbent. Normal sodium intake & stop medication for 2 days	7	320
	Resistin	5 ml Plain	Research, call	
RSV	Respiratory Syncytial Virus (RSV) Ag	Nasopharyngeal Aspirates	7	200
RC	Reticulocyte Count	3 ml EDTA	2	20
RF	Rheumatoid Factor (Quantitative)	5 ml Plain	2	44
ROMA	Risk of Ovarian Malignancy Algorithm (ROMA) (HE4 + CA-125)	5 ml Plain <i>HE4 and CA125 must be assayed with the same technology. Interference possible in patients treated with biotin (vitamin B7, B8 or H) or taking any food supplement containing biotin. Essential to stop treatment 8 days before taking the sample.</i>	7-10	500
ROTA	Rotavirus Antigen	Fresh Stool	2	140
RG	Rubella IgG	5 ml Plain	2	44
RM	Rubella IgM	5 ml Plain	5	100

S

SARS2R	SARS Coronavirus Type 2 (SARS-CoV-2) Viral RNA (by rtPCR)	Nasopharyngeal & Oropharyngeal Swab in VTM / Sputum/ Brounchial Wash/ Saliva in preservative	call	
SARS2AB	SARS-CoV-2 Antibodies IgM & IgG (by Rapid Test)	5 ml Plain	call	
SC2GS	SARS-CoV-2 S Antibody IgG (by CLIA) – antibodies against receptor binding domain (RBD) of viral spike (S) protein, which may be more likely to confer immunity.	5 ml Plain / 3ml EDTA (plasma)	3-5	160
SARS2AG	SARS-CoV-2 Antigen (by Rapid Test)	Nasopharyngeal Swab in Plain tube	call	
SEMA	Seminal Analysis	Call lab / patient to lab <i>Pre-appointment, specimen must reach lab within 1 hr (State time of collection).</i>	1	80
SRT	Serotonin	5 ml Plain	Research, call	
SHBG	Sex Hormone Binding Globulin (SHBG)	5 ml Plain <i>Interference possible in patients treated with biotin (vitamin B7, B8 or H) or taking any food supplement containing biotin. Essential to stop treatment 8 days before taking the sample.</i>	7	140

Sodium (refer to Electrolytes)

Code	Test Description	Specimen Type	TAT	RPP (RM)
	Somatomedin C (refer to IGF-I)			
SPV	Spermatozoa for Post-vasectomy	Call lab/ patient to lab Specimen must reach lab in 2 hours.	1	24
SMAG	Spinal Muscular Atrophy (SMA) Genetic Assay	3 ml EDTA	7-10	500
	Sputum Culture & Sensitivity (refer to Culture)			
SCC	Squamous Cell Carcinoma Ag (SCC Ag/ TA-4)	5 ml Plain Interference possible in patients treated with biotin (vitamin B7, B8 or H) or taking any food supplement containing biotin. Essential to stop treatment 8 days before taking the sample.	5	160
STOA	Stone (Calculi) Analysis	Kidney/Biliary Stone	14-21	180
STBD	Streptococcus Group B DNA Detection	Swab/ 3ml Blood	5-7	280
	Swab Culture & Sensitivity (refer to Culture)			
SYFEM	Synovial Fluid FEME - Cell count, Gram's stain, protein & crystals	Synovial Aspirate	2	90
	Syphilis (refer to Treponema)			
T				
	Turbocluosis (TB) Tests (AFB stain refer to page C-1, C&S refer to Culture page C-6, PCR & TB-IGRA test refer to Mycobacterium page C-12)			
FTES1	Testosterone, Free	5 ml Plain	7	340
TES	Testosterone, Total	5 ml Plain Interference possible in patients treated with biotin (vitamin B7, B8 or H) or taking any food supplement containing biotin. Essential to stop treatment 8 days before taking the sample.	5	44
THLDA	Thalassaemia Alpha only DNA Analysis (7 mutations)	3 ml Whole/ Cord Blood in EDTA or 10 ml Amniotic Fluid, Avoid contamination from mother's blood.	14	900
THLDB	Thalassaemia Beta only DNA Analysis (22 mutations)	3 ml Whole/ Cord Blood in EDTA or 10 ml Amniotic Fluid, Avoid contamination from mother's blood	14	1300
THLD1	Thalassaemia Alpha & Beta DNA Analysis (adult)	3 ml Whole/ Cord Blood in EDTA or 10 ml Amniotic Fluid, Avoid contamination from mother's blood)	14	1800
THLD2	Thalassaemia Alpha & Beta DNA Analysis (prenatal/neonate)	3 ml Whole/ Cord Blood in EDTA or 10 ml Amniotic Fluid, Avoid contamination from mother's blood	14	1800
TT	Thrombin Time (TT)	2 ml Citrate Specify the current anticoagulant treatment. Attach the clinical data relating to the patient.	5	90
THG	Thyroglobulin	5 ml Plain Interference possible in patients treated with biotin (vitamin B7, B8 or H) or taking any food supplement containing biotin. Essential to stop treatment 8 days before taking the sample.	10	160
	Thyroglobulin Antibody (refer to Anti-Thyroglobulin)			
	Thyroid Anti-Microsomal Ab (refer to Anti-TPO)			
TBG	Thyroid Binding Globulin (TBG)	5 ml Plain	Research, call	
	Thyroid Peroxidase Ab (refer to Anti-TPO)			
TSH	Thyroid Stimulating Hormone (TSH)	5 ml Plain	1	32

Code	Test Description	Specimen Type	TAT	RPP (RM)
TSHR	TSH Receptor Antibody	5 ml Plain <i>Interference possible in patients treated with biotin (vitamin B7, B8 or H) or taking any food supplement containing biotin. Essential to stop treatment 8 days before taking the sample.</i>	7	160
FT4	Thyroxine, Free (FT4)	5 ml Plain <i>Interference possible in patients treated with biotin (vitamin B7, B8 or H) or taking any food supplement containing biotin. Essential to stop treatment 8 days before taking the sample.</i>	1	32
TPA	Tissue Polypeptide Ag (TPA) (non-specific tumour marker)	5 ml Plain	7-10	160
Total Iron Binding Capacity (TIBC) (refer to Transferrin / Iron Status Evaluation Profile)				
TOXG	Toxoplasma IgG	5 ml Plain	5	70
TOXM	Toxoplasma IgM	5 ml Plain	5	90
TOXD	<i>Toxoplasma gondii</i> DNA	5 ml Plain/ CSF/ Amniotic Fluid	5-7	440
TPHA (refer to Treponema Pallidum-Ab test)				
TRA	Transferrin (suggest Iron Status Evaluation profil)	5 ml Plain, <i>Fasting sample.</i>	3	50
Transforming Growth Factor Beta-1 (TGF β-1)			Research, call	
TPD	<i>Treponema pallidum</i> DNA (by PCR)	Swab / CSF	7	180
TPAB	<i>Treponema pallidum</i> Ab (TP-Ab)	5 ml Plain	3	30
FTAM	<i>Treponema</i> , FTA-Abs IgM	5 ml Plain	10	280
FTAG	<i>Treponema</i> , FTA-Abs total (IgG)	5 ml Plain	10	160
RPR	VDRL, Non-treponemal Test (TRUST/RPR Method)	5 ml Plain	1	14
TVD	<i>Trichomonas vaginalis</i> DNA (by PCR)	Genital Swab / FVU	7	160
TG	Triglycerides	5 ml Plain, <i>Fasting sample.</i>	1	10
FT3	Triiodothyronine, Free (FT3)	5 ml Plain <i>Interference possible in patients treated with biotin (vitamin B7, B8 or H) or taking any food supplement containing biotin. Essential to stop treatment 8 days before taking the sample.</i>	2	44
TNIH	Troponin I, High-Sensitive, Quantitative (CMIA)	5 ml Plain	1-3	120
TNIHF	Tropinin I, High-Sensitive, Quantitative (FIA) – Ipoh only	5 ml Plain	1-3	120
TRI	Troponin I (cTnI), Rapid Qualitative	5 ml Plain	1	120
Tumour Necrosis Factor-Alpha (TNF-α)			Research, call	
Typhoid Ab (refer to Widal Test)				
Typhus Ab (refer to Weil-Felix Test)				
U				
UREA	Urea	5 ml Plain	1	10
Urea Breath Test (refer to <i>H Pylori</i>)				
UUREA	Urea, Urine	24 hr Urine, <i>in Refrigeration without acid preservative</i>	3	20
URPD	<i>Ureaplasma spp.</i> DNA Detection	Swab	5-7	240
UA	Uric Acid	5 ml Plain	1	10

Code	Test Description	Specimen Type	TAT	RPP (RM)
UUA	Uric Acid, Urine	24 hr Urine, <i>in Refrigeration without acid preservative</i>	3	20
	Urine Culture & Sensitivity (refer to Culture)			
UFEM	Urine FEME / Urinalysis - 10 chemical parameters & microscopic examination	20 ml MSU	1	16
UPT	Urine Pregnancy Test (Urine β -hCG), rapid	20 ml Urine	1	14
V				
VMA	Vanillylmandelic Acid (VMA)	24 hr Urine, <i>in Acid preservative and Refrigeration</i>	10	140
VMAR	Vanillylmandelic Acid (VMA)/Creatinine Ratio	20 ml Urine (random)	7	140
VZG	Varicella-zoster IgG (Chicken Pox)	5 ml Plain	8	140
VZM	Varicella-zoster IgM (Chicken Pox)	5 ml Plain	12	220
VZD	Varicella-zoster Virus (VZV) DNA (by PCR)	Swab / Urine / CSF / Serum	7	320
	Vascular Endothelial Growth Factor (VEGF)			Research, call
	VDRL (refer to <i>Treponema</i>)			
VA	Vitamin A (Retinol)	5 ml Plain, fasting, wrap, avoid light. Separate serum & freeze.	14-21	400
VB12	Vitamin B12 (Cobalamin)	5 ml Plain <i>Interference possible in patients treated with biotin (vitamin B7, B8 or H) or taking any food supplement containing biotin. Essential to stop treatment 8 days before taking the sample.</i>	7-10	80
VDT	Vitamin D, Total (25-OH D2 & D3)	5 ml Plain, Separate serum & refrigerate a.s.a.p.	5-7	140
W				
WEIL	Weil-Felix Tests (<i>Rickettsia</i> Ab /Typhus)	5 ml Plain	1	30
WET	Wet Mount Examination (refer to <i>Trichomonas</i>)	Vaginal Swab in medium	1	16
	Total White Cell Differential Count, (TWDC) (refer to Full Blood Count)			
WWF	Widal & Weil-Felix Tests (WWF)	5 ml Plain	1	60
WT	Widal Tests (<i>Salmonella</i> Ab/Typhoid)	5 ml Plain	1	40
Z				
ZIKAR	Zika Virus RNA	5 ml Plain/ Urine/ CSF	7-10	600

Service Catalogue LI-SC/01/001 Rev 03(j) Effective: 1 Jan 2026

Code	Test Description	Specimen Type	TAT	RPP (RM)
Therapeutic Drug Assays				
ACET	Acetaminophen / Paracetamol	5 ml Plain	5	180
AMIK	Amikacin	5 ml Plain	5	240
CARB	Carbamazepine (Tegretol)	5 ml Plain	5	160
CCSA	Cyclosporin A	3 ml EDTA Plasma	5	220
DIG	Digoxin (Lanoxin)	5 ml Plain	5	180
FK506	FK506 (Tacrolimus)	3 ml EDTA	5	340
GENT	Gentamycin	5 ml Plain	5	140
LIT	Lithium	5 ml Plain	7	100
MTT	Methotrexate (MTX)	5 ml Plain	5	320
PNB	Phenobarbital (Luminal)	5 ml Plain	5	140
PNT	Phenytoin (Dilantin)	5 ml Plain	5	140
SIRO	Sirolimus (Rapamycin)	3 ml EDTA	7	700
TPL	Theophylline (Aminophylline)	5 ml Plain	5	140
VALA	Valproic Acid (Depakine)	5 ml Plain	5	140
VANC	Vancomycin	5 ml Plain	5	140

Code	Test Description	Specimen Type	TAT	RPP (RM)
	<i>Chemical Hazards</i>			
ACETU	Acetone, Urine	20 ml Urine	7-14	280
CHL1	Cholinesterase, RBC (long term exposure to Pesticides)	3 ml EDTA/Heparin	7-14	320
CHL	Cholinesterase, Serum (acute exposure to Pesticides)	5 ml Plain	3	100
CYANU	Cyanide, Urine (may consider plasma Lactate)	20 ml Urine	7-14	240
SPMA	Benzene (as S-Phenylmercapturic Acid SPMA in Urine)	20 ml Urine	7-14	440
BENZ	Benzene (as Phenol in Urine)	20 ml Urine	7-14	380
BEMA	Benzene (as t,t-Muconic Acid in Urine)	20 ml Urine	7-14	440
FLUU	Fluoride, Urine	20 ml Urine	7-14	280
FORMU	Formaldehyde, Urine	20 ml Urine	7-14	300
MEKU	Methyl Ethyl Ketone (MEK) in Urine	20 ml Urine	7-14	440
HEX	n-Hexane (as 2,5-Hexanedione in Urine)	20 ml Urine	7-14	560
PRQ	Paraquat, Serum	5 ml Plain	7-10	320
PRQU	Paraquat, Urine	20 ml Urine	7-10	320
STYRU	Styrene (as Mandelic Acid in Urine)	20 ml Urine	7-14	240
TOLU	Toluene (as Hippuric Acid in Urine)	20 ml Urine	7-14	200
TCA	Trichloroethylene (as Trichloroacetic Acid in Urine)	20 ml Urine	7-14	440
XYL	Xylene (as Methylhippuric Acid), Urine	20 ml Urine	7-14	240
	<i>Trace & Toxic Elements</i>			
ALU	Aluminium (Al), Blood	5 ml TET Heparin	7-10	180
ALUU	Aluminium (Al), Urine	10 ml Urine	7-14	240
ARS	Arsenic (As), Blood	5 ml EDTA	14-20	260
ARSU	Arsenic (As), Urine	10 ml Urine	14-20	260
ARSU2	Arsenic (AS), Inorganic in Urine	10 ml Urine	7-14	1200
CADM	Cadmium (Cd), Blood	5 ml EDTA	7-14	220
CADMU	Cadmium (Cd), Urine	10 ml Urine	7-14	220
CHRB	Chromium (Cr), Blood	5 ml EDTA	7-14	240
CHRU	Chromium (Cr), Urine	10 ml Urine	7-14	240
COBB	Cobalt (Co), Blood	5 ml in EDTA	7-14	240
COBU	Cobalt (Co), Urine	10 ml Urine	7-14	240
COP	Copper (Cu), Serum	5 ml TET Heparin	7-14	180
COPU	Copper (Cu), Urine	10 ml Urine	7-14	240
PB	Lead (Pb), Blood	5 ml EDTA	7-10	76
UPB	Lead (Pb), Urine	10 ml Urine	7-10	240

Code	Test Description	Specimen Type	TAT	RPP (RM)
MNBL	Manganese (Mn), Blood	5 ml EDTA	7-14	240
MNU	Manganese (Mn), Urine	10 ml Urine	7-14	240
HG	Mercury (Hg), Blood	3 ml EDTA	7-14	200
UHG	Mercury (Hg), Urine	10 ml Urine	7-14	220
NIBL	Nickel (Ni), Blood	5 ml EDTA	7-14	160
NIU	Nickel (Ni), Urine	10 ml Urine in TET Plain	7-14	240
ZN	Zinc (Zn), Serum	5 ml TET Heparin	7-14	160
ZNU	Zinc (Zn), Urine	10 ml Urine	7-14	260

* TET = Trace Element Tube (Royal Blue Cap), Plain (Red label), EDTA (Lavender label), Heparin (Royal Blue Label)

If urine collection (24hrs/Random) via other bottle other than TET, the bottle must be acid washed.

Nutrient & Toxic Element Assay by ICP-MS Method

FNTU6	6 Items Toxic Elements Analysis, Urine ~ Al, As, Cd, Pb, Hg, Ni	Urine, refer below	10-20	720
FNTCU	Toxic Elements Clearance Profile, Urine ~ Al, As, Cd, Pb, Hg & Ni	Urine, refer below	10-20	1440
FNTEB	Toxic Elements Analysis, RBC	Blood, refer below	10-20	1140
FNNEB	Nutrient Elements Analysis, RBC	Blood, refer below	10-20	1140
FNNTB	Nutrient & Toxic Elements Analysis, RBC	Blood, refer below	10-20	1820
FNNTH	Nutrient & Toxic Elements Analysis, Hair	Hair, refer below	10-20	1520

Available for specimen, **H**: Hair, **R**: RBC, **U**: Urine

Toxic Elements	H	R	U		H	R	U		H	R	U
Aluminium (Al)				Germanium (Ge)*				Tellurium (Te)*			
Antimony (Sb)				Lead (Pb)				Thallium (Tl)*			
Arsenic (As)				Mercury (Hg)				Thorium (Th)*			
Barium (Ba)				Nickel (Ni)				Tin (Sn)			
Beryllium (Be)				Palladium (Pd)				Titanium (Ti)			
Bismuth (Bi)				Platinum (Pt)				Tungsten (W)*			
Cadmium (Cd)				Rhodium (Rh)*				Uranium (U)*			
Gadolinium (Gd)*				Rubidium (Rb)*							
Gallium (Ga)*				Silver (Ag)							
Nutrient / Trace Elements											
Boron (B)*				Lithium (Li)*				Selenium (Se)			
Calcium (Ca)				Magnesium (Mg)				Sodium (Na)*			
Chromium (Cr)				Manganese (Mn)				Strontium (Sr)*			
Cobalt (Co)				Molybdenum (Mo)				Sulphur (S)*			
Copper (Cu)				Phosphorous (P)*				Vanadium (V)*			
Iron (Fe)				Potassium (K)				Zinc (Zn)			

Specimen type:

Blood- in 5ml Trace Element (K₂EDTA) Tube (Royal Blue cap and Lavender label)

Urine- First morning urine directly fills in 2 x 5ml Trace Element (Plain) Tube, or urinate in acid washed bottles then transfer to 2x 5ml Trace Element (Plain) Tube (Royal Blue and Red label)

(all blood & urine sample preferably fasting, refrigerated, do not freeze)

Hair- Hair in special collection kit.

Cytopathology

Code	Test Description	Specimen Type	TAT	RPP (RM)
PAP	PAP Smear	Fixed smear	4-6	26
PALP	PAP Test, Liquid-based preparation	Cervical brush in collection kit	4-6	64
CY04	Smear - sputum, brushing etc.	Fixed smear	5-7	240
CY05	Fluid - urine, washing, secretion, body fluid etc.	In Sterile Container	5-7	240
CY06	Fine Needle Aspirate Cytology	Relevant fixed smears	5-7	300

Histopathology

Code	Test Description	Specimen Type	TAT	RPP (RM)
HP01	Small Specimen/Biopsy (<3 cm) - Endometrial curetings, Appendix, Fallopian tubes, Gall bladder, Vas deferens, Lymph nodes, Skin biopsies, Prostatic biopsies, single Ovary, single Tonsil, Cervical punch/ Colposcopic biopsies, Gastro-intestinal tract biopsies.	10% buffered formalin fixed tissue <i>Specimen must be accompanied with appropriate and relevant clinical data. The specimen container must be labelled with patient's identification and anatomic site of the sample.</i>	5-7	330
HP02	Medium Specimen (<6 cm) - Prostatic chips (TURP), Cervical Cone biopsies, Simple Mastectomy, Uterus (without ovaries), Brest Lump, Lumps etc	10% buffered formalin fixed tissue <i>Specimen must be accompanied with appropriate and relevant clinical data. The specimen container must be labelled with patient's identification and anatomic site of the sample.</i>	5-7	480
HP03	Large Specimen (>6 cm) - Bladder, Breast Mastectomy, Caecum, Cervical Cone Biopsy, Colectomy, Gastrectomy, Intestine, Kidney, Liver, Uterus with tubes & ovaries, etc	10% buffered formalin fixed tissue <i>Specimen must be accompanied with appropriate and relevant clinical data. The specimen container must be labelled with patient's identification and anatomic site of the sample.</i>	7-10	830
HP04	Large Complex/Radical Specimen Contact the lab for specific charges - Wertheim's hysterectomy, TAHBSO with entire cervix, etc.	10% buffered formalin fixed tissue <i>Specimen must be accompanied with appropriate and relevant clinical data. The specimen container must be labelled with patient's identification and anatomic site of the sample.</i>	Call	1000
	Immunohistochemistry (IHC) & Special Stains	Contact us for more information on stains available and charges.	Call	Call

