Web Appendix for "A Bayesian Critique of a Rank-Based Method for Surrogate Marker Evaluation"

A. Derivation of θ and δ as Expectations over the Covariate Distribution

This appendix derives expressions for θ and δ as expectations over the covariate distribution.

A.1 *Derivation of θ as an Expectation over the Covariate Distribution*

Under the model in Equation (3), it is easy to see that

$$Y_{1i} - Y_{0i} \mid X_i = x \overset{\text{ind.}}{\sim} \mathcal{N}\left(\frac{\mu^x_{Y_1} - \mu^x_{Y_0}}{\sqrt{\Sigma^x_{Y_1Y_1} + \Sigma^x_{Y_0Y_0} - 2\Sigma^x_{Y_1Y_0}}}\right), \quad i = 1, \ldots, n, \ \forall\, x \in \mathcal{X}.$$

Therefore, we obtain this analytical expression for V^x_Y:

$$\begin{aligned}
V^x_Y &= \mathbb{P}\left(Y_{1i} > Y_{0i} \mid X_i = x\right) \\
&= \mathbb{P}\left(Y_{1i} - Y_{0i} > 0 \mid X_i = x\right) \\
&= \mathbb{P}\left(\frac{Y_{1i} - Y_{0i} - \mu^x_{Y_1} + \mu^x_{Y_0}}{\sqrt{\Sigma^x_{Y_1Y_1} + \Sigma^x_{Y_0Y_0} - 2\Sigma^x_{Y_1Y_0}}} > \frac{-\mu^x_{Y_1} + \mu^x_{Y_0}}{\sqrt{\Sigma^x_{Y_1Y_1} + \Sigma^x_{Y_0Y_0} - 2\Sigma^x_{Y_1Y_0}}} \mid X_i = x\right) \\
&= \mathbb{P}\left(W > \frac{-\mu^x_{Y_1} + \mu^x_{Y_0}}{\sqrt{\Sigma^x_{Y_1Y_1} + \Sigma^x_{Y_0Y_0} - 2\Sigma^x_{Y_1Y_0}}}\right) \\
&= 1 - \Phi\left(\frac{-\mu^x_{Y_1} + \mu^x_{Y_0}}{\sqrt{\Sigma^x_{Y_1Y_1} + \Sigma^x_{Y_0Y_0} - 2\Sigma^x_{Y_1Y_0}}}\right) \\
&= \Phi\left(\frac{\mu^x_{Y_1} - \mu^x_{Y_0}}{\sqrt{\Sigma^x_{Y_1Y_1} + \Sigma^x_{Y_0Y_0} - 2\Sigma^x_{Y_1Y_0}}}\right), \ \forall\, x \in \mathcal{X},
\end{aligned}$$

where $W \sim \mathcal{N}(0,1)$ and Φ is the standard normal CDF. Then, by the law of total probability, we can derive an analytical expression for V_Y as follows:

$$\begin{aligned}
V_Y &= \mathbb{P}\left(Y_{1i} > Y_{0i}\right) \\
&= \int_{\mathcal{X}} V^x_Y \, f(x) \, \mathrm{d}x \\
&= \int_{\mathcal{X}} \Phi\left(\frac{\mu^x_{Y_1} - \mu^x_{Y_0}}{\sqrt{\Sigma^x_{Y_1Y_1} + \Sigma^x_{Y_0Y_0} - 2\Sigma^x_{Y_1Y_0}}}\right) f(x) \, \mathrm{d}x.
\end{aligned}$$

Applying a similar logic to V_S, we obtain this analytical expression for θ:

$$\theta = \int_{\mathcal{X}} \theta^x \, f(x) \, \mathrm{d}x,$$

where we denote

$$\theta^x = \Phi\left(\frac{\mu^x_{Y_1} - \mu^x_{Y_0}}{\sqrt{\Sigma^x_{Y_1Y_1} + \Sigma^x_{Y_0Y_0} - 2\Sigma^x_{Y_1Y_0}}}\right) - \Phi\left(\frac{\mu^x_{S_1} - \mu^x_{S_0}}{\sqrt{\Sigma^x_{S_1S_1} + \Sigma^x_{S_0S_0} - 2\Sigma^x_{S_1S_0}}}\right), \ \forall\, x \in \mathcal{X}.$$

A.2 *Derivation of δ as an Expectation over the Covariate Distribution*

Under the model in Equation (3), it is easy to see that

$$Y_{1i} - Y_{0j} \mid X_i = x_i, X_j = x_j \sim \mathcal{N}\left(\frac{\mu_{Y_1}^{x_i} - \mu_{Y_0}^{x_j}}{\sqrt{\Sigma_{Y_1Y_1}^{x_i} + \Sigma_{Y_0Y_0}^{x_j}}}\right), \ \forall\, x_i, x_j \in \mathcal{X},\ i, j = 1, \ldots, n.$$

Therefore, we obtain this analytical expression for $U_Y^{x_ix_j}$:

$$\begin{aligned}
U_Y^{x_ix_j} &= \mathbb{P}\left(Y_{1i} > Y_{0j} \mid X_i = x_i, X_j = x_j\right) \\
&= \mathbb{P}\left(\frac{Y_{1i} - Y_{0j} - \mu_{Y_1}^{x_i} + \mu_{Y_0}^{x_j}}{\sqrt{\Sigma_{Y_1Y_1}^{x_i} + \Sigma_{Y_0Y_0}^{x_j}}} > \frac{-\mu_{Y_1}^{x_i} + \mu_{Y_0}^{x_j}}{\sqrt{\Sigma_{Y_1Y_1}^{x_i} + \Sigma_{Y_0Y_0}^{x_j}}} \mid X_i = x_i, X_j = x_j\right) \\
&= \mathbb{P}\left(W > \frac{-\mu_{Y_1}^{x_i} + \mu_{Y_0}^{x_j}}{\sqrt{\Sigma_{Y_1Y_1}^{x_i} + \Sigma_{Y_0Y_0}^{x_j}}}\right) \\
&= 1 - \Phi\left(\frac{-\mu_{Y_1}^{x_i} + \mu_{Y_0}^{x_j}}{\sqrt{\Sigma_{Y_1Y_1}^{x_i} + \Sigma_{Y_0Y_0}^{x_j}}}\right) \\
&= \Phi\left(\frac{\mu_{Y_1}^{x_i} - \mu_{Y_0}^{x_j}}{\sqrt{\Sigma_{Y_1Y_1}^{x_i} + \Sigma_{Y_0Y_0}^{x_j}}}\right), \quad \forall\, x_i, x_j \in \mathcal{X},\ i, j = 1, \ldots, n.
\end{aligned}$$

Then, by the law of total probability, we can derive an analytical expression for U_Y as follows:

$$\begin{aligned}
U_Y &= \mathbb{P}\left(Y_{1i} > Y_{0j}\right) \\
&= \int_{\mathcal{X}}\int_{\mathcal{X}} U_Y^{x_i,x_j}\, f(x_i)\, f(x_j)\, \mathrm{d}x_i\, \mathrm{d}x_j \\
&= \int_{\mathcal{X}}\int_{\mathcal{X}} \Phi\left(\frac{\mu_{Y_1}^{x_i} - \mu_{Y_0}^{x_j}}{\sqrt{\Sigma_{Y_1Y_1}^{x_i} + \Sigma_{Y_0Y_0}^{x_j}}}\right) f(x_i)\, f(x_j)\, \mathrm{d}x_i\, \mathrm{d}x_j.
\end{aligned}$$

Applying a similar logic to U_S, we obtain this analytical expression for δ:

$$\delta = \int_{\mathcal{X}}\int_{\mathcal{X}} \delta^{x_ix_j}\, f(x_i)\, f(x_j)\, \mathrm{d}x_i\, \mathrm{d}x_j,$$

where we denote

$$\delta^{x_ix_j} = \Phi\left(\frac{\mu_{Y_1}^{x_i} - \mu_{Y_0}^{x_j}}{\sqrt{\Sigma_{Y_1Y_1}^{x_i} + \Sigma_{Y_0Y_0}^{x_j}}}\right) - \Phi\left(\frac{\mu_{S_1}^{x_i} - \mu_{S_0}^{x_j}}{\sqrt{\Sigma_{S_1S_1}^{x_i} + \Sigma_{S_0S_0}^{x_j}}}\right), \ \forall\, x_i, x_j \in \mathcal{X}.$$

A.3 *Derivation of θ and δ in the Condionally Orthogonal and Homoscedastic Setting*

Under the assumption of conditional orthogonality stated in the paper, it follows that

$$V_Y^x = U_Y^x \text{ and } V_S^x = U_S^x,\ \forall\, x \in \mathcal{X},$$

so that

$$\theta^x = \delta^{xx}\ \forall\, x \in \mathcal{X}. \tag{A.1}$$

Under the assumption of conditional homoscedasticity stated in the paper, it follows that

$$\theta = \mathbb{E}_{X_i \sim f}(g(X_i, X_i))\,, \quad \delta = \mathbb{E}_{(X_i, X_j) \sim f^2}(g(X_i, X_j))\,,$$

where we denote

$$g(x_i, x_j) = \Phi\left(\mu_{Y_1}^{x_i} - \mu_{Y_0}^{x_j}\right) - \Phi\left(\mu_{S_1}^{x_i} - \mu_{S_0}^{x_j}\right)\ \forall\, x_i, x_j \in \mathcal{X}.$$

B. Derivation of $|\theta - \delta|$ in the Binary Covariate Setting

Under the setting described in Section 2.4.1 of the main paper, θ and δ reduce to

$$\theta = \frac{\theta^0 + \theta^1}{2}, \quad \delta = \frac{\delta^{00} + \delta^{11} + \delta^{01} + \delta^{10}}{4}.$$

Based on Equation (A.1), we have that

$$|\theta - \delta| = \frac{|\delta^{00} + \delta^{11} - \delta^{01} - \delta^{10}|}{4}.$$

This expression highlights that the distance between the two parameters is driven by differences between comparisons within the same covariate group and comparisons across different covariate groups. Importantly, this effect does not arise from treatment effect heterogeneity.

To derive a formula for $|\theta - \delta|$, let us further simplify the setting by assuming that S is a perfect surrogate for Y with homogeneous treatment effects across covariate groups:

$$\mu_{Y_1}^x - \mu_{Y_0}^x = \mu_{S_1}^x - \mu_{S_0}^x = \Delta > 0 \quad \forall\, x \in \{0, 1\}.$$

Under this setting, we can easily compute the four terms in the expression for $|\theta - \delta|$:

$$\delta^{00} = \Phi(\Delta) - \Phi(\Delta) = 0,$$

$$\delta^{11} = \Phi(\Delta) - \Phi(\Delta) = 0,$$

$$\delta^{01} = \Phi\big(\Delta + \mu^0_{Y_0} - \mu^1_{Y_0}\big) - \Phi\big(\Delta + \mu^0_{S_0} - \mu^1_{S_0}\big)\,,$$

$$\delta^{10} = \Phi\big(\Delta + \mu^1_{Y_0} - \mu^0_{Y_0}\big) - \Phi\big(\Delta + \mu^1_{S_0} - \mu^0_{S_0}\big)\,.$$

Hence, we conclude that

$$|\theta - \delta| = \frac{1}{4}\left|\Phi(\Delta + d_Y) - \Phi(\Delta + d_S) + \Phi(\Delta - d_Y) - \Phi(\Delta - d_S)\right|,$$

where we denote $d_Y = \mu^0_{Y_0} - \mu^1_{Y_0}$ and $d_S = \mu^0_{S_0} - \mu^1_{S_0}$.

We now prove analytically that $\sup_{d_Y, d_S}\{|\theta - \delta|\} = \frac{1}{4}|1 - 2\Phi(\Delta)|$.

DO THE MATH!

C. Derivation of θ and δ in the Gaussian Covariate Setting

Under the setting described in Section 2.4.2 of the main paper, let $f = \mathcal{N}(m, s^2)$ and $\mu^x = x\,(\beta_{Y_1}, \beta_{S_1}, \beta_{Y_0}, \beta_{S_0})^\top$. We use the following standard identity throughout:

$$\mathbb{E}_{Z\sim\mathcal{N}(0,1)}(\Phi(a + bZ)) = \Phi\left(\frac{a}{\sqrt{1+b^2}}\right), \quad \forall\, a, b \in \mathbb{R}. \tag{A.2}$$

Derivation of δ. Writing $X_i\beta_{Y_1} - X_j\beta_{Y_0} = m(\beta_{Y_1} - \beta_{Y_0}) + s(\beta_{Y_1}Z_i - \beta_{Y_0}Z_j)$ where $Z_i, Z_j \overset{\text{i.i.d.}}{\sim} \mathcal{N}(0,1)$, and applying Equation (A.2) twice (once over Z_j, once over Z_i):

$$\begin{aligned} U_Y &= \mathbb{E}_{X_i, X_j}(\Phi(X_i\beta_{Y_1} - X_j\beta_{Y_0})) \\ &= \Phi\left(\frac{m(\beta_{Y_1} - \beta_{Y_0})}{\sqrt{1 + s^2(\beta^2_{Y_1} + \beta^2_{Y_0})}}\right). \end{aligned}$$

An analogous calculation for U_S yields:

$$\delta = \Phi\left(\frac{m(\beta_{Y_1} - \beta_{Y_0})}{\sqrt{1 + s^2(\beta^2_{Y_1} + \beta^2_{Y_0})}}\right) - \Phi\left(\frac{m(\beta_{S_1} - \beta_{S_0})}{\sqrt{1 + s^2(\beta^2_{S_1} + \beta^2_{S_0})}}\right).$$

Derivation of θ.. Writing $X_i(\beta_{Y_1} - \beta_{Y_0}) = m(\beta_{Y_1} - \beta_{Y_0}) + s(\beta_{Y_1} - \beta_{Y_0})Z_i$ and applying Equation (A.2) over Z_i:

$$V_Y = \mathbb{E}_{X_i}(\Phi(X_i(\beta_{Y_1} - \beta_{Y_0})))$$

$$= \Phi\left(\frac{m(\beta_{Y_1} - \beta_{Y_0})}{\sqrt{1 + s^2(\beta_{Y_1} - \beta_{Y_0})^2}}\right).$$

An analogous calculation for V_S yields:

$$\theta = \Phi\left(\frac{m(\beta_{Y_1} - \beta_{Y_0})}{\sqrt{1 + s^2(\beta_{Y_1} - \beta_{Y_0})^2}}\right) - \Phi\left(\frac{m(\beta_{S_1} - \beta_{S_0})}{\sqrt{1 + s^2(\beta_{S_1} - \beta_{S_0})^2}}\right).$$

Supremum of $|\delta|$ when S is a perfect surrogate.. Setting $\beta_{Y_1} - \beta_{Y_0} = \beta_{S_1} - \beta_{S_0} = \Delta > 0$ so that $\theta = 0$, we have:

$$|\delta| = \left|\Phi\left(\frac{m\Delta}{\sqrt{1 + s^2((\beta_{Y_0} + \Delta)^2 + \beta_{Y_0}^2)}}\right) - \Phi\left(\frac{m\Delta}{\sqrt{1 + s^2((\beta_{S_0} + \Delta)^2 + \beta_{S_0}^2)}}\right)\right|.$$

This is maximized when one of β_{Y_0}, β_{S_0} equals zero and the other diverges to $\pm\infty$. Setting $\beta_{Y_0} = 0$ and $|\beta_{S_0}| \to +\infty$ (or vice versa):

$$\sup_{\beta_{Y_0}, \beta_{S_0}} \{|\delta|\} = \left|\Phi\left(\frac{m\Delta}{\sqrt{1 + s^2\Delta^2}}\right) - \frac{1}{2}\right|.$$

Maximizing further over Δ by taking $|\Delta| \to +\infty$:

$$\sup_{\beta_{Y_0}, \beta_{S_0}, \Delta} \{|\delta|\} = \left|\Phi\left(\frac{m}{s}\right) - \frac{1}{2}\right|,$$

which can be arbitrarily close to $\frac{1}{2}$ whenever $|m/s|$ is large.

D. Regularity and Posterior Consistency

Definition 1 (Regularity of a Bayesian model): Let a Bayesian model be defined as

$$D_i \mid \phi \overset{\text{i.i.d.}}{\sim} f_\phi, \quad i = 1, \ldots, n,$$

$$\phi \sim \pi,$$

where D_i is the observed data for unit i, f_ϕ is the likelihood function parametrized by ϕ, and π is a prior distribution over the parameter ϕ. We say that the Bayesian model is regular if the following conditions hold:

- **Kullback-Leibler support of the prior**: The prior π assigns positive mass to every Kullback-Leibler neighborhood of the true parameter ϕ^*, i.e., for every $\varepsilon > 0$,

$$\pi\left(\{\phi : D_{\mathrm{KL}}(f_{\phi^*} \| f_\phi) < \varepsilon\}\right) > 0,$$

where $D_{\mathrm{KL}}(f_{\phi^*} \| f_\phi)$ is the Kullback-Leibler divergence between f_{ϕ^*} and f_ϕ.

- **Identifiability**: The model is identifiable, i.e., for every $\phi \neq \phi'$, we have that $f_\phi \neq f_{\phi'}$.

THEOREM 1 (Schwartz's theorem (Schwartz, 1965)): *Let $D_i \overset{i.i.d.}{\sim} f_{\phi^*}$ and let the Bayesian model be*

$$D_i \mid \phi \overset{i.i.d.}{\sim} f_\phi, \quad i = 1, \ldots, n,$$

$$\phi \sim \pi.$$

If the model is regular according to Definition 1, the posterior concentrates around ϕ^* as $n \to \infty$:

$$\pi\left(U^c \mid D_{1:n}\right) \overset{P_{\phi^*}}{\to} 0,$$

for every neighborhood U of ϕ^*.

E. Derivation of the Bayes Factor for the Validation Threshold

Under the model described in Section 3.3 of the main paper, the marginal likelihoods under the null and alternative hypotheses are:

$$m\Big(\widehat{V}_S \mid H_0\Big) = \binom{n}{n\,\widehat{V}_S} \frac{1}{2^n},$$

$$m\Big(\widehat{V}_S \mid H_1\Big) = \binom{n}{n\,\widehat{V}_S} \frac{B\Big(a + n\,\widehat{V}_S,\, b + n - n\,\widehat{V}_S\Big)\left(1 - F_{\mathrm{Beta}\left(a+n\,\widehat{V}_S,\, b+n-n\,\widehat{V}_S\right)}\left(\frac{1}{2}\right)\right)}{B(a,b)\left(1 - F_{\mathrm{Beta}(a,b)}\left(\frac{1}{2}\right)\right)},$$

where the expression for $m(\widehat{V}_S \mid H_1)$ follows by integrating the binomial likelihood against the truncated Beta prior over $\left(\frac{1}{2}, 1\right)$. Taking the ratio and cancelling the binomial coefficient yields:

$$BF_n = \frac{B\Big(a + n\,\widehat{V}_S,\, b + n - n\,\widehat{V}_S\Big)\left(1 - F_{\mathrm{Beta}\left(a+n\,\widehat{V}_S,\, b+n-n\,\widehat{V}_S\right)}\left(\frac{1}{2}\right)\right)}{B(a,b)\left(1 - F_{\mathrm{Beta}(a,b)}\left(\frac{1}{2}\right)\right)} \cdot 2^n,$$

where $B(a,b)$ is the Beta function and $F_{\mathrm{Beta}(a,b)}$ is the CDF of the $\mathrm{Beta}(a,b)$ distribution.

The distribution of BF_n under H_0 has closed form: since $n\,\widehat{V}_S \in \{0, \ldots, n\}$, the Bayes

factor takes at most $n+1$ distinct values $x_0, \ldots, x_n$ with

$$\mathbb{P}(BF_n = x_k \mid H_0) = \binom{n}{k}\frac{1}{2^n}, \quad k = 0, \ldots, n,$$

where

$$x_k = \frac{B(a+k,\, b+n-k)\left(1 - F_{\mathrm{Beta}(a+k,\, b+n-k)}\left(\frac{1}{2}\right)\right)}{B(a,b)\left(1 - F_{\mathrm{Beta}(a,b)}\left(\frac{1}{2}\right)\right)} \cdot 2^n, \quad k = 0, \ldots, n.$$

The same formulas apply to the distribution of BF_n under any $V_S = v_S$, by replacing the binomial probabilities $\binom{n}{k}2^{-n}$ with $\binom{n}{k}v_S^k(1-v_S)^{n-k}$.

F. Simulation Study: Data-Generating Processes and Implementation Details

Setting 1 (Useless surrogate).. Potential outcomes are drawn independently from:

$$P_i \overset{\text{i.i.d.}}{\sim} \mathcal{N}_4(\mu^*, \Sigma^*), \quad \mu^* = \left(5,\, \tfrac{2}{3},\, 3,\, \tfrac{2}{3}\right)^\top, \quad \Sigma^* = \mathrm{diag}(3,\, 1,\, 3,\, 1).$$

The surrogate potential outcomes are independent of the primary outcome potential outcomes, so S carries no information about the treatment effect on Y.

Setting 2 (Perfect surrogate).. The primary outcome potential outcomes $(Y_{1i}, Y_{0i}) \overset{\text{i.i.d.}}{\sim} \mathcal{N}_2\big((6, 2.5)^\top,\, 3I_2\big)$, and the surrogate is defined as $S_{gi} = Y_{gi} + \varepsilon_{gi}$, where $(\varepsilon_{1i}, \varepsilon_{0i}) \overset{\text{i.i.d.}}{\sim} \mathcal{N}_2(0,\, 0.1\, I_2)$ independently. The resulting joint distribution of $P_i = (Y_{1i}, S_{1i}, Y_{0i}, S_{0i})^\top$ is $\mathcal{N}_4(\mu^*, \Sigma^*)$ with

$$\mu^* = (6,\, 6,\, 2.5,\, 2.5)^\top, \qquad \Sigma^* = \begin{bmatrix} 3 & 3 & 0 & 0 \\ 3 & 3.1 & 0 & 0 \\ 0 & 0 & 3 & 3 \\ 0 & 0 & 3 & 3.1 \end{bmatrix}.$$

The noise is negligible ($\sigma_\varepsilon^2 = 0.1$), making S nearly identical to Y and therefore a near-perfect surrogate.

Setting 3 (Imperfect surrogate).. The surrogate potential outcomes $(S_{1i}, S_{0i}) \overset{\text{i.i.d.}}{\sim} \mathcal{N}_2\big((5, 3)^\top,\, 3I_2\big)$, and the primary outcome is $Y_{gi} = S_{gi} + \varepsilon_{gi}$, where $\varepsilon_{1i} \overset{\text{i.i.d.}}{\sim} \mathcal{N}(1.5,\, 0.6)$ and $\varepsilon_{0i} \overset{\text{i.i.d.}}{\sim} \mathcal{N}(0,\, 0.6)$

independently. The resulting joint distribution is $\mathcal{N}_4(\mu^*, \Sigma^*)$ with

$$\mu^* = (6.5,\ 5,\ 3,\ 3)^\top, \qquad \Sigma^* = \begin{bmatrix} 3.6 & 3 & 0 & 0 \\ 3 & 3 & 0 & 0 \\ 0 & 0 & 3.6 & 3 \\ 0 & 0 & 3 & 3 \end{bmatrix}.$$

The noise is more substantial ($\sigma_\varepsilon^2 = 0.6$), making S a valid but imperfect surrogate.

Setting 4 (Misspecified model).. The surrogate potential outcomes are log-normal: $S_{1i} \sim \exp(\mathcal{N}(2.5,\ 1.5^2))$ and $S_{0i} \sim \exp(\mathcal{N}(0.5,\ 1.5^2))$. The primary outcome is a non-linear function of the surrogate plus log-normal noise:

$$Y_{1i} = 2.0 + 1.2\sqrt{S_{1i}} + 0.3\exp(S_{1i}/500) + \exp(\varepsilon_{1i}),$$
$$Y_{0i} = 0.8\sqrt{S_{0i}} + 0.2\exp(S_{0i}/50) + \exp(\varepsilon_{0i}),$$

where $\varepsilon_{gi} \overset{\text{i.i.d.}}{\sim} \mathcal{N}(0,\ 0.3)$. This setting is designed to violate the Gaussian parametric assumptions of our Bayesian model.

Setting 5.. A binary baseline covariate $X_i \overset{\text{i.i.d.}}{\sim}$ Bernoulli(0.5) is generated, and the distribution of $P_i \mid X_i$ is the following mixture:

$$P_i \mid X_i \overset{\text{ind.}}{\sim} X_i\, \mathcal{N}_4\big(\mu^1, \Sigma^1\big) + (1 - X_i)\, \mathcal{N}_4\big(\mu^0, \Sigma^0\big), \quad i = 1, \ldots, n,$$

with

$$\mu^0 = (5, 5, 0, 0)^\top, \qquad \mu^1 = (5, -5, 0, -10)^\top,$$
$$\Sigma^0 = \Sigma^1 = \begin{bmatrix} A & 0 \\ 0 & A \end{bmatrix}, \qquad A = \begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix}.$$

Setting 6.. A Gaussian baseline covariate $X_i \overset{\text{i.i.d.}}{\sim} \mathcal{N}(3, 1)$ is generated, and the distribution of $P_i \mid X_i$ follows a linear Gaussian model:

$$P_i \mid X_i \overset{\text{ind.}}{\sim} \mathcal{N}_4(X_i\,\beta,\ \Sigma), \quad i = 1, \ldots, n,$$

with

$$\beta = (1,\ 7,\ 0,\ 6)^\top, \qquad\qquad \Sigma = \tfrac{1}{2}\, I_4.$$

Implementation details.. For the Bayesian method, we ran $T = 500$ posterior sampling iterations with a burn-in of $b = 125$. Prior parameters were set as follows across both models:

$$\mu_0 = (0,0,0,0)^\top, \qquad \Sigma_0 = 10\, I_4, \qquad s = 2, \qquad \tau = 1,$$

$$\mu_\beta = (0,0,0,0)^\top, \qquad \Sigma_\beta = 10\, I_4.$$

The significance level was set to $\alpha = 0.05$. To compute the surrogate validation threshold η, we set the desired power to $1 - \beta = 0.8$ and used a uniform prior for V_S by setting $a = b = 1$.

References

Schwartz, L. (1965). On bayes procedures. *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete* **4,** 10–26.