Biometrics 000, 000–000 DOI: 000
000 0000

A Bayesian Critique of Rank-Based Methods for Surrogate Marker Evaluation

Pietro Carlotti

Department of Statistics and Data Sciences, University of Texas at Austin

pietro.carlotti@utexas.edu

and

Layla Parast

Department of Statistics and Data Sciences, University of Texas at Austin

parast@austin.utexas.edu

Summary: Surrogate markers are often employed in clinical trials to replace primary outcomes that may be difficult, expensive, or time-consuming to measure directly. These markers can accelerate the evaluation of new treatments, provided they reliably capture the causal relationship between treatment and true clinical benefit. Parast et al. (2024) recently proposed a rank-based approach for evaluating surrogate markers, characterized by its nonparametric nature and minimal assumptions. While this method is useful in small-sample model-agnostic settings, it has several limitations, including a lack of clear causal interpretation, low statistical power, and insufficient robustness to different data-generating mechanisms. In this paper, we propose a Bayesian approach that addresses these shortcomings by focusing on causal treatment effect estimands and, in doing so, improves power through covariate adjustment. To demonstrate the advantages of our proposed method, we conduct a comprehensive evaluation using both a simulation study, designed to highlight gains in accuracy and statistical power, and an application to a real clinical trial dataset.
Key words: Bayesian Inference; Surrogate Markers; Clinical Trials; Causal Inference.

This paper has been submitted for consideration for publication in *Biometrics*

1. Introduction

Surrogate markers are often employed in clinical trials to replace primary outcomes that may be difficult, costly, or time-consuming to measure (Elliott, 2023a). These markers can accelerate the evaluation of promising new treatments, provided that they reliably capture the causal relationship between treatment and true clinical benefit. Notably, in the United States, the Food and Drug Administration's (FDA) Accelerated Approval Program, established in 1992, allows new treatments to receive early approval based on demonstrated effects on surrogate markers, and many drugs have been approved, and continue to be approved, through this pathway over time (FDA, 1992; FDA, 2025).

A valid surrogate can substantially reduce trial burden, duration, and cost while still providing meaningful information about treatment efficacy, thereby enabling more timely access to effective treatments. However, establishing *surrogate validity* is a challenging statistical problem, requiring methods that are both robust and interpretable. On this account, surrogate evaluation has been an active area of research, particularly because the FDA does not provide a formal statistical definition of a valid surrogate. A recent review by Elliott (2023b) summarizes the state of proposed and currently used methods, including the proportion of treatment effect explained (PTE) framework (Freedman et al., 1992; Wang and Taylor, 2002; Parast et al., 2016; Parast, 2025), the principal stratification framework (Frangakis and Rubin, 2002; Gilbert and Hudgens, 2008), and the meta-analytic framework (Buyse and Molenberghs, 1998; Burzykowski et al., 2005; Huang and Gilbert, 2011).

Unfortunately, these methods often perform poorly in small trials, especially when using robust techniques such as kernel-based estimation within the PTE framework. To address this, Parast et al. (2024) proposed a rank-based approach for evaluating surrogate markers in such settings. Despite its attractive nonparametric nature and minimal assumptions,

this method has notable limitations, including unclear causal interpretation for its target estimands, low power, and limited robustness across data-generating mechanisms.

In this paper, we improve upon this existing approach by proposing an imputation-based Bayesian alternative that extends the rank-based framework of Parast et al. (2024). Our approach is grounded in Bayesian causal inference, which treats unobserved potential outcomes as latent variables and samples them from predictive distributions (Li et al., 2023). By using this perspective, our method not only provides a causal interpretation but also allows for covariate adjustment, improved efficiency, and a flexible framework for inference.

The rest of this work is organized as follows. In Section 2, we review the existing rank-based approach for surrogate evaluation, outline its limitations, and highlight how the choice of treatment effect estimands can affect conclusions about surrogate validity. In Section 3, we present our Bayesian methodology, including two parametric models and a new criterion for the surrogate validation threshold. In Section 4, we present a simulation study comparing our proposed method with the existing approach, and in Section 5, we apply our method to a clinical trial dataset. Finally, Section 6 discusses potential future extensions.

2. Methods

2.1 *Notation*

Let Y denote the primary outcome and S the surrogate marker. Let $Z \in \{0, 1\}$ be the treatment indicator, assigned at random, where $Z = 1$ indicates treatment and $Z = 0$ control. Without loss of generality, assume higher values of Y and S correspond to greater clinical benefit. We adopt the potential outcomes framework to define causal effects (Rubin, 1974). For each unit $i \in \{1, \ldots, n\}$, let Y_{gi} and S_{gi} denote the potential outcomes for the primary and surrogate under treatment $g \in \{0, 1\}$. The potential outcomes for unit i are

$$P_i = (Y_{0i}, Y_{1i}, S_{0i}, S_{1i})^\top.$$

We assume that for each unit i, the vector of potential outcomes P_i is independently drawn from a joint distribution F_i, which may differ across units, and that the treatment assignment Z_i is independently drawn from a Bernoulli distribution, with $Z_i \perp\!\!\!\perp P_i$ by design. Only the potential outcomes corresponding to the observed treatment assignment are observed. That is, each unit i contributes $P_{1i} = (Y_{1i}, S_{1i})^\top$ when $Z_i = 1$, and $P_{0i} = (Y_{0i}, S_{0i})^\top$ when $Z_i = 0$. The hierarchical process that generates the data is summarized as follows:

$$
\begin{aligned}
(Y_i, S_i)^\top &= \begin{cases} P_{1i}, & \text{if } Z_i = 1, \\ P_{0i}, & \text{if } Z_i = 0, \end{cases} \qquad i = 1, \ldots, n, \\
P_i &\overset{\text{ind.}}{\sim} F_i, \quad i = 1, \ldots, n, \\
Z_i &\overset{\text{i.i.d.}}{\sim} \text{Bernoulli}(p), \quad i = 1, \ldots, n,
\end{aligned}
\tag{1}
$$

where p is the known treatment assignment probability (e.g., $p = 0.5$ for a balanced trial).

In the next section, we describe the existing rank-based approach for evaluating a surrogate marker and call attention to its key limitations.

2.2 *Existing Frequentist Rank-Based Approach*

The main idea of Parast et al. (2024) is that if S is a valid surrogate for Y, then the treatment Z should have comparable effects on both, on a probabilistic scale that is invariant to differences in the measurement units of Y and S. Their method focuses on small-sample settings, where nonparametric kernel methods are infeasible and parametric models are difficult to validate. Motivated by this, they used the following probabilistic index

$$
U_Y = \mathbb{P}(Y_{1i} > Y_{0j}) + \frac{1}{2}\mathbb{P}(Y_{1i} = Y_{0j}), \tag{2}
$$

as a measure of the treatment effect on the primary outcome Y, where i and j index two different units. This quantity is often referred to as the Mann-Whitney parameter (Mann and Whitney, 1947; Thas et al., 2012; Hoeffding, 1992; Buyse, 2010), as it is the estimand targeted by the corresponding hypothesis test. In this work, we restrict our attention to

continuous outcomes, thereby excluding ties in Equation (2). Therefore, a positive treatment effect on Y is concluded if $U_Y > 0.5$. The same reasoning applies to the surrogate S, with U_S defined analogously. The discrepancy in treatment effects is then measured by their difference $\delta = U_Y - U_S$. Based on this definition, Parast et al. (2024) propose that, if S is a valid surrogate for Y, δ should be small, leading to the following hypothesis test

$$\begin{cases} H_0 : \delta \geq \varepsilon, \\ H_1 : \delta < \varepsilon, \end{cases} \tag{3}$$

where ε is a pre-specified threshold for surrogate validation.

To estimate the parameter of interest δ, Parast et al. (2024) compute $\widehat{U}_Y$ and $\widehat{U}_S$ using plug-in estimators, commonly known as the Mann–Whitney U-statistics:

$$\begin{aligned} \widehat{U}_Y &= \frac{1}{n_1 n_0} \sum_{i=1}^{n_1} \sum_{j=1}^{n_0} \mathbb{1}(Y_i > Y_j), \\ \widehat{U}_S &= \frac{1}{n_1 n_0} \sum_{i=1}^{n_1} \sum_{j=1}^{n_0} \mathbb{1}(S_i > S_j), \end{aligned} \tag{4}$$

where n_1 and n_0 are the treated and control sample sizes, and then estimate δ by $\widehat{\delta} = \widehat{U}_Y - \widehat{U}_S$. A confidence interval for δ follows from the central limit theorem for U-statistics and closed-form variance estimators for linear combinations of U-statistics. Parast et al. (2024) conclude that S is valid if the upper bound of this interval is less than the pre-specified threshold ε.

2.3 *Limitations of the Existing Approach*

The nonparametric approach of Parast et al. (2024) suffers from several notable limitations. Firstly, it is worth questioning whether the treatment effect quantities considered truly capture the causal targets. In the causal inference literature, rank-based statistics have in fact been criticized for lacking a clear causal interpretation (Imbens and Rubin, 2015). For example, the Mann–Whitney test assesses whether two variables are equal in distribution, but this condition is not necessarily informative about causal effects. To illustrate this point, note that since U_Y represents the probability that a randomly selected treated unit has a higher Y

than an independently selected control, it captures global distributional differences between groups, rather than a directly interpretable causal effect. This motivates the consideration of alternative estimands that more directly target causal effects, such as

$$V_Y = \mathbb{P}(Y_{1i} > Y_{0i}), \tag{5}$$

which represents the probability that a randomly selected unit has a better outcome Y under treatment than under control. If we define V_S analogously for the surrogate S, then we can measure the discrepancy in treatment effects via $\theta = V_Y - V_S$. Notably, U_Y and U_S are not equivalent to V_Y and V_S, as discussed in detail by Fay et al. (2018). This difference propagates directly to the discrepancy parameters δ and θ. As we illustrate in Section 2.4, they can differ substantially even under relatively simple and well-behaved data-generating mechanisms, and therefore the choice between them may lead to opposing conclusions.

Secondly, the assumptions underlying the Mann–Whitney statistic may be too restrictive. In particular, the test assumes potential outcomes for each observation are independent and identically distributed. The identical distribution assumption is often unrealistic, as potential outcomes are typically influenced by covariates that vary across units.

Moreover, the numerical studies in Parast et al. (2024) show that their procedure achieves high coverage across different settings, but suffers from low power in testing surrogate validity. Approaches that prioritize higher power, even if they rely on parametric assumptions, may better detect valid surrogates, particularly in small-sample settings. Along these lines, leveraging baseline covariates is a natural strategy to increase power, as covariate adjustment improves efficiency and power by reducing residual outcome variability unrelated to the treatment (Zhang et al., 2008; Colantuoni and Rosenblum, 2015; Zhang and Ma, 2019). However, the rank-based framework of Parast et al. (2024) does not readily accommodate covariate adjustment, and incorporating baseline information would likely require a rank regression-type approach, representing a substantial departure from the original framework.

Hence, in Section 3, we describe our Bayesian imputation-based methodology addressing these limitations. In particular, our approach focuses on the causal treatment effect quantities V_Y and V_S, improves power, and allows incorporation of baseline covariates. The following section clarifies the discrepancy parameters and may be skipped without loss of continuity.

2.4 *Comparison between Discrepancy Parameters*

As noted in Section 2.3, the parameters U_Y and V_Y are not equivalent, which carries over to δ and θ. This means that, under the same data-generating mechanism, the two discrepancy parameters can take substantially different values, potentially leading to opposing conclusions about surrogate validity. To illustrate this point, it is useful to consider the analytical forms of δ and θ in a simple setting where the potential outcomes are drawn from a Gaussian distribution whose parameters that depend on a baseline covariate X:

$$\begin{aligned} P_i &\overset{\text{ind.}}{\sim} \mathcal{N}_4\big(\mu^{X_i},\ \Sigma^{X_i}\big)\,, \quad i = 1, \ldots, n, \\ X_i &\overset{\text{i.i.d.}}{\sim} f, \quad i = 1, \ldots, n, \end{aligned} \tag{6}$$

where μ^{X_i} and Σ^{X_i} are the mean vector and covariance matrix that depend on X_i, and f is the distribution of the covariate over some space $\mathcal{X}$. Under the model in Equation (6), θ and δ admit a convenient analytical expression as expectations over the covariate distribution.

Specifically, consider the following two assumptions:

- **Conditional Orthogonality**:

$$\Sigma^x_{Y_1Y_0} = \Sigma^x_{S_1S_0} = 0 \quad \forall\, x \in \mathcal{X}.$$

- **Conditional Homoscedasticity**:

$$\Sigma^x_{Y_1Y_1} = \Sigma^x_{S_1S_1} = \Sigma^x_{Y_0Y_0} = \Sigma^x_{S_0S_0} = \frac{1}{2} \quad \forall\, x \in \mathcal{X}.$$

Then, it follows that

$$\delta = \mathbb{E}_{(X_i,X_j)\sim f\times f}(g(X_i, X_j))\,, \quad \theta = \mathbb{E}_{X_i\sim f}(g(X_i, X_i))\,, \tag{7}$$

where

$$g(x_i, x_j) = \Phi\left(\mu_{Y_1}^{x_i} - \mu_{Y_0}^{x_j}\right) - \Phi\left(\mu_{S_1}^{x_i} - \mu_{S_0}^{x_j}\right) \quad \forall\, x_i, x_j \in \mathcal{X}. \tag{8}$$

Without the two assumptions above, the formula for g still involves a difference of Gaussian CDFs with arguments depending on conditional variances and covariances, leading to a more cumbersome expression. The full derivation is given in Appendix A.

Importantly, we introduce the stronger assumptions stated above for ease of exposition, as they allow us to highlight more clearly that θ and δ may differ substantially whenever $\mathbb{E}_{X_j \sim f}\left(g(X_i, X_j)\right)$ and $g(X_i, X_i)$ differ in expectation. To illustrate this point further, we consider two analytically tractable settings.

2.4.1 *Binary Covariate Setting.* Let us start by considering the simple setting where $f = \text{Bernoulli}\left(\frac{1}{2}\right)$. Although this choice may seem trivial, it can still have a meaningful interpretation in a clinical trial setting (e.g. X represents smoker vs non-smoker status). To simplify the analysis even further, let us also assume treatment effect homogeneity across X:

$$\mu_{Y_1}^{x} - \mu_{Y_0}^{x} = \mu_{S_1}^{x} - \mu_{S_0}^{x} = \Delta, \quad x = 0, 1, \tag{9}$$

for some constant $\Delta > 0$. As we show in Appendix B, the discrepancy between θ and δ is driven by Δ and the difference between means of Y and S for control units across X groups:

$$|\theta - \delta| = \frac{1}{4}|\Phi(\Delta + d_Y) - \Phi(\Delta + d_S) + \Phi(\Delta - d_Y) - \Phi(\Delta - d_S)|, \tag{10}$$

where $d_Y = \mu_{Y_0}^0 - \mu_{Y_0}^1$ and $d_S = \mu_{S_0}^0 - \mu_{S_0}^1$. To understand how the distance between θ and δ changes as a function of d_Y and d_S, it is useful to consider the plot of this quantity in Figure 1 for a fixed value of Δ. It appears that $|\theta - \delta|$ is very large whenever

$$|d_Y| < \Delta \text{ and } |d_S| > \Delta \quad \text{or} \quad |d_Y| > \Delta \text{ and } |d_S| < \Delta. \tag{11}$$

[Figure 1 about here.]

This implies that

$$\sup_{d_Y, d_S} \{|\theta - \delta|\} = \frac{1}{4}|1 - 2\Phi(\Delta)| \Rightarrow \sup_{d_Y, d_S, \Delta} \{|\theta - \delta|\} = \frac{1}{4}, \tag{12}$$

where the first supremum is proved analytically in Appendix B, and the second follows from the principle of iterated suprema (Rudin, 1976). In particular, $\theta = 0$ while $|\delta|$ can be as large as $\frac{1}{4}$, so even though S is a perfect surrogate for Y, δ may be substantially different from 0, while θ correctly reflects surrogate validity.

2.4.2 *Gaussian Covariate Setting.* Let us now consider a setting where $f = \mathcal{N}(m, s^2)$ and the mean potential outcomes are linear functions of the covariate:

$$\mu^x = x\boldsymbol{\beta} \quad \forall\, x \in \mathbb{R}, \quad \text{where} \quad \boldsymbol{\beta} = (\beta_{Y_1}, \beta_{S_1}, \beta_{Y_0}, \beta_{S_0})^\top. \tag{13}$$

As we show in Appendix C, the general formulas for θ and δ are non-trivial functions of the slope parameter $\boldsymbol{\beta}$ and the covariate distribution parameters m and s. Then, to simplify the analysis, let us assume perfect surrogacy by imposing the following assumption:

$$\beta_{Y_1} - \beta_{Y_0} = \beta_{S_1} - \beta_{S_0} = \Delta. \tag{14}$$

This implies that the treatment effect is the same for both Y and S across all values of the covariate. Under this assumption, $\theta = 0$ and therefore $|\theta - \delta| = |\delta|$, which is given by

$$|\theta - \delta| = \left| \Phi\left(\frac{m\Delta}{\sqrt{1 + s^2\left((\beta_{Y_0} + \Delta)^2 + \beta_{Y_0}^2\right)}} \right) - \Phi\left(\frac{m\Delta}{\sqrt{1 + s^2\left((\beta_{S_0} + \Delta)^2 + \beta_{S_0}^2\right)}} \right) \right|. \tag{15}$$

In particular, for $m\Delta \neq 0$, we observe that

$$\sup_{\boldsymbol{\beta}, \Delta}\{|\theta - \delta|\} = \left| \Phi\left(\frac{m\sqrt{2}}{s} \right) - \frac{1}{2} \right| \Rightarrow \sup_{\boldsymbol{\beta}, \Delta, m, s} \{|\theta - \delta|\} = \frac{1}{2},$$

An analytic proof for the first supremum is given in Appendix C, while the second supremum follows by the principle of iterated suprema (Rudin, 1976), and it is approached when $|\frac{m}{s}| \gg 0$, which occurs when the Gaussian covariate is highly concentrated away from zero.

This means that in this setting, even though S is a perfect surrogate for Y, δ can be substantially different from 0 and its value depends arbitrarily on the distribution of the covariate. In contrast, θ correctly captures the surrogate validity by being equal to 0.

Both settings above illustrate that the choice of discrepancy parameter is crucial, as different choices can lead to very different conclusions about surrogate validity. The rank-based approach focuses on δ, which may not capture the causal relationship between Y and S even in simple settings. We therefore advocate focusing on θ instead.

3. Proposed Bayesian Imputation-Based Approach

The central idea of our approach is to estimate the parameter of interest and perform inference through its posterior distribution via Bayesian imputation. Specifically, we use the observed data to impute the unobserved potential outcomes from their posterior distribution. After imputing the missing potential outcomes, we can perform either between-unit or within-unit comparisons between treatment and control groups using any chosen discrepancy metric. Therefore, following the discussion in Section 2.3, we focus on the following quantities:

$$V_Y = \mathbb{P}(Y_{1i} > Y_{0i}), \quad V_S \;\; = \mathbb{P}(S_{1i} > S_{0i}), \quad \theta = V_Y - V_S. \tag{16}$$

As is typically done in Bayesian analysis, we perform inference on θ by sampling from its posterior distribution. At each iteration of our sampling algorithm, we impute the unobserved Y_{gi} and S_{gi} values (e.g., Y_{0i} and S_{0i} for treated units) from their posterior distribution. Using both the observed and imputed outcomes, we compute the probabilities in Equation (16) via Monte Carlo estimators, and obtain a posterior sample for θ by taking their difference. Then, using posterior draws of θ, we can perform inference and assess whether S is a valid surrogate. Specifically, we compute a one-sided credible interval for θ using a quantile of the T posterior draws, after discarding the first b iterations as burn-in:

$$\widehat{\theta}_{1-\alpha} = \text{quantile}\Big(\widehat{\theta}^{(b+1:T)},\, 1-\alpha\Big)\,, \tag{17}$$

and perform the hypothesis test:

$$\begin{cases} H_0 : \theta \geq \eta, \\ H_1 : \theta < \eta. \end{cases} \tag{18}$$

where η is a pre-specified threshold for surrogate validation. We reject the null hypothesis if $\widehat{\theta}_{1-\alpha} < \eta$; otherwise, we fail to reject it. The full procedure is summarized in Algorithm 1.

The primary challenge of this methodology lies in sampling the unobserved potential outcomes. Clearly, for each unit i, the sampling strategy to draw the missing potential outcomes from their posterior distribution crucially depends on the assumed data-generating mechanism, denoted as F_i in Equation (1). Therefore, in the next section, we propose two flexible parametric models that can be used for this sampling step.

3.1 *Parametric Models for the Data Generating Mechanism*

The first model we consider excludes covariates, and therefore more closely aligns with the setting of Parast et al. (2024). Specifically, we define F_i from Equation (1) as follows:

$$\begin{aligned} P_i &\overset{\text{i.i.d.}}{\sim} \mathcal{N}_4(\mu, \Sigma)\,, \quad i = 1, \ldots, n, \\ \mu &\sim \mathcal{N}_4(\mu_0, \Sigma_0), \\ \Sigma &= \operatorname{diag}(\sigma_{1:4})\,\Omega\,\operatorname{diag}(\sigma_{1:4}), \\ \sigma_k &\overset{\text{i.i.d.}}{\sim} \text{Half-Normal}(0, s), \quad k = 1, 2, 3, 4, \\ \Omega &\sim \text{LKJ}(\tau), \end{aligned} \tag{19}$$

where we place a Gaussian prior on the mean vector μ and we adopt a separation strategy (Barnard et al., 2000) for the covariance matrix Σ, decomposing it into a correlation matrix Ω and a scale vector $\sigma_{1:4}$. This allows us to specify easily interpretable priors: an LKJ prior (Lewandowski et al., 2009) for the correlation and a half-normal prior for the scale.

However, one of the main advantages of our proposed Bayesian framework is its ability to naturally incorporate baseline covariates into the data-generating mechanism. To illustrate this feature, we consider a second model that introduces covariates through a linear regression

Algorithm 1 Bayesian Imputation-Based Approach.

Require: Observed data $\{Y_i, S_i, Z_i, X_i\}_{i=1}^n$, prior parameters, number of posterior iterations T, burn-in b, surrogate threshold η, significance level α

```
for t = 1 to T do
    for i = 1 to n do
        if Z_i = 1 then
            Sample P_{0i}^{(t)} ~ p(P_{0i}^{(t)} | P_{1i}, X_i, Z_i = 1)
        else
            Sample P_{1i}^{(t)} ~ p(P_{1i}^{(t)} | P_{0i}, X_i, Z_i = 0)
        end if
    end for
    \widehat{V}_Y^{(t)} ← (1/n) Σ_{i=1}^n 𝟙(Y_{1i}^{(t)} > Y_{0i}^{(t)})
    \widehat{V}_S^{(t)} ← (1/n) Σ_{i=1}^n 𝟙(S_{1i}^{(t)} > S_{0i}^{(t)})
    \widehat{θ}^{(t)} ← \widehat{V}_Y^{(t)} − \widehat{V}_S^{(t)}
end for
\widehat{θ}_{1−α} ← quantile(\widehat{θ}^{(b+1:T)}, 1 − α)
if \widehat{θ}_{1−α} < η then
    Reject H_0
else
    Fail to reject H_0
end if
```

structure, specifying F_i from Equation (1) as follows:

$$
\begin{aligned}
P_i &\overset{\text{ind.}}{\sim} \mathcal{N}_4(BX_i, \Sigma)\,, \quad i = 1, \ldots, n, \\
B &= \left[\beta_1\ \beta_2\ \beta_3\ \beta_4\right]^\top, \\
\beta_k &\overset{\text{i.i.d.}}{\sim} \mathcal{N}_d(\mu_\beta, \Sigma_\beta), \quad k = 1, 2, 3, 4, \\
\Sigma &= \operatorname{diag}(\sigma_{1:4})\,\Omega\,\operatorname{diag}(\sigma_{1:4}), \\
\sigma_k &\overset{\text{i.i.d.}}{\sim} \text{Half-Normal}(0, s), \quad k = 1, 2, 3, 4,
\end{aligned}
\tag{20}
$$

where we place a Gaussian prior on the regression coefficients in the matrix B and we adopt the same separation strategy for the covariance matrix Σ as in the first model.

3.2 *Identifiability of θ and Interpretation of its Posterior*

Both formulations in Equations (19) and (20) are quite simple and intuitive. However, they are both plagued by a common non-identifiability issue. Specifically, it is a well-known fact in the causal inference literature that the joint distribution of the potential outcomes is not identifiable from the observed data, since only one potential outcome is observed for each unit. This issue is typically referred to as the "fundamental problem of causal inference" (Holland, 1986). Therefore, to understand how this issue manifests in our setting, it is useful to take a closer look at the correlation matrix Ω in the two models:

$$\Omega = \begin{bmatrix} 1 & \rho_{Y_1S_1} & \rho_{Y_1Y_0} & \rho_{Y_1S_0} \\ \rho_{Y_1S_1} & 1 & \rho_{S_1Y_0} & \rho_{S_1S_0} \\ \rho_{Y_1Y_0} & \rho_{S_1Y_0} & 1 & \rho_{Y_0S_0} \\ \rho_{Y_1S_0} & \rho_{S_1S_0} & \rho_{Y_0S_0} & 1 \end{bmatrix}. \tag{21}$$

It is easy to see that the green entries in the correlation matrix from Equation (21) are identifiable from the data, since they involve correlations between potential outcomes that are observed together in the data. In contrast, the remaining entries are not identifiable from the data, since they involve correlations between potential outcomes that are never observed together for any unit. Moreover, as we showed in Equation (7) for the Gaussian case, the red entries directly affect the computation of θ, while the orange entries do not.

In principle, one could try to address this non-identifiability issue by imposing further assumptions on the correlation matrix Ω. For example, one could assume that all the non-identifiable entries are equal to zero, which would imply that the potential outcomes are independent of each other. However, this assumption is quite strong and is likely unreasonable in many applications. But if we are not willing to make such strong assumptions, how can

we still make sense of the posterior distribution of θ obtained from our Bayesian imputation approach?

Under standard regularity conditions — specifically, that the prior assigns positive mass to every KL-neighborhood of the truth and that the model is identifiable — the posterior concentrates around the true parameter value (Schwartz's theorem (Schwartz, 1965); see Appendix D for the formal statement and definition of regularity). Our model satisfies the first condition but fails identifiability, and therefore we cannot guarantee that the posterior of θ concentrates around its true value. A characterization of what happens instead is given by Gustafson (2010):

THEOREM 1: *Let the true data-generating process be defined as*

$$D_i \overset{i.i.d.}{\sim} f_{\phi^*}, \quad i = 1, \ldots, n, \tag{22}$$

where D_i is the observed data for unit i and f_{ϕ^*} is the true distribution of the data, parametrized by ϕ^*. Let the parameter ϕ be decomposed as follows:

$$\phi = (\phi_I, \phi_N), \tag{23}$$

where ϕ_I is the identifiable part of the parameter and ϕ_N is the non-identifiable part of the parameter. Let the Bayesian model be defined as

$$\begin{aligned} D_i \mid \phi &\overset{i.i.d.}{\sim} f_{\phi_I}, \quad i = 1, \ldots, n, \\ \phi &\sim \pi, \end{aligned} \tag{24}$$

where π is a prior distribution over the parameter ϕ. Then, if the identifiable component of the Bayesian model is regular according to Definition 1, the posterior distribution of ϕ concentrates around the set of values that are compatible with the observed data and the model assumptions as the sample size goes to infinity:

$$\pi\left(\phi_I \in U_I \mid D_{1:n}\right) \overset{P_{\phi^*}}{\to} 1, \tag{25}$$

for every neighborhood U_I of the identifiable component ϕ_I^*, where P_{ϕ^*} is the probability measure induced by the true distribution f_{ϕ^*}. Whereas, the posterior distribution of the non-

identifiable component ϕ_N will behave as follows:

$$\pi\left(\phi_N \in U_N \mid D_{1:n}\right) \overset{P_{\phi^*}}{\rightarrow} \pi\left(\phi_N \in U_N \mid \phi_I = \phi_I^*\right), \tag{26}$$

for every neighborhood U_N of the non-identifiable component ϕ_N^*, where $\pi\left(\phi_N \in U_N \mid \phi_I = \phi_I^*\right)$ is the conditional prior distribution of the non-identifiable component given the identifiable component evaluated at its true value.

What this last result tells us is that, even though the posterior distribution of θ is not fully identified from the data, there is still a process of Bayesian learning that takes place as the sample size increases, and the posterior distribution of θ will concentrate around a set of values that are compatible with the observed data and the model assumptions. Therefore, it is still possible to meaningfully perform inference on θ for the purpose of surrogate validation. However, it is important to keep in mind that due to the lack of concentration for the non-identifiable component of the parameter, there is a non-vanishing uncertainty in the posterior distribution of θ that persists even in the asymptotic limit. This feature of the proposed Bayesian imputation approach may appear to constitute an inherent limitation. Nevertheless, we argue that the methodology retains merit for at least two reasons.

First, the identifiability issue is not unique to our proposed Bayesian imputation approach, but rather it is a fundamental issue that arises in any approach that relies on the observed data to make inference about the joint distribution of the potential outcomes. Therefore, any method that attempts to assess surrogate validity using the observed data will have to contend with this issue in some form. Then, the non-vanishing uncertainty in the posterior distribution of θ can be seen as a reflection of the inherent uncertainty in the problem of surrogate validation. However, since the amount of uncertainty injected by the prior does not disappear even asymptotically, it becomes of crucial importance to choose reasonable priors that reflect our beliefs about the data-generating process. In this regard, given the prior specifications chosen in Equations (19) and (20) and, more specifically, the choice of the

prior for the correlation matrix Ω, unless we have strong prior beliefs about the correlation structure of the potential outcomes, it is reasonable to choose a weakly informative prior, such as an LKJ prior with $\tau = 1$, which corresponds to a uniform distribution over the space of correlation matrices.

Second, even though our proposed Bayesian imputation approach does not showcase the desirable property of consistency, it still provides useful information about surrogate validity and, most importantly, it may still be preferable to alternative approaches that may not suffer by this non-identifiability issue but are unable to correctly capture the causal relationship between the surrogate and the primary outcome. For example, as we showed in Section 2.3, the discrepancy parameter δ used by Parast et al. (2024) may fail to capture the causal relationship between Y and S even in very simple settings, while our approach targets a parameter θ that is able to correctly reflect surrogate validity. Hence, this suggests that using our approach can lead to more accurate assessments of surrogate validity than the rank-based approach of Parast et al. (2024) and especially in small-sample settings where the lack of asymptotic consistency may not be as problematic.

3.3 *Selection Procedure for the Validation Threshold*

We now turn to the challenge of selecting the prespecified threshold used to assess surrogate validity, denoted ε by Parast et al. (2024) and η in Equation (18). As noted by Parast et al. (2024), the distribution of $\widehat{\delta}$ is not fully specified under the null in Equation (3), so choosing the threshold ε is not a trivial task. They suggest choosing it to ensure a desired power to detect a treatment effect on S. Here, we outline an alternative procedure for selecting η in Equation (18), similar in spirit but better suited to our Bayesian framework, which relies on a Bayes factor, the standard tool for Bayesian hypothesis testing.

The intuition is that we aim to set η such that when V_S is close to V_Y i.e., within η, S is a valid replacement of Y. To do this, we aim to identify v_s, which is defined as the value of

V_S that ensures a power of $(1-\beta)100\%$ for the following hypothesis test:

$$\begin{cases} H_0 : V_S = \frac{1}{2}, \\ H_1 : V_S > \frac{1}{2} \end{cases} \tag{27}$$

implemented using a Bayes factor. Let v_Y denote the hypothesized treatment effect on Y. One reasonable way to choose v_Y is by using the posterior mean of V_Y obtained from our Bayesian imputation framework. Then $v_Y - v_S$ reflects how far we are willing to let v_S be from v_Y, while still ensuring a $(1-\beta)100\%$ power. To this end, the value of η is defined as:

$$\eta = \max\{v_Y - v_S, 0\}. \tag{28}$$

We now describe the details behind obtaining the value v_s. To compute a Bayes factor that can be used to conduct the hypothesis test described in Equation (27), we first need to specify a Bayesian model taking V_S as its unknown parameter:

$$n\,\widehat{V}_S \mid V_S \sim \text{Binomial}(n, V_S), \quad \text{and} \quad V_S \quad \sim p(V_S), \tag{29}$$

where the prior for V_S is a Beta distribution renormalized on $\left(\frac{1}{2}, 1\right)$:

$$p(V_S) = \frac{\text{Beta}(V_S \mid a, b)}{\int_{\frac{1}{2}}^{1} \text{Beta}(v \mid a, b)\, dv} \;\forall\, V_S \in \left(\tfrac{1}{2}, 1\right), \tag{30}$$

with $a, b > 0$ suitably chosen prior parameters, and $\widehat{V}_S = \frac{1}{n}\sum_{i=1}^{n} \mathbb{1}(S_{1i} > S_{0i})$. Note that $\widehat{V}_S$ cannot be computed directly from the observed data, since S_{1i} and S_{0i} are never jointly observed for any unit i. However, as we shall see below, this is not an issue, because the procedure to select v_s only requires the sampling distribution of $\widehat{V}_S$ under the null hypothesis and alternative hypothesis, rather than an observed value for it.

The resulting Bayes factor BF_n is derived and given in closed form in Appendix E. As usual, we would like to ensure a type I error probability of α:

$$\mathbb{P}(BF_n > BF_{n,\alpha} \mid H_0) = \alpha. \tag{31}$$

Since $n\,\widehat{V}_S$ takes values in $\{0, \ldots, n\}$, BF_n has at most $n+1$ distinct values, so $BF_{n,\alpha}$ can be obtained as the $(1-\alpha)$-quantile of BF_n (see Appendix E for the closed-form distribution

of BF_n under H_0). Given $BF_{n,\alpha}$, we can define v_S as the value such that

$$\mathbb{P}\left(BF_n > BF_{n,\alpha} \mid V_S = v_S > \frac{1}{2}\right) = 1 - \beta. \tag{32}$$

It is useful to observe that the map

$$v_S \mapsto \mathbb{P}\big(BF_n > BF_{n,\alpha} \mid V_S = v_S\big) \tag{33}$$

is monotone on $\left(\frac{1}{2}, 1\right)$. Therefore, there exists a unique v_S satisfying Equation (32). The probability can be evaluated using the same procedure described in Appendix E for the distribution of BF_n given $V_S = v_S$. Since the function is continuous and monotone, v_S can be obtained using standard numerical root-finding algorithms, such as `uniroot` in `R`.

4. Simulation Study

We evaluated the performance of our proposed Bayesian imputation-based method against the frequentist rank-based approach of Parast et al. (2024) in terms of test coverage and power via a simulation study in six settings.

Specifically, in Settings 1 to 4, we replicated the data-generating mechanisms used by Parast et al. (2024) to facilitate direct comparison. In Settings 1 to 3 the potential outcomes for the primary outcome were generated as Gaussian variables, but in setting 1 the surrogate was designed to be useless, since its potential outcomes were generated independently of the primary outcome, while in settings 2 and 3 the potential outcomes for the surrogate were obtained by applying a linear transformation of the primary outcome, with the addition of Gaussian noise. Moreover, in Setting 2, the surrogate was designed to be perfect, since the noise was negligible, while in Setting 3, the surrogate was designed to be imperfect, since the noise was more substantial. In contrast, in Setting 4 the outcome was generated from a non-Gaussian distribution and the surrogate was generated by applying a non-linear transformation with the addition of non-Gaussian noise. In all of these settings, we implemented the model in Equation (19) without covariates. Given the true data generating

mechanism, one would expect our proposed method to work well in Settings 1-3, but possibly not work well in Setting 4.

In Settings 5 and 6, we purposefully aimed to break the frequentist rank-based method and highlight the advantage of our proposed approach based on the model in Equation (20) with covariates. Setting 5 features a binary baseline covariate, with generating parameters chosen to maximize the discrepancy between δ and θ, which are approximately 0.25 and 0, respectively. Conditionally on the covariate, the potential outcomes are not identically distributed across units, violating a key assumption of the Mann-Whitney test. In this setting the surrogate is a replica of the primary outcome with the addition of Gaussian noise, with treatment effect homogeneity across the two subpopulations and uncorrelated potential outcomes; therefore, any valid surrogate validation method should correctly identify S as valid. Setting 6 features a Gaussian baseline covariate with potential outcomes following a linear Gaussian model. The conditional treatment effect on Y and S is identical (implying $\theta = 0$), but the marginal rank-based discrepancy δ is nonzero due to the covariate structure, further illustrating the limitation of rank-based methods in covariate-dependent settings. Full details of all data-generating processes, prior parameters, and MCMC settings are given in Appendix F.

We ran 500 simulations for each setting, with a sample size of $n = 50$. We quantify performance in terms of coverage of the confidence/credible intervals of the true parameter (δ for the frequentist method and θ for the Bayesian method) and power, by which we mean the proportion of simulation iterations where the method deemed the surrogate to be valid. In Setting 1, the surrogate is not valid, and thus, this is the Type 1 error rate.

Results are shown in Table 1. In Settings 1 and 2, the rank-based method of Parast et al. (2024) and the Bayesian imputation-based method perform similarly. In in Setting 3,improved performance is observed for the Bayesian method with higher coverage and higher

power. As expected, in Setting 4, where the data-generating process is non-Gaussian and non-linear, the nonparametric rank-based method outperforms the Bayesian method in terms of power. Settings 5 and 6 highlight the true strength of our proposed Bayesian imputation-based method, where the frequentist method very rarely identifies the surrogate as valid, while our method correctly recognizes the surrogate as valid in all iterations. This is not surprising, since the data-generating process in Settings 5 and 6 were purposely designed to illustrate the limitations of the the rank-based method. The proposed Bayesian method is able to adjust for the binary covariate in these settings, and correctly capture surrogacy. The results demonstrate the power gains of our proposed methodology, though we also intentionally explore its limitations under model misspecification.

5. Real Data Application

We illustrate our proposed method using data from the Diabetes Control and Complications Trial (DCCT). This trial was a multicenter, randomized, clinical study designed to determine the effect of an intensive treatment protocol directed at maintaining blood glucose concentrations on progression of early vascular complications in patients with Type 1 diabetes, compared to a control group that received usual care (DCCT, 1987; Lachin and Braffett, 2021; DCCT, 1993). Specifically, the intensive therapy was designed to achieve normal blood glucose levels with three or more insulin injections or an insulin pump. Usual care consisted of 1 or 2 insulin injections per day. Our primary outcome of interest was change in hemoglobin A1c (HbA1c) from baseline to 4.5 years years post-randomization, while the surrogate was an earlier measurement of change in HbA1c from baseline to 1.5 years post-randomization.

Our analysis is focused on a small subgroup of high-risk participants, specifically those in the secondary-intervention cohort (with baseline mild-to-moderate retinopathy) with a body mass index over 25.0 with a history of smoking. This resulted in an analytic sample of 47 participants. Figure 2 shows the distribution of the primary outcome and the surrogate by

treatment arm. Figure 3 illustrates the relationship between the surrogate and the primary outcome and the empirical CDFs, whose horizontal gap directly corresponds to the rank-based quantities U_Y and U_S.

[Figure 2 about here.]

[Figure 3 about here.]

We first applied our proposed method without covariates as described above, with a specified power of 0.70 used to determine the corresponding threshold. The resulting posterior distribution of $\widehat{\theta}$ is shown in the left panel of Figure 4, with an average estimate of 0.027, and a credible interval upper bound of 0.149. The calculated threshold was 0.164 and thus, we conclude that change in HbA1c from baseline to 1.5 years is a surrogate for change in HbA1c from baseline to 4.5 years post-randomization. Next, we applied our proposed procedure incorporating the covariates of age (numeric) and sex (male or female). The resulting posterior distribution of $\widehat{\theta}$ incorporating these covariates is shown in the right panel of Figure 4, with an average estimate of 0.010, and a credible interval upper bound of 0.149. The calculated threshold was 0.158 and thus, we similarly conclude that this is a valid surrogate. For comparison, the method of Parast et al. (2024) without covariates (because covariates are not possible to incorporate) resulted in $\widehat{\delta} = 0.044$, a confidence interval upper bound of 0.120, and with a threshold of 0.132, concluding that this is a surrogate marker. While the conclusions are similar, it is certainly worth noting that the estimated difference quantities ($\widehat{\theta}$ vs $\widehat{\delta}$) are smaller using the proposed definition and estimation approaches. Of course, the true values and true surrogacy here are unknown. Nonetheless, this application highlights the potential utility of our proposed approach and, in particular, demonstrates improved characterization of surrogacy and increased efficiency compared to the existing approach.

[Figure 4 about here.]

6. Discussion

We proposed a Bayesian imputation framework for surrogate evaluation in randomized clinical trials. Our method overcomes limitations of the frequentist rank-based approach, including lack of causal interpretation, inability to use baseline covariates, and low power, particularly in in covariate-dependent settings. By imputing unobserved potential outcomes, it enables flexible treatment-control comparisons and inference on surrogate-primary associations. Incorporating baseline covariates improves applicability to complex scenarios, as shown in simulations where our method achieved perfect power versus 0.075 for the existing approach. Our proposed methods are implemented in the R package `BSET` available at `https://github.com/PietroCarlotti/BSET`, which also contains all reproducible code.

While our approach overcomes some limitations of the rank-based approach, there is room for further improvement, which we are actively pursuing. A key limitation is the reliance on parametric assumptions about the data-generating mechanism, which are vulnerable to model mis-specification. One possible direction, which we plan for future work, is to relax these assumptions by extending the framework to a Bayesian nonparametric setting (Ghosal and Van der Vaart, 2017), allowing a data-driven characterization of both the surrogate-outcome relationship and the covariate-outcome association. However, this flexibility comes at a cost of increased computational complexity and larger required sample sizes (Miller and Harrison, 2018). Finally, sensitivity analyses examining robustness of our conclusions to unidentified covariance parameters would also be useful, and we aim to incorporate such results into our framework and future software.

Funding and Acknowledgments

This work was supported by NIDDK grant R01DK118354 (PI:Parast) and [Pietro add funding..?].

The Diabetes Control and Complications Trial (DCCT) and its follow-up the Epidemiology of Diabetes Interventions and Complications (EDIC) study were conducted by the DCCT/EDIC Research Group and supported by National Institutes of Health (NIH) grants and contracts and by the General Clinical Research Center Program (GCRC), the National Center for Research Resources (NCRR). The resources from the DCCT/EDIC study were supplied by NIDDK Central Repository (NIDDK-CR). This manuscript was not prepared under the auspices of the DCCT/EDIC study and does not represent analyses or conclusions of the DCCT/EDIC study group, NIDDK-CR, or NIH.

References

Barnard, J., McCulloch, R., and Meng, X.-L. (2000). Modeling covariance matrices in terms of standard deviations and correlations, with application to shrinkage. *Statistica Sinica* pages 1281–1311.

Burzykowski, T., Molenberghs, G., and Buyse, M. (2005). *The evaluation of surrogate endpoints*. Springer.

Buyse, M. (2010). Generalized pairwise comparisons of prioritized outcomes in the two-sample problem. *Statistics in medicine* **29,** 3245–3257.

Buyse, M. and Molenberghs, G. (1998). Criteria for the validation of surrogate endpoints in randomized experiments. *Biometrics* pages 1014–1029.

Colantuoni, E. and Rosenblum, M. (2015). Leveraging prognostic baseline variables to gain precision in randomized trials. *Statistics in medicine* **34,** 2602–2617.

DCCT (1987). Diabetes control and complications trial (dcct): results of feasibility study. the dcct research group. *Diabetes Care* **10,** 1–19.

DCCT (1993). Diabetes control and complications trial research group: The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *New England journal of medicine* **329,** 977–986.

Elliott, M. R. (2023a). Surrogate endpoints in clinical trials. *Annual Review of Statistics and Its Application* **10,** 75–96.

Elliott, M. R. (2023b). Surrogate endpoints in clinical trials. *Annual Review of Statistics and its Application* **10,** 75–96.

Fay, M. P., Brittain, E. H., Shih, J. H., Follmann, D. A., and Gabriel, E. E. (2018). Causal estimands and confidence intervals associated with wilcoxon-mann-whitney tests in randomized experiments. *Statistics in Medicine* **37,** 2923–2937.

FDA (1992). U.s. food and drug administration, accelerated approval program. FDA — NDA and BLA Approvals.

FDA (2025). Table of surrogate endpoints that were the basis of drug approval or licensure. https://www.fda.gov/drugs/development-resources/table-surrogate-endpoints-were-basis-drug-approval-or-licensure.

Frangakis, C. E. and Rubin, D. B. (2002). Principal stratification in causal inference. *Biometrics* **58,** 21–29.

Freedman, L. S., Graubard, B. I., and Schatzkin, A. (1992). Statistical validation of intermediate endpoints for chronic diseases. *Statistics in medicine* **11,** 167–178.

Ghosal, S. and Van der Vaart, A. W. (2017). *Fundamentals of nonparametric Bayesian inference*, volume 44. Cambridge University Press.

Gilbert, P. B. and Hudgens, M. G. (2008). Evaluating candidate principal surrogate endpoints. *Biometrics* **64,** 1146–1154.

Gustafson, P. (2010). Bayesian inference for partially identified models. *The international journal of biostatistics* **6,**.

Hoeffding, W. (1992). A class of statistics with asymptotically normal distribution. In *Breakthroughs in statistics: Foundations and basic theory*, pages 308–334. Springer.

Holland, P. W. (1986). Statistics and causal inference. *Journal of the American statistical*

Association **81,** 945–960.

Huang, Y. and Gilbert, P. B. (2011). Comparing biomarkers as principal surrogate endpoints. *Biometrics* **67,** 1442–1451.

Imbens, G. W. and Rubin, D. B. (2015). *Causal inference in statistics, social, and biomedical sciences*. Cambridge university press.

Lachin, J. and Braffett, B. (2021). Diabetes Control and Complications Trial / Epidemiology of Diabetes Interventions and Complications (DCCT/EDIC). Dataset (Version 21). Available from the NIDDK Central Repository.

Lewandowski, D., Kurowicka, D., and Joe, H. (2009). Generating random correlation matrices based on vines and extended onion method. *Journal of multivariate analysis* **100,** 1989–2001.

Li, F., Ding, P., and Mealli, F. (2023). Bayesian causal inference: a critical review. *Philosophical Transactions of the Royal Society A* **381,** 20220153.

Mann, H. B. and Whitney, D. R. (1947). On a test of whether one of two random variables is stochastically larger than the other. *The annals of mathematical statistics* pages 50–60.

Miller, J. W. and Harrison, M. T. (2018). Mixture models with a prior on the number of components. *Journal of the American Statistical Association* **113,** 340–356.

Parast, L. (2025). Surrogate marker evaluation: A tutorial using r. *Statistics in Medicine* **44,** e70048.

Parast, L., Cai, T., and Tian, L. (2024). A rank-based approach to evaluate a surrogate marker in a small sample setting. *Biometrics* **80,** ujad035.

Parast, L., McDermott, M. M., and Tian, L. (2016). Robust estimation of the proportion of treatment effect explained by surrogate marker information. *Statistics in Medicine* **35,** 1637–1653.

Rubin, D. B. (1974). Estimating causal effects of treatments in randomized and nonrandom-

ized studies. *Journal of educational Psychology* **66,** 688.

Rudin, W. (1976). *Principles of Mathematical Analysis*, volume 25 of *International series in pure and applied mathematics*. McGraw-Hill, New York, 3rd edition.

Schwartz, L. (1965). On bayes procedures. *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete* **4,** 10–26.

Thas, O., Neve, J. D., Clement, L., and Ottoy, J.-P. (2012). Probabilistic index models. *Journal of the Royal Statistical Society Series B: Statistical Methodology* **74,** 623–671.

Wang, Y. and Taylor, J. M. (2002). A measure of the proportion of treatment effect explained by a surrogate marker. *Biometrics* **58,** 803–812.

Zhang, M., Tsiatis, A. A., and Davidian, M. (2008). Improving efficiency of inferences in randomized clinical trials using auxiliary covariates. *Biometrics* **64,** 707–715.

Zhang, Z. and Ma, S. (2019). Machine learning methods for leveraging baseline covariate information to improve the efficiency of clinical trials. *Statistics in Medicine* **38,** 1703–1714.

Table 1: Simulation results comparing the frequentist rank-based method and the proposed Bayesian imputation-based method in terms of coverage and power in Settings 1–5, summarized over 500 replications with $n = 50$ per replication. Bolding indicates the better performing method.

Setting	**Valid Surrogate**	**Coverage**		**Power**	
		Frequentist	Bayesian	Frequentist	Bayesian
1	No	0.98	**1.00**	0.00	0.00
2	Yes	0.98	**1.00**	1.00	1.00
3	Yes	0.90	**0.98**	0.54	**0.64**

(continued on next page)

(continued)

Setting	**Valid Surrogate**	**Coverage**		**Power**	
		Frequentist	Bayesian	Frequentist	Bayesian
4 (Misspecified)	Yes	0.96	**1.00**	**0.56**	0.06
5 (Binary Covariate)	Yes	0.925	**0.975**	0.075	**1.00**
6 (Gaussian Covariate)	Yes				



Figure 1: Plot of the distance $|\theta - \delta|$ as a function of d_Y and d_S in the setting where the distribution of the potential outcomes is Gaussian and its parameters depend on a binary covariate. Here we set $\Delta = 5$.



(a) Primary outcome

(b) Surrogate

Figure 2: Diabetes Control and Complications Trial (DCCT) data illustration of the distribution of the primary outcome of interest, change in hemoglobin A1c (HbA1c) from baseline to 4.5 years post-randomization (left panel), and the surrogate, change in HbA1c from baseline to 1.5 years post-randomization (right panel), in the treated and control groups within the high-risk subgroup.



(a) Surrogate vs. primary outcome

(b) Empirical CDFs

Figure 3: Left panel: scatter plot of the surrogate (change in HbA1c at 1.5 years) against the primary outcome (change in HbA1c at 4.5 years) by treatment arm, with LOESS curves. Right panel: empirical cumulative distribution functions of the primary outcome and the surrogate by treatment arm. The horizontal gap between the solid and dashed curves of the same color reflects the treatment effect on each outcome, directly corresponding to the rank-based quantities U_Y and U_S.



(a) Without covariates

(b) With covariates (sex and BMI)

Figure 4: Posterior distributions of $\widehat{\theta}$ when applying the method with no covariates (left panel) and with age and sex (right panel) to the Diabetes Control and Complications Trial (DCCT) data where the primary outcome of interest was change in hemoglobin A1c (HbA1c) from baseline to 4.5 years post-randomization and the surrogate was change in HbA1c from baseline to 1.5 years post-randomization.