

# Algorithms for Random Variate Generation

Pietro Colaguri 1936709

November 2023

## 1 Algorithms

A random variate is a variable whose possible values are outcomes of a random process. In statistics and probability theory, a random variate is often used to represent the numerical outcome associated with a random experiment. The term is used to emphasize that the value of the variable is subject to randomness or uncertainty.

When simulating processes driven by random influences, such as in stochastic processes, modern applications often involve generating random variates corresponding to a specified probability distribution. This task is accomplished using computer procedures designed to produce random variates representing a uniform distribution. These procedures essentially yield values selected from a uniform distribution of pseudorandom numbers.

Procedures dedicated to generating random variates aligned with a specific distribution are known as either procedures for (uniform) random number generation or non-uniform pseudo-random variate generation.

In the realm of probability theory, a random variable is a measurable function mapping from a probability space to a measurable space of possible values. Within this framework, the values taken on by the variable are also termed random variates or random variates. This broader perspective encompasses more than just the concept associated with pseudorandom numbers, expanding the understanding of random variates in a probabilistic context.

Let's proceed by considering some algorithms for random variate generation:

1. **Inverse Transform Method:** Suppose you have  $F$  which is the CDF of  $X$ . If  $F$  is continuous and strictly increasing one can prove that  $X = F^{-1}(U)$ . Where  $U$  is a continuous uniform distribution between 0 and 1. This means that I can simply generate  $u_1, \dots, u_N$  independent random observations from a uniform distribution between 0 and 1 and then compute:

$$x_1 = F^{-1}(u_1), \dots, x_N = F^{-1}(u_N)$$

Then  $x_1, \dots, x_N$  are  $N$  independent random observations of the variables  $X$ .

2. **Convolution Method:** This method is based on "adding things up", as deduced from the name. We start by considering  $n$  independent random variables  $X_1, \dots, X_N$  with known PDFs  $f_1(x), \dots, f_N(x)$ . Let:

$$Y = \sum_{i=1}^N X_i$$

Then the PDF of  $Y$  is given by the convolution of all the  $f_i$  PDFs.

$$f_Y(y) = \int_{-\infty}^{\infty} f_1(y - x_2) f_2(x_2) \dots \int_{-\infty}^{\infty} f_n(y - x_n) f_n(x_n) dx_n \dots dx_2$$

3. **Acceptance-Rejection Method:** In this case we leverage the idea of acceptance or rejection based on a comparison with another simpler, easily generated distribution. This method is particularly useful when it is challenging to directly sample from a target distribution. Let's see the various steps. First we choose the simple distribution, also known as "envelope distribution", then we generate a random variate  $x$  from the envelope distribution. Let  $f(x)$  be the PDF of the target's distribution,  $g(x)$  the PDF of the envelope distribution and  $M$  is a constant s.t.  $M \cdot g(x)$  is a valid upper-bound for  $f(x), \forall x$ ; we can define

$$AcceptanceProbability = \frac{f(x)}{M \cdot g(x)}$$

. At this point we generate a uniform random variate  $u$  from the interval  $[0, 1]$  and proceed as follows: if

$$u \leq AcceptanceProbability$$

then we accept  $x$  as a sample from the target distribution, otherwise reject  $x$  and try generating a new random sample from the envelope distribution. We keep going until we have obtained the desired number of samples.