

## 5.3 GRAM-SCHMIDT

By now we are used to finding bases, but recall that orthogonal, or even better, orthonormal bases are preferred.

**Definition 1.** The vectors  $q_1, \dots, q_n$  are orthonormal if

$$q_i^T q_j = \begin{cases} 0 & \text{if } i \neq j & \text{(giving orthogonality),} \\ 1 & \text{if } i = j & \text{(giving the normalization);} \end{cases} \quad (1)$$

We can also create matrices out of these bases. Notice that the standard basis for an Euclidean space is in the columns of the identity matrix. However, if we want a generic orthonormal basis we need to apply the Gram-Schmidt orthogonalization procedure.

**Theorem 1.** If  $Q$  (square or rectangular) has orthonormal columns, then  $Q^T Q = I$ .

**Definition 2.** An orthogonal matrix is a square matrix with orthonormal columns.

**Theorem 2.** For orthogonal matrices, the transpose is the inverse.

Ex: Consider

$$Q = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \Rightarrow Q^T = Q^{-1} = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}$$

which we can verify by multiplying.

Ex: Any permutation matrix  $P$  (consisting of only row exchanges) is an orthogonal matrix. The criteria of orthonormal columns and square are trivially satisfied. Then we check  $P^{-1} = P^T$  by checking  $PP^T = I$ .

**Theorem 3.** Multiplication by any  $Q$  preserves lengths:  $\|Qx\| = \|x\|$  for all  $x$ .

It also preserves inner products and angles:  $(Qx)^T(Qy) = x^T Q^T Q y = x^T y$ .

Consider  $Qx = b$  where  $q_i$  are the columns of  $Q$ . Then we can write

$$b = x_1 q_1 + x_2 q_2 + \dots + x_i q_i + \dots + x_{n-1} q_{n-1} + x_n q_n$$

If we multiply both sides by  $q_i^T$  we get

$$q_i^T b = 0 + \dots + x_i q_i^T q_i + \dots + 0 = x_i \Rightarrow x = Q^T b.$$

So if your  $A$  is an orthogonal matrix, you don't have to do Gaussian Elimination.

#### The Gram-Schmidt Process

Suppose you are given three independent vectors  $\vec{a}, \vec{b}, \vec{c}$ . If they are orthonormal we can project a vector  $\vec{v}$  onto  $\vec{a}$  by doing  $(\vec{a}^T \vec{v})\vec{a}$ . To project onto the  $\vec{a} - \vec{b}$  plane we do  $(\vec{a}^T \vec{v})\vec{a} + (\vec{b}^T \vec{v})\vec{b}$ , etc.

Process: We are given  $\vec{a}, \vec{b}, \vec{c}$  and we want  $\vec{q}_1, \vec{q}_2, \vec{q}_3$ . No problem with  $q_1$ ; i.e.,  $q_1 = a/\|a\|$  (we don't have to change its direction, just normalize.) The problem begins with  $q_2$ , which has to be orthogonal to  $q_1$ . If the vector  $b$  has any component in the direction of  $q_1$  (i.e., direction of  $a$ ) it has to be subtracted:  $B = b - (q_1^T b)q_1$ , then  $q_2 = B/\|B\|$ , and this continues for  $q_3$ :  $C = c - (q_1^T c)q_1 - (q_2^T c)q_2$ , then  $q_3 = C/\|C\|$ , so on and so forth.

Ex:  $a = (1, 0, 1)$ ,  $b = (1, 0, 0)$ , and  $c = (2, 1, 0)$  for  $A = [a \quad b \quad c]$ .

#### Solution:

Step 1: Make the first vector into a unit vector:  $q_1 = a/\sqrt{2} = (1/\sqrt{2}, 0, 1/\sqrt{2})$ .

Step 2a: Subtract from the second vector its component in the direction of the first:  $B = b - (q_1^T b)q_1 = (1/2, 0, -1/2)$ .

Step 2b: Divide  $B$  by its magnitude:  $q_2 = B/\|B\| = (1/\sqrt{2}, 0, -1/\sqrt{2})$ .

Step 3a: Subtract from the third vector its component in the first and second directions:  $C = c - (q_1^T c)q_1 - (q_2^T c)q_2 = (0, 1, 0)$ .

Step 3b: We normalize  $C$ , but  $C$  is already a unit vector so  $q_3 = (0, 1, 0)$

Then we can write  $Q$  as the matrix

$$Q = \begin{pmatrix} | & | & | \\ q_1 & q_2 & q_3 \\ | & | & | \end{pmatrix} = \begin{pmatrix} 1/\sqrt{2} & 1/\sqrt{2} & 0 \\ 0 & 0 & 1 \\ 1/\sqrt{2} & -1/\sqrt{2} & 0 \end{pmatrix}$$

From the matrix  $Q$  we can get a  $A = QR$  factorization. This means that  $A = QR \Rightarrow Q^T A = R$ , then

$$R = \begin{pmatrix} - & - & q_1^T & - & - \\ - & - & q_2^T & - & - \\ - & - & q_3^T & - & - \end{pmatrix} \begin{pmatrix} | & | & | \\ a & b & c \\ | & | & | \end{pmatrix} = \begin{pmatrix} q_1^T a & q_1^T b & q_1^T c \\ 0 & q_2^T b & q_2^T c \\ 0 & 0 & q_3^T c \end{pmatrix} \quad (2)$$

Now lets do a bunch of examples. I skip some of the computation steps in the notes, but they are done in detail in the lectures.

Ex: Are the vectors orthonormal?

$$\left\{ \begin{bmatrix} 2 \\ -4 \end{bmatrix}, \begin{bmatrix} 2 \\ 1 \end{bmatrix} \right\}$$

They are orthogonal but not normal, so they are not orthonormal.

Ex: Are the vectors orthonormal?

$$\left\{ \begin{bmatrix} 4 \\ -1 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \\ 4 \end{bmatrix}, \begin{bmatrix} -4 \\ -17 \\ -1 \end{bmatrix} \right\}$$

They are orthogonal but not normal, so they are not orthonormal.

Ex: Orthonormalize the following set of vectors

$$\left\{ \begin{bmatrix} 3 \\ 4 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right\}$$

**Solution:** We can get  $q_1 = (3, 4)/5$  immediately. Then

$$B = b - (q_1^T b)q_1 = (1, 0) - \frac{3}{5}(3, 4)/5 = \boxed{(16/25, -12/25)} \Rightarrow q_2 = \frac{(4^2/5^2, -12/5^2)}{\sqrt{(4^4/5^4) + (3^2 \cdot 4^2)/5^4}} = \boxed{(4/5, -3/5)}.$$

Ex: Orthonormalize the following set of vectors

$$\left\{ \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 5 \end{bmatrix} \right\}$$

**Solution:**  $q_1 = (0, 1)$ . Then

$$B = b - (q_1^T b)q_1 = (2, 5) - 5(0, 1) = \boxed{(2, 0)} \Rightarrow \boxed{q_2 = (1, 0)}.$$

Ex: Orthonormalize the following set of vectors

$$\left\{ \begin{bmatrix} 2 \\ 1 \\ -2 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}, \begin{bmatrix} 2 \\ -2 \\ 1 \end{bmatrix} \right\}$$

**Solution:** The vectors are already orthogonal, so just divide by the magnitude.

Ex: Orthonormalize the following set of vectors

$$\left\{ \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} \right\}$$

**Solution:**  $q_1 = (0, 1, 1)/\sqrt{2}$ . Then

$$B = b - (q_1^T b)q_1 = (1, 1, 0) - \frac{1}{\sqrt{2}}(0, 1/\sqrt{2}, 1/\sqrt{2}) = (1, 1/2, -1/2) \Rightarrow q_2 = (1, 1/2, -1/2)/\sqrt{3/2} = (\sqrt{2/3}, \sqrt{2/3}/2, -\sqrt{2/3}/2).$$

And

$$\begin{aligned} C &= c - (q_1^T c)q_1 - (q_2^T c)q_2 = (1, 0, 1) - \frac{1}{\sqrt{2}}(0, 1/\sqrt{2}, 1/\sqrt{2}) - \frac{\sqrt{2/3}}{2}(\sqrt{2/3}, \sqrt{2/3}/2, -\sqrt{2/3}/2) \\ &= (1, 0, 1) - (0, 1/2, 1/2) - (1/3, 1/6, -1/6) = (2/3, -2/3, 2/3). \end{aligned}$$

So,

$$q_3 = (2/3, -2/3, 2/3)/(2\sqrt{2/3}) = (1/\sqrt{3}, -1/\sqrt{3}, 1/\sqrt{3}).$$