

MOMENTS AND MOMENT- GENERATING FUNCTIONS



CALCULUS AHEAD!!

Moments of random variables

- Definition 1: The k th MOMENT [taken about the origin] of a random variable is $\mu'_k = E[Y^k]$.
- The first moment of a random variable is $\mu'_1 = E[Y] = \mu$.
- The second moment is $\mu'_2 = E[Y^2]$




third $E[Y^3]$

Moments of random variables

- Definition 2: The k th CENTRAL MOMENT (or k th moment about the mean) of a random variable is $\mu_k = E[(Y - \mu)^k]$.


- Most common moments:

- Mean: $\mu = \mu_1 = E[Y]$
- Variance: $\sigma^2 = \mu_2 = E[(Y - \mu)^2]$
- Skewness: $\mu_3 = E[(Y - \mu)^3]$
- Kurtosis: $\mu_4 = E[(Y - \mu)^4]$



Moment generating functions (mgfs)

- Definition 3: The MOMENT GENERATING FUNCTION for a random variable Y is $m(t) = E[e^{tY}]$.
- NOTE: the mgf exists if there is some $b > 0$ such that $m(t)$ is finite for $|t| < b$.
- This is an “open neighborhood” of 0.
- If there are no open neighborhoods of 0 for which $E[e^{tY}]$ exists (is finite) then we say that the mgf does not exist.



What is so great about moment generating functions?

- First, we like mgfs because they generate moments . . .
- Second, we like mgfs because they characterize a probability distribution. Each unique probability distribution has its own mgf. So if I can show that some random variable has a mgf associated with some distribution, then the random variable must have that distribution!

mgfs same \Leftrightarrow distributions same

Generating moments with moment generating functions

- Theorem 1: If $m(t)$ exists, then for any positive integer k ,

$$\underline{\mu'_k} = \frac{d^k m(t)}{dt^k} \bigg|_{t=0} = \underline{m^{(k)}(0)}$$

mean $E[Y] = \mu'_1 = \frac{d}{dt} m(t) \bigg|_{t=0} = m'(0)$ $m^{(1)}(0)$

~~variance~~ $E[Y^2] = \mu'_2 = \frac{d^2}{dt^2} m(t) \bigg|_{t=0} = m''(0)$

$$\text{direct } E(Y) = 1 \cdot \frac{1}{10} + 2 \cdot \frac{2}{10} + 3 \cdot \frac{3}{10} + 4 \cdot \frac{4}{10} = 3$$

Generating moments with moment generating functions

y	1	2	3	4
p(y)	$\frac{1}{10}$	$\frac{2}{10}$	$\frac{3}{10}$	$\frac{4}{10}$

- Example 1: If Y is a random variable with probability distribution $p(y) = \frac{y}{10}$ for $y = 1, 2, 3, 4$, find $m(t)$ and use $m(t)$ to calculate $E[Y]$.

$$m(t) = E[e^{ty}] = \sum_{y=1}^4 e^{ty} p(y)$$

$$= e^t \cdot \frac{1}{10} + e^{2t} \cdot \frac{2}{10} + e^{3t} \cdot \frac{3}{10} + e^{4t} \cdot \frac{4}{10}$$

Does this "exist"?

$$|m(t)| < \infty \text{ for all } t \in (-b, b)?$$

Any finite b will work
 \Rightarrow mgf exists!

$$E(Y) = \left. \frac{d}{dt} m(t) \right|_{t=0}$$

$$= \left. e^t \cdot \frac{1}{10} + 2e^{2t} \cdot \frac{2}{10} + 3e^{3t} \cdot \frac{3}{10} + 4e^{4t} \cdot \frac{4}{10} \right|_{t=0}$$

$$= \frac{1}{10} (e^t + 4e^{2t} + 9e^{3t} + 16e^{4t}) \Big|_{t=0} = \frac{1}{10} (1 + 4 + 9 + 16) = \frac{30}{10} = 3$$

$$\boxed{\text{FACT} \quad e^x = \sum_{i=0}^{\infty} \frac{x^i}{i!} \quad \text{A1.11}}$$

Generating moments with moment generating functions

- Example 2: Find the mgf of a Poisson random variable with rate parameter λ and use it to find the mean and variance.

$$m(t) = E[e^{ty}] = \sum_{y=0}^{\infty} e^{ty} p(y) = \sum_{y=0}^{\infty} e^{ty} \frac{\lambda^y e^{-\lambda}}{y!} = e^{-\lambda} \sum_{y=0}^{\infty} \frac{e^{ty} \lambda^y}{y!}$$

$$= e^{-\lambda} e^{\lambda e^t} = e^{\lambda(e^t - 1)}$$

exist? $b > 0$ is $m(t) < \infty$ for $-b < t < b$?
Yes - any finite b

$$\boxed{e^{\lambda e^t - \lambda}}$$

$$\begin{aligned} \text{Let } x &= \lambda e^t \\ \text{FACT:} \\ e^{\lambda e^t} &= \sum_{y=0}^{\infty} \frac{(\lambda e^t)^y}{y!} \\ &= \sum_{y=0}^{\infty} \frac{e^{ty} \lambda^y}{y!} \end{aligned}$$

$$E[Y] = \left. \frac{d}{dt} m(t) \right|_{t=0} = \left. e^{\lambda(e^t-1)} \lambda e^t \right|_{t=0} = \left. \frac{m(t) = e^{\lambda(e^t-1)}}{\lambda \times 1} \right|_{t=0} = \boxed{\lambda}$$

mean

$$E[Y^2] = \left. \frac{d^2}{dt^2} m(t) \right|_{t=0} = \left. \frac{d}{dt} \left(\lambda e^t e^{\lambda(e^t-1)} \right) \right|_{t=0} =$$

$$= \lambda e^t e^{\lambda(e^t-1)} \lambda e^t + e^{\lambda(e^t-1)} \lambda e^t \Big|_{t=0}$$

$$= \lambda e^t e^{\lambda(e^t-1)} (1 + \lambda e^t) \Big|_{t=0} = m''(t)$$

$$= \lambda \cdot 1 \cdot 1 (1 + \lambda) = \lambda + \lambda^2$$

$$\text{Var}(Y) = E[Y^2] - \mu^2 = \lambda + \lambda^2 - (\lambda)^2 = \boxed{\lambda}$$

Poisson dist
mean = variance

Characterizing distributions with moment generating functions

- If a probability distribution $p(y)$ for a random variable Y has (existing) mgf $m(t)$, then it is the unique mgf for Y .
- If two random variables have the same mgf then they must have the same probability distribution.
- Example 3: If a random variable Y has mgf $e^{7.1e^t - 7.1}$, then what is the distribution of Y ?

Poisson $\lambda = 7.1$

$m(t) =$