

FUNCTIONS OF RANDOM VARIABLES

Method of cdfs

Y, pdf f , cdf F


Functions of random variables

- We will now study how to find the distribution of functions of random variables.
 - Functions of a single random variable: Y^2 , \sqrt{Y} , $\text{sign}(Y)$, ..., $g(Y)$ for a fixed known function g
 - Functions of multiple random variables: $\frac{1}{n}(Y_1 + Y_2 + \dots + Y_n)$, $\frac{1}{n-1} \sum_{i=1}^n (Y_i - \bar{Y})^2$, $\min(Y_1, Y_2, \dots, Y_n)$, ..., $h(Y_1, Y_2, \dots, Y_n)$ for a fixed known function h
- Three approaches:
 - Method of cdfs
 - Method of transformation
 - Method of mgfs

Method of cdfs

- We start with a random variable Y and its distribution: either $F(y)$ or $f(y)$ (if Y is continuous) or $p(y)$ (if Y is discrete).
- We want to find the distribution of a “new” random variable U that is a function of Y : $U = U(Y)$.

$$F_u(u) = \begin{cases} 0, & u \leq -1 \\ \left(\frac{u+1}{3}\right)^2, & -1 < u \leq 2 \\ 1, & u > 2 \end{cases}$$



Method of cdfs: Univariate example

- Example 1: Y has pdf $f(y) = 2y$ for $0 \leq y \leq 1$. Find the distribution of $3Y - 1$.


$\textcircled{1}$ Get cdf of Y $P(Y \leq y) = \int_0^y 2x dx = x^2 \Big|_0^y = y^2$

Be careful / clear when expressing cdf (pdf)

$$\underline{F_Y(y)} = \begin{cases} 0 & y \leq 0 \\ y^2 & 0 < y \leq 1 \\ 1 & y > 1 \end{cases}$$

- $\textcircled{2}$ Set $u = 3Y - 1 \rightarrow$ find cdf of u

$$\begin{aligned} F_u(u) &= P(u \leq u) = P(3Y - 1 \leq u) = P(3Y \leq u + 1) \\ &= P(Y \leq \frac{u+1}{3}) = F_Y\left(\frac{u+1}{3}\right) = \begin{cases} 0 & \frac{u+1}{3} \leq 0 \\ \left(\frac{u+1}{3}\right)^2 & 0 < \frac{u+1}{3} \leq 1 \\ 1 & \frac{u+1}{3} \geq 1 \end{cases} \end{aligned}$$



Method of cdfs: Another univariate example

- Example 2: Y has pdf $f(y) = 6y(1 - y)$ for $0 \leq y \leq 1$. Find the distribution of Y^3 .

① Find cdf of Y

$$F_Y(y) = P(Y \leq y) = \int_0^y 6x(1-x)dx = \int_0^y (6x - 6x^2)dx$$

$$= 3x^2 - 2x^3 \Big|_0^y = 3y^2 - 2y^3$$

② $U = Y^3$

$$F_U(u) = P(U \leq u) = P(Y^3 \leq u) = P(Y \leq u^{1/3})$$

$$= F_Y(u^{1/3})$$

$$F_Y(y) = \begin{cases} 0 & y < 0 \\ 3y^2 - 2y^3 & 0 \leq y < 1 \\ 1 & y \geq 1 \end{cases}$$

$$F_U(u) = \begin{cases} 0 & u < 0 \\ 3u^{2/3} - 2u & 0 \leq u < 1 \\ 1 & u \geq 1 \end{cases}$$

pdf $f_U(u) = 2u^{-1/3} - 2$
 $0 \leq u \leq 1$

Method of cdfs: Bivariate case


- We start with two random variables Y_1 and Y_2 and their joint density $f(y_1, y_2)$
- We want to find the distribution of a “new” random variable U that is a function of Y : $U = U(Y_1, Y_2)$.
- Method of cdfs in the bivariate case:
 - Determine possible values of U
 - For u in this domain, integrate $f(y_1, y_2)$ to get $F_U(u)$
 - Differentiate $F_U(u)$ to get $f_U(u)$

$$F_U(u)$$



$$y_2 = Y_1 - u$$

$$y_1 = y_2$$



Method of cdfs: Bivariate example

- Example 3: Random variables Y_1 and Y_2 have joint density $f(y_1, y_2) = 3y_1$ for $0 \leq y_2 \leq y_1 \leq 1$. Find the density of $Y_1 - Y_2$.

$$F_U(u) = P(U \leq u) = P(Y_1 - Y_2 \leq u)$$

$0 \leq u \leq 1$

event!
draw this event in
diagram

$$P(\text{red}) = 1 - P(\text{black})$$

$$= 1 - \int_u^1 \int_0^{y_1-u} 3y_1 \, dy_2 \, dy_1 = 1 - \int_u^1 \left(3y_1 y_2 \Big|_{y_2=0}^{y_1-u} \right) \, dy_1$$


$$= 1 - \int_u^1 3y_1(y_1-u) \, dy_1 = 1 - \int_u^1 \left(3y_1^2 - 3uy_1 \right) \, dy_1 = 1 - \left(y_1^3 - \frac{3}{2}uy_1^2 \right) \Big|_{y_1=u}$$

$$= 1 - \left(1 - \frac{3}{2}u - u^2 + \frac{3}{2}u^3 \right) = \frac{3}{2}u + u^3 - \frac{3}{2}u^3 = \frac{1}{2}(3u - u^3)$$


$0 \leq u \leq 1$

$$f_u(u) = \frac{d}{du} F_u(u) = \frac{1}{2}(3 - 3u^2) = \frac{3}{2}(1 - u^2)$$

$$P\left(\frac{1}{4} < U < \frac{3}{4}\right)$$



- For your own amusement: calculate $E[U]$



Method of cdfs: Another bivariate example

- Example 4: Random variables $\underline{Y_1}$ and $\underline{Y_2}$ have joint density $f(y_1, y_2) = \underline{6e^{-3y_1-2y_2}}$ for $y_1 > 0, y_2 > 0$. Find the $\frac{\text{pdf}}{\text{pdf}}$ of $\underline{Y_1 + Y_2} = u$

Possible values for u ? $0 \leq u < \infty$

$$F_u(u) = P(U \leq u) = P(\underbrace{Y_1 + Y_2 \leq u}_{\text{event}})$$

$$\begin{aligned}
 &= \int_0^u \int_0^{u-y_1} 6e^{-3y_1-2y_2} dy_2 dy_1 = \\
 &= - \int_0^u 6e^{-3y_1} \left(\frac{1}{2} e^{-2y_2} \Big|_0^{u-y_1} \right) dy_1 \\
 &= - \int_0^u 3e^{-3y_1} \left(e^{-2u+2y_1} - 1 \right) dy_1
 \end{aligned}$$

$$= -3 \int_0^1 (e^{-2u-y_1} - e^{-3y_1}) dy_1 = -3 \left[e^{-2u} e^{-y_1} + \frac{1}{3} e^{-3y_1} \Big|_0^1 \right]$$

$$= -3 \left(-e^{-2u} e^{-u} + \frac{1}{3} e^{-3u} + e^{-2u} - \frac{1}{3} \right)$$


$$= -3 \left(-e^{-3u} + \frac{1}{3} e^{-3u} + e^{-2u} - \frac{1}{3} \right)$$

$$= 1 - 3e^{-2u} + 2e^{-3u} = F_u(u),$$

$$F_u(y) = \begin{cases} 0 & u < 0 \\ 1 - 3e^{-2u} + 2e^{-3u} & u \geq 0 \end{cases}$$

$$f_u(u) = \frac{d}{du} (1 - 3e^{-2u} + 2e^{-3u})$$

$$= 6e^{-2u} - 6e^{-3u} = 6(e^{-2u} - e^{-3u}), \quad u \geq 0$$






Transforming a $U(0,1)$ random variable: The inverse cdf method

- If $U \sim U(0,1)$, then the cdf is

$$F(u) = \begin{cases} 0, & u < 0 \\ u, & 0 \leq u \leq 1 \\ 1, & u > 1 \end{cases}$$



- From a uniform random variable we can transform to any distribution F_Y by applying the inverse cdf: $F^{-1}(y)$

pdf of exp $f(y) = \frac{1}{\beta} e^{-\frac{1}{\beta}y}, y > 0$

cdf

$$F(y) = 1 - e^{-\frac{1}{\beta}y}, y > 0$$

Inverse cdf method

- Example 5: Starting with a random variable $U \sim U(0,1)$ find some function h of U so that $Y = h(U) \sim \exp(\beta)$.

TRICK Find the dist of $Y = F^{-1}(u)$
Step 1. Find F^{-1} $1 - e^{-\frac{1}{\beta}y} = u$ solve for y

Step 2 Get dist of $Y = F^{-1}(u)$

$$\begin{aligned} F_Y(y) &= P(Y \leq y) \\ &= P(-\beta \log(1-u) \leq y) \\ &= P(\log(1-u) \geq -\frac{y}{\beta}) \\ &= P(1-u \geq e^{-y/\beta}) \\ &= P(-u \geq e^{-y/\beta} - 1) = P(u \leq 1 - e^{-y/\beta} !!) \end{aligned}$$

$$\begin{aligned} -e^{-\frac{1}{\beta}y} &= u-1 \\ e^{-\frac{y}{\beta}} &= 1-u \\ -\frac{y}{\beta} &= \log(1-u) \\ y &= -\beta \log(1-u) = F^{-1}(u) \end{aligned}$$

Check: $F^{-1}(F(y))$ should get y
 $F(F^{-1}(u))$ s.g. u