


INTRODUCTION TO PROBABILITY:

Counting, Permutations, and Combinations

Counting



- It can help if we can formalize "counting" by reviewing combinatorial analysis.
- This is especially useful when many events are equally likely.
- **Theorem 2.1:** If one set has n elements and another set has m elements then there are $n \times m$ possible pairs with one element from each set.
- Note 1: This can be extended: if $A = \{a_1, a_2, \dots, a_n\}$, $B = \{b_1, b_2, \dots, b_m\}$, and $C = \{c_1, c_2, \dots, c_p\}$, then there are $n \times m \times p$ possible triplets.
- Note 2: The sets could be the same. $A \times A$ $n \times n$

Some simple examples

- Example 1: Three coin tosses. How many possible outcomes?

$$A = \{H, T\} \quad B = \{H, T\} \quad C = \{H, T\}$$
$$2 \quad \times \quad 2 \quad \times \quad 2 = 8$$

- Example 2: Three questions on the Likert scale: one 5-level question, one 7-level question, one 9-level question. How many unique combinations?

$$A = \{1, 2, 3, 4, 5\} -$$
$$B = \{1, 2, \dots, 7\} -$$
$$C = \{1, \dots, 9\} -$$

$$5 \times 7 \times 9$$

Permutations

- PERMUTATION: Ordered arrangement of a specified number of distinct objects. ("Order matters")
- How many ways are there to order exactly r of n objects?

~~A~~ a b c ~~d~~ e $n = 5$
 $r = 2$

ab, ba, ac . . .

A formula for permutations

- Notation: The number of ways to order exactly r of n objects is P_r^n .
- A formula for P_r^n (Theorem 2.2):

$$P_r^n = \frac{n!}{(n-r)!} = \underbrace{n \times (n-1) \times \dots \times (n-r+1)}_{\substack{(n-r) \times (n-r-1) \times \dots \times 1 \\ \text{cancel out}}} \times \cancel{(n-r)} \times \cancel{(n-r-1)} \times \dots \times 1$$

$$A = \{a_1, a_2, \dots, a_n\}$$

choose 1 n possibilities $\leftarrow n$

\Rightarrow A except for one chosen
choose 1 $n-1$ possibilities $\times (n-1)$

\Rightarrow A except for two chosen
 $n-2$ possibilities $\times (n-2) \dots \times (n-r+1)$

Example

- Example 3: Eight patients present at an emergency room. In how many ways (sequences) can the first five patients be triaged?

$$P_5^8 = \frac{8 \times 7 \times 6 \times 5 \times 4}{(8-5)!} = \frac{8 \cdot 7 \cdot 6 \cdot 5 \cdot 4 \cdot \cancel{3!}}{\cancel{3!}}$$

Partitioning


- The number of ways to partition n distinct objects into k distinct groups of sizes n_1, n_2, \dots, n_k , respectively (each object appears in exactly one group) is

$$\binom{n}{n_1 n_2 \dots n_k} = \frac{n!}{n_1! n_2! \dots n_k!}$$

where $n = n_1 + n_2 + \dots + n_k$

Note: Counting exercises when order does not matter involves computing combinations.

Example



- Example 4: How many ways can 10 patients be assigned to 3 treatment groups of size 5, 3, and 2?

$$\binom{10}{5 \ 3 \ 2}$$

$$= \frac{10!}{5! \ 3! \ 2!}$$

$$= \frac{\overset{5}{\cancel{10}} \cdot \cancel{9} \cdot \cancel{8} \cdot \cancel{7} \cdot \cancel{6} \cdot \cancel{5}!}{\cancel{5!} \cdot \cancel{3} \cdot \cancel{2} \cdot \cancel{2}}$$

$$5 \times 9 \times 8 \times 7$$

Combinations with 1 group

- A special case of group partitioning is when there is only 1 group.
- If k objects are chosen from among n distinct objects to be in a group, the number of possible ways is

$$\binom{n}{k} = \frac{n!}{k! (n-k)!}$$

- (If k are in the first group, then the other group must be size $n-k$.)

- Notation: $\binom{n}{k} = \frac{n!}{k!(n-k)!}$. “ n choose k ”

Example

- Example 5: How many ways are there to choose 4 patients out of 9 to receive an experimental treatment?

$$\binom{9}{4} = \frac{9!}{4!5!} = \frac{9 \cdot \cancel{8} \cdot 7 \cdot \cancel{6} \cdot \cancel{5!}}{\cancel{4} \cdot \cancel{3} \cdot \cancel{2} \cdot \cancel{5!}} = 9 \cdot 2 \cdot 7$$