Appendix 1. Sequential Binary Partition

The SBP (Sequential Binary Partition) is a partitioning scheme that iteratively divides a composition into subgroups to facilitate the construction of isometric log ratio (ILR) coordinates. Each partition defines how the parts are grouped into the numerator (+1) and the denominator (-1), while the parts not involved are coded as 0. Each partitioning step generates a row in a matrix called the sign matrix, where each row contains the values +1, -1, and 0 for that particular partition. The +1 values indicate the parts of the composition that will increase, the -1 values indicate the parts that will decrease, and the 0 values indicate the parts that do not change (i.e., they do not participate in that coordinate).

In this SBP, the first partition balances moderate-to-vigorous physical activity (MVPA) against the remaining movement behaviors, with MVPA placed in the numerator and sleep, sedentary behavior (SB), and light physical activity (LPA) placed in the denominator. The second partition balances sleep against SB and LPA, while the final partition balances SB against LPA. Subsequent partitions are therefore performed hierarchically within the denominator subgroup. This hierarchical partitioning generates $D-1$ILR coordinates for a composition with Dparts (Formulas 1, 2, and 3).

The advantage of the SBP is that the parts in the numerator and denominator can be selected theoretically according to the researcher's hypothesis or conceptual framework. Regardless of the ILR generation method used, the statistical results such as coefficients, significance, or model fit do not vary. The method only helps improve their interpretability.

$$1\sqrt{\frac{3}{4}}\log\left(\frac{\text{MVPA}}{(\text{sleep}\cdot\text{SB}\cdot\text{LPA})^{1/3}}\right)$$

$$2\sqrt{\frac{2}{3}}\log\left(\frac{\text{sleep}}{(\text{SB}\cdot\text{LPA})^{1/2}}\right)$$

$$3\sqrt{\frac{1}{2}}\log\left(\frac{\text{SB}}{\text{LPA}}\right)$$

Appendix 2. Missing Data Assessment and Analytic Sample Considerations

2.1 Accelerometer Validity and Analytic Sample


Of the 135 enrolled participants, 81 (60%) met the accelerometer validity criteria, defined as having more **than 180 minutes** of valid sleep data and more than 8 valid waking hours on at least three measurement days. Data loss related to accelerometer validity criteria was consistent with previous studies in preschool-aged children, where attrition rates between 20% and 40% are commonly reported due to device wear-time compliance, removal during sleep, or insufficient valid recording periods.

Missing accelerometry data were primarily attributable to predefined methodological validity criteria rather than participant withdrawal or selective refusal. Because movement-behaviour variables were analyzed within a compositional framework, only participants with complete valid accelerometer compositions could be included in the primary analyses.

2.2 Exploration of Missing Data Patterns

To evaluate the potential impact of missing data on study findings, exploratory analyses of missingness patterns were conducted for both movement-behaviour and developmental outcome variables. Visual inspection of missingness patterns did not reveal strong systematic clustering among accelerometer-derived variables or sociodemographic characteristics, suggesting no clear evidence of highly structured missingness mechanisms within the observed data.

Figure S1. Patterns of missingness across movement-behaviour, developmental, and sociodemographic variables.



Blue cells indicate observed values, whereas pink cells indicate missing values. Numbers shown on the left represent the frequency of observations with the corresponding pattern of missing variables. LPA = light physical activity; MVPA = moderate-to-vigorous physical activity; SES = socioeconomic status; BMI = body mass index.

For developmental outcomes, missingness was more frequent in executive function measures, particularly inhibitory control and working memory. Given the observed missingness in executive function outcomes, additional analyses were conducted to examine whether missingness could be predicted by observed participant characteristics. Logistic regression models were fitted using missingness indicators for executive function outcomes as dependent variables.

Results suggested that younger age and rural locality were significantly associated with a higher probability of missing executive function data. These findings may reflect contextual or procedural challenges, such as connectivity limitations in rural settings or greater difficulty among younger children in completing tablet-based tasks.

Although these findings suggest that missingness may not have been completely random for some developmental outcomes, the observed associations should be interpreted cautiously given the relatively small sample size and the wide confidence intervals observed in several estimates.

Although multiple imputation approaches such as Multiple Imputation by Chained Equations (MICE) could theoretically be considered, their implementation in the present study posed important methodological challenges. The movement-behaviour variables were analyzed as compositional data using isometric log-ratio (ILR) transformations and Dirichlet-based modelling approaches, which complicate standard imputation procedures and require specialized strategies to preserve the compositional structure of the data.

Furthermore, given the exploratory nature of this pilot study and the relatively small analytic sample, multiple imputation was unlikely to substantially improve statistical precision and could

introduce additional model-dependent assumptions. Therefore, complete-case analyses were retained as the primary analytic strategy.

Findings should nevertheless be interpreted cautiously, as selection bias related to missing developmental outcomes cannot be fully excluded. Future longitudinal studies with larger samples should explore compositional multiple imputation approaches and investigate contextual determinants of missingness in early childhood movement-behaviour research.

Appendix 3. Descriptive compositional analysis

Table TS2. Variation matrix

	MVPA	Sleep	Sedentary behavior	LPA
MVPA	0			
Sleep	0.108	0		
Sedentary behavior	0.135	0.033	0	
LPA	0.069	0.042	0.053	0


MVPA = moderate-to-vigorous physical activity, LPA = Light physical activity

Box-and-Whisker Plots

The ratio between pairs of behaviors quantifies the relative variability of movement behaviors across participants and is an appropriate way to interpret variability in compositional data. A ratio equal to 1 indicates that the time spent in behavior x_1 is the same as in behavior x_2; a ratio of 2 means that the time in x_1 is twice that in x_2; and a ratio of 0.5 indicates that the time in x_1 is half that in x_2. However, when plotted directly, these ratios are asymmetric, making it


impossible to apply measures such as variance. By applying the logarithm of the ratios, the relationship becomes symmetric. For example, log(2) = 0.693, whereas log(0.5) = −0.693. This transformation allows for consistent variance calculation and proper assessment of variability between pairs.

Figure FS2. Box-and-whisker plots of the log-ratios between pairs of movement behaviors.



MVPA = Moderate-to-vigorous physical activity, LPA = Light physical activity, SB = Sedentary behavior. The blue dashed line represents the reference point where the ratio equals 1.

Figure FS3. Box-and-whisker plots of the logarithmic ratios between pairs of movement behaviors.



MVPA = Moderate-to-vigorous physical activity, LPA = Light physical activity, SB = Sedentary behavior. The x-axis scale is shown in the original ratio units rather than logarithmic values, so the displayed values correspond to their direct ratio counterparts. The red dashed line represents the reference point where the ratio equals 1.

Appendix 4.

Table TS3. Results from the Dirichlet regression model of SUNRISE Mexico Pilot Study Participants, 2022

	Relative fraction ratio (95% CI)					
	LPA vs MVPA	LPA vs SB	LPA vs Sleep	SB vs MVPA	SB vs Sleep	Sleep vs MVPA
Sex						
Boys	1 (reference)					

Girls	**1.18**** **(1.03, 1.36)**	0.97 (0.87, 1.08)	1.0 (0.90, 1.11)	**1.23**** **(1.09, 1.38)**	1.03 (0.95, 1.12)	**1.19**** **(1.06, 1.33)**
Age (months), SD	1.02 (0.95, 1.11)	0.98 (0.92, 1.03)	1.02 (0.96, 1.07)	1.04* (0.98, 1.10)	**1.04**** **(1.00, 1.09)**	1.00 (0.96, 1.09)
BMI z-score, SD	0.99 (0.93, 1.07)	1.02 (0.97, 1.08)	1.01 (0.96, 1.07)	0.98 (0.92, 1.04)	0.99 (0.95, 1.03)	0.99 (0.93, 1.05)
SES						
High	1 (reference)					
Medium	1.02 (0.81, 1.08)	1.10 (0.98, 1.23)	1.10 (0.98, 1.22)	0.94 (0.83, 1.06)	1.00 (0.92, 1.08)	0.94 (0.84, 1.05)
Low	1.03 (0.89, 1.19)	**1.34**** **(1.10, 1.73)**	1.22* (0.99, 1.50)	**0.74**** **(0.59, 0.93)**	0.89 (0.75, 1.05)	0.84 (0.68, 1.04)
Sector						
Rural	1 (reference)					
Urban	0.94 (0.81, 1.08)	0.92 (0.83, 1.04)	0.89* (0.80, 1.02)	1.01 (0.90, 1.14)	0.97 (0.90, 1.06)	1.04 (0.93, 1.17)
Reference relative fraction	1.09 (0.94, 1.28)	0.42*** (0.36, 0.47)	0.31*** (0.28, 0.35)	2.63*** (2.31, 3.00)	0.75*** (0.68, 0.82)	3.52*** (3.10, 3.99)

Continuous variables were standardized (mean = 0, SD =1), LPA = Light physycal activity, MVPA = Moderate-to-vigorous physical activity, SB = Sedentary behavior, SD = Standar deviation, BMI = Body mass index. SES = Socioeconomic status. *p<0.1, **p<0.05, ***p<0.01

Appendix 5.

Table S1. Logistic regression analyses examining associations between participant characteristics and missing executive function outcomes.

	Missing inhibitory control		Missing working memory		Missing upper-limb strength		Missing fine motor function	
Characteristic[1]	**OR**[1]	**95% CI**[1]	**OR**[1]	**95% CI**[1]	**OR**[1]	**95% CI**[1]	**OR**[1]	**95% CI**[1]
Age (months)	0.80	0.66, 0.93	0.76	0.59, 0.91	0.78	0.49, 1.06	0.84	0.54, 1.16
Sex								
Boy	—	—	—	—	—	—	—	—
Girl	1.18	0.21, 7.31	1.07	0.12, 10.1	0.00		0.00	
SES								
High	—	—	—	—	—	—	—	—
Low	6.17	0.29, 135	0.68	0.01, 28.5	17.4	0.15, 27,713	0.00	
Medium	0.65	0.10, 3.58	0.58	0.06, 4.53	0.73	0.00, 121	0.57	0.00, 62.9
MVPA	0.99	0.96, 1.01	0.99	0.97, 1.02	1.00	0.94, 1.06	1.01	0.95, 1.07
Locality								
Urban	—	—	—	—	—	—	—	—
Rural	6.78	1.38, 43.7	37.1	4.10, 1,080	1.61	0.03, 172	1.24	0.01, 88.8
Sleep	1.01	1.00, 1.02	1.01	1.00, 1.03	1.02	1.00, 1.06	1.02	1.00, 1.06

[1]Values are presented as odds ratios (OR) with 95% confidence intervals. Bold estimates indicate $p < 0.05$.

Abbreviations: CI = Confidence Interval, OR = Odds Ratio

Table TS4. Compositional isotemporal substitution of 24h-MB and their association with motor/executive functions of SUNRISE Mexico Pilot Study Participants, 2022

	Upper Limb Strength (n=78)		
Compositional Isotemporal Substitutions of 24h-MB	**10 min**	**20 min**	**30 min**
↑ MVPA ↓ Sleep	**0.053** (0.003, 0.104)	**0.102** (0.004, 0.201)	**0.148** (0.004, 0.292)
↑ MVPA ↓ SB	**0.070** (0.017, 0.122)	**0.135** (0.032, 0.239)	**0.198** (0.046, 0.350)
↑ MVPA ↓ LPA	**0.180** (0.050, 0.311)	**0.363** (0.100, 0.627)	**0.550** (0.150, 0.950)
↑ Sleep ↓ MVPA	**-0.058** (-0.111, -0.004)	**-0.120** (-0.230, -0.011)	**-0.189** (-0.358, -0.019)
↑ Sleep ↓ SB	0.016 (-0.022, 0.054)	0.032 (-0.044, 0.108)	0.048 (-0.067, 0.162)
↑ Sleep ↓ LPA	**0.127** (0.025, 0.229)	**0.260** (0.051, 0.468)	**0.400** (0.078, 0.721)
↑ SB ↓ MVPA	**-0.074** (-0.129, -0.018)	**-0.152** (-0.266, -0.039)	**-0.236** (-0.411, -0.061)
↑ SB ↓ Sleep	-0.016 (-0.054, 0.021)	-0.033 (-0.108, 0.043)	-0.049 (-0.162, 0.063)
↑ SB ↓ LPA	**0.111** (0.008, 0.213)	**0.228** (0.018, 0.438)	**0.352** (0.029, 0.676)
↑ LPA ↓ MVPA	**-0.179** (-0.307, -0.050)	**-0.357** (-0.613, -0.101)	**-0.536** (-0.919, -0.152)
↑ LPA ↓ Sleep	**-0.121** (-0.219, -0.024)	**-0.237** (-0.428, -0.046)	**-0.349** (-0.630, -0.068)
↑ LPA ↓ SB	**-0.105** (-0.203, -0.006)	**-0.204** (-0.398, -0.011)	**-0.299** (-0.584, -0.015)

	Lower Limb Strength (n=79)		
Compositional Isotemporal Substitutions of 24h-MB	**10 min**	**20 min**	**30 min**
↑ MVPA ↓ Sleep	-0.418 (-1.335, 0.500)	-0.833 (-2.642, 0.975)	-1.247 (-3.923, 1.430)
↑ MVPA ↓ SB	0.240 (-0.484, 0.965)	0.496 (-0.933, 1.926)	0.768 (-1.353, 2.888)
↑ MVPA ↓ LPA	-0.244 (-1.863, 1.376)	-0.486 (-3.770, 2.797)	-0.729 (-5.734, 4.276)
↑ Sleep ↓ MVPA	0.422 (-0.527, 1.370)	0.849 (-1.085, 2.782)	1.282 (-1.680, 4.244)
↑ Sleep ↓ SB	0.654 (-0.232, 1.539)	1.311 (-0.464, 3.087)	1.973 (-0.698, 4.645)
↑ Sleep ↓ LPA	0.170 (-1.314, 1.653)	0.329 (-2.705, 3.362)	0.476 (-4.184, 5.136)

↑ SB ↓ MVPA	-0.224 (-0.972, 0.524)	-0.431 (-1.956, 1.095)	-0.619 (-2.959, 1.720)
↑ SB ↓ Sleep	-0.650 (-1.531, 0.231)	-1.297 (-3.056, 0.461)	-1.942 (-4.575, 0.691)
↑ SB ↓ LPA	-0.476 (-2.051, 1.099)	-0.951 (-4.165, 2.263)	-1.425 (-6.350, 3.500)
↑ LPA ↓ MVPA	0.246 (-1.340, 1.833)	0.497 (-2.654, 3.648)	0.754 (-3.951, 5.459)
↑ LPA ↓ Sleep	-0.180 (-1.604, 1.244)	-0.370 (-3.165, 2.426)	-0.569 (-4.689, 3.552)
↑ LPA ↓ SB	0.478 (-1.042, 1.999)	0.960 (-2.033, 3.952)	1.446 (-2.977, 5.868)

	Balance (n=80)		
Compositional Isotemporal Substitutions of 24h-MB	**10 min**	**20 min**	**30 min**
↑ MVPA ↓ Sleep	0.020 (-0.186, 0.225)	0.037 (-0.364, 0.438)	0.052 (-0.537, 0.641)
↑ MVPA ↓ SB	0.034 (-0.188, 0.255)	0.065 (-0.372, 0.502)	0.094 (-0.552, 0.740)
↑ MVPA ↓ LPA	0.161 (-0.354, 0.676)	0.327 (-0.715, 1.369)	0.499 (-1.087, 2.084)
↑ Sleep ↓ MVPA	-0.023 (-0.238, 0.193)	-0.048 (-0.492, 0.396)	-0.077 (-0.764, 0.609)
↑ Sleep ↓ SB	0.014 (-0.170, 0.197)	0.027 (-0.343, 0.396)	0.039 (-0.517, 0.596)
↑ Sleep ↓ LPA	0.141 (-0.265, 0.547)	0.289 (-0.545, 1.122)	0.444 (-0.840, 1.728)
↑ SB ↓ MVPA	-0.036 (-0.266, 0.194)	-0.075 (-0.545, 0.394)	-0.118 (-0.839, 0.603)
↑ SB ↓ Sleep	-0.014 (-0.196, 0.168)	-0.028 (-0.391, 0.334)	-0.043 (-0.584, 0.498)
↑ SB ↓ LPA	0.127 (-0.346, 0.601)	0.261 (-0.705, 1.227)	0.403 (-1.077, 1.884)
↑ LPA ↓ MVPA	-0.157 (-0.663, 0.348)	-0.312 (-1.317, 0.694)	-0.464 (-1.967, 1.039)
↑ LPA ↓ Sleep	-0.135 (-0.522, 0.252)	-0.265 (-1.022, 0.493)	-0.390 (-1.503, 0.723)
↑ LPA ↓ SB	-0.121 (-0.578, 0.335)	-0.237 (-1.135, 0.661)	-0.347 (-1.674, 0.979)

	Locomotion (n=80)		
Compositional Isotemporal Substitutions of 24h-MB	**10 min**	**20 min**	**30 min**
↑ MVPA ↓ Sleep	-0.075 (-0.153, 0.003)	-0.148 (-0.302, 0.006)	-0.218 (-0.447, 0.010)

↑ MVPA ↓ SB	-0.058 (-0.144, 0.028)	-0.112 (-0.283, 0.058)	-0.164 (-0.418, 0.090)
↑ MVPA ↓ LPA	-0.031 (-0.208, 0.146)	-0.058 (-0.418, 0.302)	-0.080 (-0.630, 0.471)
↑ Sleep ↓ MVPA	0.078 (-0.002, 0.159)	0.160 (-0.003, 0.324)	0.247 (-0.004, 0.497)
↑ Sleep ↓ SB	0.017 (-0.079, 0.114)	0.035 (-0.158, 0.227)	0.051 (-0.239, 0.342)
↑ Sleep ↓ LPA	0.044 (-0.135, 0.223)	0.089 (-0.277, 0.455)	0.136 (-0.426, 0.697)
↑ SB ↓ MVPA	0.061 (-0.027, 0.149)	0.126 (-0.053, 0.304)	0.195 (-0.077, 0.466)
↑ SB ↓ Sleep	-0.018 (-0.114, 0.078)	-0.036 (-0.227, 0.155)	-0.055 (-0.341, 0.231)
↑ SB ↓ LPA	0.027 (-0.146, 0.199)	0.054 (-0.298, 0.407)	0.084 (-0.458, 0.625)
↑ LPA ↓ MVPA	0.036 (-0.136, 0.208)	0.076 (-0.264, 0.417)	0.122 (-0.384, 0.628)
↑ LPA ↓ Sleep	-0.043 (-0.215, 0.129)	-0.086 (-0.424, 0.253)	-0.127 (-0.626, 0.372)
↑ LPA ↓ SB	-0.025 (-0.192, 0.141)	-0.050 (-0.376, 0.277)	-0.073 (-0.555, 0.409)

	Fine Motor Function (n=79)		
Compositional Isotemporal Substitutions of 24h-MB	**10 min**	**20 min**	**30 min**
↑ MVPA ↓ Sleep	-0.014 (-0.377, 0.348)	-0.019 (-0.731, 0.693)	-0.015 (-1.065, 1.034)
↑ MVPA ↓ SB	**-0.337** (-0.650, -0.024)	**-0.671** (-1.285, -0.056)	**-1.003** (-1.909, -0.096)
↑ MVPA ↓ LPA	-0.176 (-0.921, 0.569)	-0.346 (-1.854, 1.161)	-0.513 (-2.806, 1.780)
↑ Sleep ↓ MVPA	0.025 (-0.353, 0.403)	0.061 (-0.713, 0.835)	0.110 (-1.080, 1.301)
↑ Sleep ↓ SB	**-0.321** (-0.623, -0.019)	**-0.644** (-1.250, -0.038)	**-0.970** (-1.882, -0.058)
↑ Sleep ↓ LPA	-0.159 (-0.777, 0.458)	-0.319 (-1.583, 0.944)	-0.480 (-2.423, 1.463)
↑ SB ↓ MVPA	**0.341** (0.015, 0.668)	**0.688** (0.019, 1.357)	**1.041** (0.010, 2.072)
↑ SB ↓ Sleep	**0.318** (0.018, 0.619)	**0.634** (0.035, 1.234)	**0.949** (0.051, 1.846)
↑ SB ↓ LPA	0.157 (-0.499, 0.813)	0.307 (-1.032, 1.647)	0.451 (-1.604, 2.507)
↑ LPA ↓ MVPA	0.182 (-0.551, 0.916)	0.372 (-1.087, 1.832)	0.572 (-1.612, 2.756)
↑ LPA ↓ Sleep	0.159 (-0.431, 0.750)	0.319 (-0.839, 1.477)	0.479 (-1.226, 2.184)

↑ LPA ↓ SB	-0.163 (-0.794, 0.468)	-0.333 (-1.573, 0.908)	-0.508 (-2.340, 1.323)

	Inhibitory Control (n=69)		
Compositional Isotemporal Substitutions of 24h-MB	**10 min**	**20 min**	**30 min**
↑ MVPA ↓ Sleep	**0.011** (0.000, 0.022)	**0.021** (0.000, 0.042)	**0.032** (0.001, 0.062)
↑ MVPA ↓ SB	0.003 (-0.009, 0.015)	0.006 (-0.018, 0.030)	0.009 (-0.027, 0.044)
↑ MVPA ↓ LPA	0.008 (-0.017, 0.033)	0.015 (-0.035, 0.066)	0.023 (-0.055, 0.100)
↑ Sleep ↓ MVPA	-0.011 (-0.022, 0.000)	-0.023 (-0.046, 0.000)	-0.035 (-0.070, 0.001)
↑ Sleep ↓ SB	-0.007 (-0.018, 0.003)	-0.015 (-0.035, 0.005)	-0.022 (-0.053, 0.008)
↑ Sleep ↓ LPA	-0.003 (-0.024, 0.018)	-0.006 (-0.049, 0.037)	-0.008 (-0.074, 0.058)
↑ SB ↓ MVPA	-0.004 (-0.016, 0.009)	-0.008 (-0.034, 0.018)	-0.013 (-0.053, 0.026)
↑ SB ↓ Sleep	0.007 (-0.003, 0.018)	0.015 (-0.005, 0.035)	0.022 (-0.008, 0.052)
↑ SB ↓ LPA	0.004 (-0.016, 0.025)	0.009 (-0.034, 0.051)	0.013 (-0.052, 0.078)
↑ LPA ↓ MVPA	-0.008 (-0.033, 0.017)	-0.017 (-0.066, 0.032)	-0.026 (-0.100, 0.048)
↑ LPA ↓ Sleep	0.003 (-0.017, 0.023)	0.006 (-0.033, 0.046)	0.010 (-0.049, 0.068)
↑ LPA ↓ SB	-0.004 (-0.024, 0.015)	-0.009 (-0.048, 0.030)	-0.013 (-0.071, 0.044)

	Working Memory (n=71)		
Compositional Isotemporal Substitutions of 24h-MB	**10 min**	**20 min**	**30 min**
↑ MVPA ↓ Sleep	0.045 (-0.114, 0.204)	0.081 (-0.231, 0.392)	0.109 (-0.350, 0.567)
↑ MVPA ↓ SB	**0.176** (0.024, 0.329)	**0.346** (0.047, 0.646)	**0.511** (0.069, 0.953)
↑ MVPA ↓ LPA	**0.441** (0.123, 0.759)	**0.893** (0.251, 1.535)	**1.359** (0.385, 2.332)
↑ Sleep ↓ MVPA	-0.054 (-0.220, 0.112)	-0.119 (-0.459, 0.222)	-0.195 (-0.720, 0.330)
↑ Sleep ↓ SB	0.130 (-0.003, 0.264)	0.260 (-0.008, 0.528)	0.390 (-0.014, 0.793)
↑ Sleep ↓ LPA	**0.395** (0.152, 0.638)	**0.807** (0.309, 1.304)	**1.238** (0.473, 2.003)

↑ SB ↓ MVPA	**-0.183** (-0.342, -0.024)	**-0.375** (-0.701, -0.049)	**-0.576** (-1.078, -0.075)
↑ SB ↓ Sleep	-0.131 (-0.264, 0.002)	-0.262 (-0.527, 0.003)	-0.394 (-0.791, 0.004)
↑ SB ↓ LPA	**0.266** (0.013, 0.518)	**0.550** (0.035, 1.066)	**0.856** (0.065, 1.648)
↑ LPA ↓ MVPA	**-0.433** (-0.748, -0.118)	**-0.861** (-1.489, -0.232)	**-1.285** (-2.229, -0.342)
↑ LPA ↓ Sleep	**-0.380** (-0.613, -0.148)	**-0.748** (-1.204, -0.291)	**-1.103** (-1.774, -0.432)
↑ LPA ↓ SB	**-0.249** (-0.491, -0.006)	**-0.482** (-0.959, -0.005)	-0.701 (-1.404, 0.003)

MVPA = moderate-to-vigorous physical activity; LPA = light physical activity; SB = sedentary behavior. Confidence intervals were estimated using HC1 robust standard errors. Bold coefficients are statistically significant at $p < 0.05$. Values represent mean changes adjusted for the preschooler's sex and age, socioeconomic-status category, locality (urban/rural), and BMI z-score. n in each heading is the number of participants with an observed outcome; n_model in the accompanying long file is the complete-case analytic sample.