

Ch-06 R Codes

Ping-Yang Chen

Textbook: Montgomery, D. C. (2012). *Design and analysis of experiments*, 8th Edition. John Wiley & Sons.

Online handouts: https://github.com/PingYangChen/ANOVA_Course_R_Code

2²-Design: Chemical Process Example

Read the csv file 6_ChemicalRecovery.csv in R.

```
df5 <- read.csv(file.path("data", "6_ChemicalRecovery.csv"))
```

The estimation of factor effects are:

```
# Effect of factor A (reactant concentration)
mean(df5$Recovery[df5$ReactConc == 1] - df5$Recovery[df5$ReactConc == -1])
# or 2*mean(df5$Recovery*df5$ReactConc)

# Effect of factor A (catalyst amount)
mean(df5$Recovery[df5$CataAmo == 1] - df5$Recovery[df5$CataAmo == -1])

# Effect of two-factor interaction AB
twofi <- as.numeric(df5$ReactConc)*df5$CataAmo
mean(df5$Recovery[twofi == 1] - df5$Recovery[twofi == -1])
```

```
## Effect of factor A (reactant concentration): 8.3333
```

```
## Effect of factor A (catalyst amount): -5.0000
```

```
## Effect of two-factor interaction AB: 1.6667
```

Use `aov()` to fit the ANOVA model (model description in math is omitted). Here, if the columns of the factor in the `data.frame` are not of the `factor` type. We can also specify the type of the factors as `factor` in the R model formula.

```
fit5 <- aov(Recovery ~ factor(ReactConc)*factor(CataAmo), data = df5)
summary(fit5)
```

```
##                                Df Sum Sq Mean Sq F value    Pr(>F)
## factor(ReactConc)              1  208.33   208.33   53.191 8.44e-05 ***
## factor(CataAmo)                1   75.00    75.00   19.149  0.00236 **
## factor(ReactConc):factor(CataAmo) 1    8.33     8.33    2.128  0.18278
## Residuals                      8   31.33     3.92
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

The two main effects are significant and their two-factor interaction may not exist.


To visualize the analysis result of the factorial experiment, interaction plot is commonly used tool.

```
interaction.plot(
  x.factor = df5$ReactConc, # x-axis variable
```

```

trace.factor = df5$CataAmo, # variable for lines
response = df5$Recovery, # y-axis variable
ylab = "Recovery", xlab = "Reactant Concentration",
col = c("firebrick", "dodgerblue2"),
lty = 1, lwd = 3, trace.label = "Catalyst\nAmount"
)

```



2³-Design: Plasma Etching Example

Read the csv file 6_PlasmaEtching_2³.csv in R.

```
df6 <- read.csv(file.path("data", "6_PlasmaEtching_2^3.csv"))
```

The estimation of factor effects are:

```

# Effect of main effects
2*mean(df6$EachRate*df6$Gap)
2*mean(df6$EachRate*df6$GasFlow)
2*mean(df6$EachRate*df6$Power)

# Effect of two-factor interactions
GapGasFlow <- df6$Gap * df6$GasFlow
GapPower <- df6$Gap * df6$Power
GasFlowPower <- df6$GasFlow * df6$Power
2*mean(df6$EachRate*GapGasFlow)
2*mean(df6$EachRate*GapPower)
2*mean(df6$EachRate*GasFlowPower)

# Effect of three-factor interaction
GapGasFlowPower <- df6$Gap * df6$GasFlow * df6$Power

```

```
2*mean(df6$EachRate*GapGasFlowPower)
```

```
##   Factor Est.Effect
## 1      A    -101.625
## 2      B      7.375
## 3      C    306.125
## 4     AB    -24.875
## 5     AC   -153.625
## 6     BC     -2.125
## 7     ABC      5.625
```

Use `aov()` to fit the ANOVA model (model description in math is omitted). Here, if the columns of the factor in the `data.frame` are not of the `factor` type. We can also specify the type of the factors as `factor` in the R model formula.

```
fit6 <- aov(
  EachRate ~ factor(Gap) * factor(GasFlow) * factor(Power),
  data = df6
)
summary(fit6)
```

```
##               Df Sum Sq Mean Sq F value    Pr(>F)
## factor(Gap)      1  41311    41311   18.339 0.002679
## factor(GasFlow)   1    218      218    0.097 0.763911
## factor(Power)     1 374850   374850  166.411 1.23e-06
## factor(Gap):factor(GasFlow)  1    2475    2475    1.099 0.325168
## factor(Gap):factor(Power)    1   94403   94403   41.909 0.000193
## factor(GasFlow):factor(Power)  1     18      18    0.008 0.930849
## factor(Gap):factor(GasFlow):factor(Power)  1    127    127    0.056 0.818586
## Residuals        8   18020    2253
##
## factor(Gap)                **
## factor(GasFlow)
## factor(Power)              ***
## factor(Gap):factor(GasFlow)
## factor(Gap):factor(Power)   ***
## factor(GasFlow):factor(Power)
## factor(Gap):factor(GasFlow):factor(Power)
## Residuals
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

The significant effects are main effects `Gap` and `Power`, and, the two-factor interaction `Gap:Power`.

WE can also fit a response surface model and obtain the same conclusion.

```
ols6_full <- lm(EachRate ~ Gap * GasFlow * Power, data = df6)
summary(ols6_full)
```

```
##
## Call:
## lm(formula = EachRate ~ Gap * GasFlow * Power, data = df6)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -65.50 -11.12   0.00  11.12  65.50
##
```

```
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)    776.062    11.865   65.406 3.32e-12 ***
## Gap           -50.812    11.865   -4.282 0.002679 **
## GasFlow         3.688    11.865    0.311 0.763911
## Power          153.062    11.865   12.900 1.23e-06 ***
## Gap:GasFlow    -12.438    11.865   -1.048 0.325168
## Gap:Power      -76.812    11.865   -6.474 0.000193 ***
## GasFlow:Power  -1.062    11.865   -0.090 0.930849
## Gap:GasFlow:Power 2.813    11.865    0.237 0.818586
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 47.46 on 8 degrees of freedom
## Multiple R-squared:  0.9661, Adjusted R-squared:  0.9364
## F-statistic: 32.56 on 7 and 8 DF,  p-value: 2.896e-05
```

The final RSM is the reduced model that those insignificant terms are removed from the full model. Now we can use the real value of the factor levels as the predictors' values from this reduced model for better interpretation.

```
ols6_reduced <- lm(EachRate ~ NumGap * NumPower, data = df6)
summary(ols6_reduced)


##
## Call:
## lm(formula = EachRate ~ NumGap * NumPower, data = df6)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -72.50 -15.44   2.50  18.69  66.50
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  -2461.188    349.513   -7.042 1.35e-05 ***
## NumGap       2242.344    342.725    6.543 2.76e-05 ***
## NumPower      10.743      1.063   10.107 3.19e-07 ***
## NumGap:NumPower -7.681      1.042   -7.370 8.62e-06 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 41.69 on 12 degrees of freedom
## Multiple R-squared:  0.9608, Adjusted R-squared:  0.9509
## F-statistic: 97.91 on 3 and 12 DF,  p-value: 1.054e-08

x1_grid <- seq(min(df6$NumGap), max(df6$NumGap), length = 100)
x2_grid <- seq(min(df6$NumPower), max(df6$NumPower), length = 100)
newx <- data.frame(
  NumGap = rep(x1_grid, each = 100),
  NumPower = rep(x2_grid, time = 100)
)

rsplot_data6 <- data.frame(newx, rs = predict(ols6_reduced, newx))

library(ggplot2)
#- Add color to the contour plot
```

```
ggplot(rsplot_data6) +
  geom_contour(aes(NumGap, NumPower, z = rs), colour = "white") +
  geom_contour_filled(aes(NumGap, NumPower, z = rs))
```



Unreplicated 2^k Factorial Designs: The Resin Plant Experiment

Read the csv file 6_PilotPlant.csv in R.

```
df7 <- read.csv(file.path("data", "6_PilotPlant.csv"))
```

The estimation of factor effects are:

```
# Compute the model matrix of all effect terms without intercept
mmat7 <- model.matrix( ~ Temperature*Pressure*CH20Conc*StirRate - 1, data = df7)
# Calculate the effect sizes using the +/- signs of the model matrix
eff7 <- numeric(ncol(mmat7))
for (i in 1:ncol(mmat7)) {
  eff7[i] <- 2*mean(df7$FiltrationRate*mmat7[,i])
}
names(eff7) <- colnames(mmat7)
```

##	Factor	Est.Effect
## 1	Temperature	21.625
## 2	Pressure	3.125
## 3	CH20Conc	9.875


```
## 4          StirRate      14.625
## 5      Temperature:Pressure      0.125
## 6      Temperature:CH20Conc     -18.125
## 7          Pressure:CH20Conc      2.375
## 8      Temperature:StirRate      16.625
## 9          Pressure:StirRate     -0.375
## 10         CH20Conc:StirRate     -1.125
## 11      Temperature:Pressure:CH20Conc      1.875
## 12      Temperature:Pressure:StirRate      4.125
## 13      Temperature:CH20Conc:StirRate     -1.625
## 14          Pressure:CH20Conc:StirRate     -2.625
## 15 Temperature:Pressure:CH20Conc:StirRate      1.375
```

Because there is no replicates in each treatment combination, the estimate of the random error σ^2 does not exist and the ANOVA table is not available. Of all effect terms, we try of eliminate some of them before analyzing the data. According to the model assumption, the effects that are negligible should be similar to random error which is normally distributed with zero mean and constant variance. Therefore, a QQ-plot or half-Normal plot is helpful to identify effective effects. Belows are the codes of half-Normal plot.

```
# Half Normal Plot
halfqqnorm <- function(input, tol = 0.5) {
  y <- sort(abs(input))
  nq <- qnorm(seq(0.5, 0.99, length = length(y)))
  plot(y, nq, yaxt = "n", pch = 15,
       xlab = "|Effects|", ylab = "Half Normal Probability")
  title("Half Normal Plot")
  # choose anchor point to draw a straight line
  s <- min(which(diff(y)/diff(range(y)) > 1/(length(y)-1)))
  abline(a = -y[1]*(nq[s]-nq[1])/(y[s]-y[1]), b = (nq[s]-nq[1])/(y[s]-y[1]))
  axis(2, at = qnorm(seq(0.5, 0.9999, length = 5)),
       labels = round(seq(0, 1, length = 5),2))
  loc <- sqrt((nq - (y - y[1])*(nq[s]-nq[1])/(y[s]-y[1]))^2) > tol
  if (is.null(names(y))) {
    text(y[loc], nq[loc], order(abs(input))[loc], pos = 2)
  } else {
    text(y[loc], nq[loc], names(abs(input))[order(abs(input))[loc]], pos = 2)
  }
}
```

```
halfqqnorm(eff7)
```

Half Normal Plot



By the half Normal plot, we find out that the main effects **Temperature**, **CH2OConc** and **StirRate** and interactions **Temperature:CH2OConc**, **Temperature:StirRate** appear to be large.

Based on the observation from the half Normal plot, now the ANOVA model is

$$y_{ijkl} = \mu + \tau_i + \beta_j + \gamma_k + (\tau\beta)_{ij} + (\tau\gamma)_{ik} + \varepsilon_{ijkl} \quad (1)$$

- τ_i is the effect of the i th **Temperature** level, $i = 1, 2$.
- β_j is the effect of the j th **CH2OConc** level, $j = 1, 2$.
- γ_k is the effect of the k th **StirRate** level, $j = 1, 2$.
- $(\tau\beta)_{ij}$ is the interaction effect of the i th **Temperature** level and the j th **CH2OConc** level.
- $(\tau\gamma)_{ik}$ is the interaction effect of the i th **Temperature** level and the k th **StirRate** level.
- ε_{ijkl} is the random error, $l = 1, 2$, satisfying

$$\varepsilon_{ijkl} \stackrel{i.i.d.}{\sim} N(0, \sigma^2) \text{ where } \sigma^2 \text{ is the conatnt variance.}$$

Use `aov()` to fit the ANOVA model.

```
fit7 <- aov(
  FiltrationRate ~ factor(Temperature) * (factor(CH2OConc) + factor(StirRate)),
  data = df7
)
summary(fit7)
```

	Df	Sum Sq	Mean Sq	F value	Pr(>F)	
## factor(Temperature)	1	1870.6	1870.6	95.86	1.93e-06	***
## factor(CH2OConc)	1	390.1	390.1	19.99	0.0012	**
## factor(StirRate)	1	855.6	855.6	43.85	5.92e-05	***

```
## factor(Temperature):factor(CH20Conc)  1 1314.1  1314.1   67.34 9.41e-06 ***
## factor(Temperature):factor(StirRate)  1 1105.6  1105.6   56.66 2.00e-05 ***
## Residuals                            10  195.1    19.5
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Here, we left the interpretation of the response surface model for practice.

Addition of Center Points to a 2^k Designs

Recall the Resin Plant Experiment. Read the csv file 6_PilotPlant.csv in R.

```
df7 <- read.csv(file.path("data", "6_PilotPlant.csv"))
```

```
df7_C <- data.frame(
  Temperature = rep(0, 4),
  Pressure = rep(0, 4),
  CH20Conc = rep(0, 4),
  StirRate = rep(0, 4),
  FiltrationRate = c(73, 75, 66, 69)
)
```

The calculation of $SS_{Purequadratic}$ with degree of freedom 1.

```
Yf_bar <- mean(df7$FiltrationRate)
Yc_bar <- mean(df7_C$FiltrationRate)
nf <- nrow(df7)
nc <- nrow(df7_C)
SS_pureQuad <- nf*nc*(Yf_bar - Yc_bar)^2/(nf + nc)
```

The calculation of SS_E with degree of freedom $n_c - 1$.

```
SS_E <- sum((df7_C$FiltrationRate - mean(df7_C$FiltrationRate))^2)
```

To test the significance of the Curvature, we compute the ratio of $MS_{Purequadratic}$ and MS_E as the F -statistic which follows the F distribution with degrees of freedom 1 and $n_c - 1$.

```
MS_pureQuad <- SS_pureQuad/1
MS_E <- SS_E/(nc - 1)
fval <- MS_pureQuad/MS_E
pval <- 1 - pf(fval, 1, nc - 1) # p-value
```

The part of the testing the significance of the Curvature of the ANOVA table.

```
print(data.frame(
  Source = c("Curvature", "Residual"),
  SS = c(SS_pureQuad, SS_E),
  DF = c(1, nc - 1),
  MS = c(MS_pureQuad, MS_E),
  "F" = c(sprintf("%.3f", fval), ""),
  "Pr(>F)" = c(sprintf("%.3f", pval), "")
))
```

```
##      Source      SS DF      MS      F Pr..F.
## 1 Curvature  1.5125  1  1.5125 0.093  0.780
## 2 Residual 48.7500  3 16.2500
```