Α				В			
Symptom frequency				Symptom frequency			
AT, survey study, % cohort				IT, survey study, % cohort			
Fever -	O 56%	• 3.8%	• 0.63%	Fever-	71%	• 3%	• 0.7%
Sore throat	O 50%	• 4%	• 1.7%	Shivering -	51%	• 2.3%	• 1.2%
Running nose -	O 43%	• 2.7%	• 1.3%	Sore throat		4.2%	• 1.2%
Shivering -	O 43%	• 0.63%	• 0.42%	Running nose		• 5.2%	• 1.6%
				_	T	0.270	
Tachypnea-	O 59%	O 37%	O 25%	Tachypnea.		O 30%	O 21%
Dry cough -	O 56%	17%	• 5.8%	Dry cough	1 Y	17%	• 5.4%
Dyspnea -	O 38%	o 17%	11%	Dyspnea -		• 14%	• 7.5%
Wet cough -	O 18%	• 4%	• 1.7%	Wet cough ·	• 13%	• 3%	• 0.47%
Joint pain -	O 66%	o 15%	• 9.4%	Joint pain ·	73%	O 22%	o 12%
Chest pain -	O 49%	0 21%	• 14%	Muscle pain	I •	0 22%	• 12%
Muscle pain -	0 58%	• 15%	• 9%	Bone pain	J	• 18%	• 9.6%
Bone pain -	O 44%	• 12%	• 7.5%	Chest pain		• 17%	• 7.7%
Abdominal pain -	O 22%	• 5.8%	• 2.3%	Abdominal pain		• 7.3%	• 2.8%
:				•			
Dim. appetite -	O 64%	• 11%	• 2.3%	Dim. appetite	1 Y	• 11%	• 2.8%
Diarrhea -	O 31%	• 4.8%	• 2.7%	Diarrhea ·	O 34%	• 4.9%	• 1.6%
Nausea -	O 27%	• 5.6%	• 2.1%	Nausea -		• 4.7%	• 2.3%
Vomiting -	• 6.1%	• 0.63%	• 0.21%	Vomiting -	• 8.7%	• 0.7%	• 0.23%
Tachycardia-	O 33%	o 16%	• 11%	Tachycardia-	0 26%	• 13%	• 8.2%
Palpitations -	O 17%	• 11%	• 8.6%	Palpitations -		9.4%	• 7.3%
		2.00/		Dad ayes	1 0000	100/	
Red eyes -	O 19%	• 6.9%	• 3.5%	Red eyes	T .	• 12%	• 6.1%
Swelling -	• 4.4%	• 2.7%	• 2.5%	Swelling -		• 4.7%	• 4%
Urticaria -	• 4.8%	• 1.7%	• 1.3%	Urticaria ·	• 8%	• 3.7%	• 3%
Blistering rash -	• 3.3%	• 1.7%	• 1%	Blistering rash	• 5.4%	• 3%	• 1.6%
Marmorated skin -	• 2.1%	• 1.5%	• 1.3%	Blue fingers/toes		• 1.6%	• 1.2%
Blue fingers/toes -	• 1.3%	• 0.84%	• 0.84%	Marmorated skin -	• 1.9%	• 1.4%	• 0.94%
Hypo/anosmia -	70%	O 40%	O 30%	Hypo/anosmia -	75%	O 39%	O 27%
Hypo/ageusia -	68%	O 36%	O 23%	Hypo/ageusia		O 36%	0 21%
Entique I	-040/	450/	070/	Entique	I — 000/	<u></u>	<u> </u>
Fatigue -	94%	O 45%	O 27%	Fatigue ·		O 54%	O 33%
Tiredness at day -	84%	O 46%	O 32%	Tiredness at day		O 49%	O 33%
Sleeplessness -	O 39%	O 19%	• 14%	Sleeplessness -	O 31%	o 16%	• 11%
Imp. concentration -	O 50%	O 32%	O 22%	Imp. concentration -		O 33%	O 27%
Forgetfulness -	O 32%	O 23%	18%	Forgetfulness -	O 35%	O 29%	O 26%
Dizziness -	O 46%	• 11%	• 6.9%	Confusion -	0 26%	17%	• 11%
Confusion -	O 18%	• 9.2%	• 5.6%	Dizziness ·		• 12%	• 5.6%
Tingling feet -	o 13%	• 5.2%	• 4.8%	Imp. walk	0 15%	• 9.8%	• 6.6%
Imp. walk	• 12%	• 4.6%	• 2.5%	Tingling hands		• 8.4%	• 6.8%
Tingling hands -							
	• 8.6% • 6.1%	• 3.5%	• 2.9%	Tingling feet		• 6.8%	• 4.9%
Numb feet - Burning feet -	• 6.1% • 5.0%	• 3.3%	• 2.9%	Numb hands · Numb feet ·		• 7%	• 5.6%
•	• 5.8%	• 2.9%	• 2.9%			• 6.6% • 4.3%	• 5.2%
Numb hands -	• 4.8%	• 3.1%	• 2.9%	Burning feet	1	4.2%	• 3.5%
Imp. FMS -	• 4.6%	• 1%	• 0.63%	Burning hands	• 4.7%	• 3%	• 2.1%
Burning hands -	• 1.7%	• 1.5%	• 1.5%	Imp. FMS	• 4.4%	• 2.8%	• 2.3%
	0 - 14 days 2	28 days 3	3 months		0 - 14 days	28 days	3 months

n = 427

n = 479