Α				В			
Symptom frequency				Symptom frequency			
AT, HACT study, % cohort				IT, HACT study, % cohort			
Fever -	O 56%	• 3.8%	• 0.63%	Fever -	71%	• 3%	• 0.7%
Sore throat -	O 50%	• 4%	• 1.7%	Shivering -	O 51%	• 2.3%	• 1.2%
Running nose -	O 43%	• 2.7%	• 1.3%	Sore throat -	O 43%	• 4.2%	• 1.2%
Shivering -	O 42%	0.63%	• 0.42%	Running nose -	O 41%	• 5.2%	• 1.6%
Tachypnea	<u></u> 59%	O 37%	O 25%	Tachypnea-	53%	30%	O 21%
Dry cough -	0 56%	0 17%	• 5.8%	Dry cough -	O 49%	• 17%	• 5.4%
Dyspnea -	O 38%	0 17%	• 5.6% • 11%	Dyspnea -	O 49% O 31%	• 14%	
Wet cough -	Ī						• 7.5%
Wei cough 7	O 18%	• 4%	• 1.7%	Wet cough -	• 13%	• 3%	• 0.47%
Joint pain -	<u></u> 66%	o 15%	• 9.4%	Joint pain -	73%	O 22%	• 12%
Chest pain -	Q 49%	21%	14%	Muscle pain -	O 68%	O 22%	o 12%
Muscle pain -	O 58%	15%	• 9%	Bone pain -	O 64%	o 18%	• 9.6%
Bone pain -	O 44%	• 12%	• 7.5%	Chest pain -	O 42%	O 17%	• 7.7%
Abdominal pain -	O 22%	• 5.8%	• 2.3%	Abdominal pain -	O 26%	• 7.3%	• 2.8%
Dim. appetite -	O 64%	o 11%	• 2.3%	Dim. appetite -	O 61%	• 11%	• 2.8%
Diarrhea -	O 31%	• 4.8%	• 2.7%	Diarrhea -	O 34%	• 4.9%	• 1.6%
Nausea -	0 27%	• 5.6%	• 2.1%	Nausea -	O 31%	• 4.7%	• 2.3%
Vomiting -	• 6.1%	• 0.63%	0.21%	Vomiting -		• 0.7%	0.23%
Tachycardia-	O 33%	o 16%	• 11%	Tachycardia-	O 26%	• 13%	• 8.2%
Palpitations -	• 17%	• 11%	• 8.6%	Palpitations -	o 17%	• 9.4%	• 7.3%
Red eyes -	o 19%	• 6.9%	• 3.5%	Red eyes -	O 31%	• 12%	• 6.1%
Swelling -	• 4.4%	• 2.7%	• 2.5%	Swelling -	• 6.3%	• 4.7%	• 4%
Urticaria -	• 4.8%	• 1.7%	• 1.3%	Urticaria -	• 8%	• 3.7%	• 3%
Blistering rash -	• 3.3%	• 1.7%	• 1%	Blistering rash -	• 5.4%	• 3%	• 1.6%
Marmorated skin -	• 2.1%	• 1.5%	• 1.3%	Blue fingers/toes -	• 2.1%	• 1.6%	• 1.2%
Blue fingers/toes -	• 1.3%	• 0.84%	• 0.84%	Marmorated skin -	• 1.9%	• 1.4%	• 0.94%
Hypo/anosmia -	70%	O 40%	O 30%	Hypo/anosmia -	75%	O 39%	O 27%
Hypo/ageusia -	68%	O 36%	0 23%	Hypo/ageusia -		O 36%	0 21%
!	-				-	ī	
Fatigue -	94%	O 45%	O 27%	Fatigue -	92%	O 54%	O 33%
Tiredness at day -	84%	O 46%	O 32%	Tiredness at day -	79%	O 49%	O 33%
Sleeplessness -	O 39%	• 19%	• 14%	Sleeplessness -	O 31%	o 16%	• 11%
Imp. concentration	O 50%	O 32%	O 22%	Imp. concentration -	O 49%	O 33%	O 27%
Forgetfulness -	O 32%	O 23%	18%	Forgetfulness -	O 35%	O 29%	O 26%
Dizziness -	O 46%	• 11%	• 6.9%	Confusion -	O 26%	o 17%	• 11%
Confusion -	o 18%	• 9.2%	• 5.6%	Dizziness -	O 31%	• 12%	• 5.6%
Tingling feet	• 13%	• 5.2%	• 4.8%	Imp. walk -	o 15%	9.8%	• 6.6%
Imp. walk -	• 12%	• 4.6%	• 2.5%	Tingling hands -	• 13%	• 8.4%	• 6.8%
Tingling hands -	• 8.6%	• 3.5%	• 2.9%	Tingling feet -	• 13%	• 6.8%	• 4.9%
Numb feet -	• 6.1%	• 3.3%	• 2.9%	Numb hands -	• 9.4%	• 7%	• 5.6%
Burning feet	• 5.8%	• 2.9%	• 2.9%	Numb feet -	• 10%	• 6.6%	• 5.2%
Numb hands -	• 4.8%	• 3.1%	• 2.9%	Burning feet -	• 7%	• 4.2%	• 3.5%
Imp. FMS	• 4.6%	• 1%	• 0.63%	Burning hands -	• 4.7%	• 3%	• 2.1%
Burning hands -	• 1.7%	• 1.5%	• 1.5%	Imp. FMS -	• 4.4%	• 2.8%	• 2.3%
Dulling names L		_		11111-1 1410-1		-	
	Acute CoV Ic	ong CoV	PASC		Acute CoV	long CoV	PASC

n = 427

n = 479