	cymptom requercy, A1, survey study					cymptom requency, 11, survey study		
survey study, % of cohort, n = 479					survey study, % of cohort, n = 427			
Fever-	5 6%	• 3.8%	• 0.63%	Fever	1	• 3%	• 0.7%	
Sore throat -	5 0%	• 4%	• 1.7%	Shivering	T .	• 2.3%	• 1.2%	
Running nose -	43 %	• 2.7%	• 1.3%	Sore throat	1	• 4.2%	• 1.2%	
Shivering -	42 %	• 0.63%	• 0.42%	Running nose	4 1%	• 5.2%	• 1.6%	
Tachypnea-	· • • • • • • • • • • • • • • • • • • •	37 %	2 5%	Tachypnea	T	3 0%	2 1%	
Dry cough -	O 56%	17%	• 5.8%	Dry cough	49%	• 17%	• 5.4%	
Dyspnea -	38 %	17%	• 11%	Dyspnea	T	• 14%	• 7.5%	
Wet cough -	• 18%	• 4%	• 1.7%	Wet cough	• 13%	• 3%	• 0.47%	
Joint pain -	66%	• 15%	• 9.4%	Joint pain	·	• 22%	• 12%	
Chest pain -	4 9%	21%	• 14%	Muscle pain	1 2	22 %	• 12%	
Muscle pain -	5 8%	15%	• 9%	Bone pain	1 Y	• 18%	• 9.6%	
Bone pain -	4 4%	• 12%	• 7.5%	Chest pain	¥ .=	• 17%	• 7.7%	
Abdominal pain -	2 2%	• 5.8%	• 2.3%	Abdominal pain	• 26%	• 7.3%	• 2.8%	
Dim. appetite -	64%	• 11%	• 2.3%	Dim. appetite	1 Y	• 11%	• 2.8%	
Diarrhea -	3 1%	• 4.8%	• 2.7%	Diarrhea -	1 7 7	• 4.9%	• 1.6%	
Nausea -	27 %	• 5.6%	• 2.1%	Nausea	¥ 0.70	• 4.7%	• 2.3%	
Vomiting -	• 6.1%	• 0.63%	• 0.21%	Vomiting -	• 8.7%	• 0.7%	• 0.23%	
Tachycardia-		• 16%	• 11%	Tachycardia	1	• 13%	• 8.2%	
Palpitations -	• 17%	• 11%	• 8.6%	Palpitations -	• 17%	• 9.4%	• 7.3%	
Red eyes -	• 19%	• 6.9%	• 3.5%	Red eyes	1 7	• 12%	• 6.1%	
Swelling -	• 4.4%	• 2.7%	• 2.5%	Swelling -		• 4.7%	• 4%	
Urticaria -	• 4.8%	• 1.7%	• 1.3%	Urticaria -	• 8%	• 3.7%	• 3%	
Blistering rash -	• 3.3%	• 1.7%	• 1%	Blistering rash		• 3%	• 1.6%	
Marmorated skin -	• 2.1%	• 1.5%	• 1.3%	Blue fingers/toes		• 1.6%	• 1.2%	
Blue fingers/toes -	• 1.3%	• 0.84%	• 0.84%	Marmorated skin	• 1.9%	• 1.4%	• 0.94%	
OD -	70%	40 %	3 0%	OD :	1 2,0,0	3 9%	27 %	
Hypogeusia/ageusia -	68%	3 6%	23 %	Hypogeusia/ageusia		3 6%	2 1%	
Fatigue -	Q 94%	45 %	27 %	Fatigue	· ·	5 4%	33 %	
Tiredness at day -	8 4%	4 6%	32%	Tiredness at day	1 -	4 9%	33%	
Sleeplessness -	3 9%	• 19%	14%	Sleeplessness	• 31%	• 16%	• 11%	
Imp. concentration -	5 0%	32 %	22 %	Imp. concentration	T T	3 3%	27 %	
Forgetfulness -	32 %	23%	18%	Forgetfulness	1 7 77.1	2 9%	2 6%	
Dizziness -	4 6%	• 11%	• 6.9%	Confusion	T = * · ·	17%	11%	
Confusion -	• 18%	• 9.2%	• 5.6%	Dizziness	• 31%	• 12%	• 5.6%	
Tingling feet -	• 13%	• 5.2%	• 4.8%	Imp. walk		• 9.8%	• 6.6%	
Imp. walk -	• 12%	• 4.6%	• 2.5%	Tingling hands		• 8.4%	• 6.8%	
Tingling hands -	• 8.6%	• 3.5%	• 2.9%	Tingling feet		• 6.8%	• 4.9%	
Numb feet -	• 6.1%	• 3.3%	• 2.9%	Numb hands		• 7%	• 5.6%	
Burning feet -	• 5.8%	• 2.9%	• 2.9%	Numb feet		• 6.6%	• 5.2%	
Numb hands -	• 4.8%	• 3.1%	• 2.9%	Burning feet	1	• 4.2%	• 3.5%	
Imp. FMS -	• 4.6%	• 1%	• 0.63%	Burning hands		• 3%	• 2.1%	
Burning hands -	• 1.7%	• 1.5%	• 1.5%	Imp. FMS	• 4.4%	• 2.8%	• 2.3%	
-	0 - 14 days	28 days 3	months		0 - 14 days	28 days 3	3 months	

Symptom frequency, IT, survey study

Symptom frequency, AT, survey study