

Mathematics for Computer Science

Linear Algebra (Part 2)

Eigendecomposition

Karl Southern

Durham University

February 3rd, 2025

Thanks to Andrei Krokhin and William Moses for use of some slides.

Outline

- 1 Recap of Last Week & Plan for Today
- 2 Eigendecomposition - Motivation
- 3 Similarity of Matrices
- 4 Diagonalisation of Matrices
- 5 Eigendecomposition
- 6 Wrapping Things Up

Recap of Last Two Weeks

- For an $n \times n$ matrix, a **non-zero** vector $\mathbf{x} \in \mathbb{R}^n$ is called an **eigenvector** of A if $A\mathbf{x} = \lambda\mathbf{x}$.
- In this case, λ is called an **eigenvalue** of A , and \mathbf{x} is an **eigenvector corresponding to λ** .

Recap of Last Two Weeks

- For an $n \times n$ matrix, a **non-zero** vector $\mathbf{x} \in \mathbb{R}^n$ is called an **eigenvector** of A if $A\mathbf{x} = \lambda\mathbf{x}$.
- In this case, λ is called an **eigenvalue** of A , and \mathbf{x} is an **eigenvector corresponding to λ** .
- The polynomial $\det(\lambda I - A)$ is called the **characteristic polynomial** of A and the equation $\det(\lambda I - A) = 0$ the **characteristic equation** of A .
- The eigenvalues of A are the solutions of $\det(\lambda I - A) = 0$.

Recap of Last Two Weeks

- For an $n \times n$ matrix, a **non-zero** vector $\mathbf{x} \in \mathbb{R}^n$ is called an **eigenvector** of A if $A\mathbf{x} = \lambda\mathbf{x}$.
- In this case, λ is called an **eigenvalue** of A , and \mathbf{x} is an **eigenvector corresponding to λ** .
- The polynomial $\det(\lambda I - A)$ is called the **characteristic polynomial** of A and the equation $\det(\lambda I - A) = 0$ the **characteristic equation** of A .
- The eigenvalues of A are the solutions of $\det(\lambda I - A) = 0$.
- For an eigenvalue λ_0 of A , the null space of matrix $\lambda_0 I - A$ is the **eigenspace** of A corresponding to λ_0 . The non-zero vectors in this subspace are the eigenvectors of A corresponding to λ_0 .

Recap of Last Two Weeks

- For an $n \times n$ matrix, a **non-zero** vector $\mathbf{x} \in \mathbb{R}^n$ is called an **eigenvector** of A if $A\mathbf{x} = \lambda\mathbf{x}$.
- In this case, λ is called an **eigenvalue** of A , and \mathbf{x} is an **eigenvector corresponding to λ** .
- The polynomial $\det(\lambda I - A)$ is called the **characteristic polynomial** of A and the equation $\det(\lambda I - A) = 0$ the **characteristic equation** of A .
- The eigenvalues of A are the solutions of $\det(\lambda I - A) = 0$.
- For an eigenvalue λ_0 of A , the null space of matrix $\lambda_0 I - A$ is the **eigenspace** of A corresponding to λ_0 . The non-zero vectors in this subspace are the eigenvectors of A corresponding to λ_0 .
- Eigenvalues and eigenvectors play a major role in PCA (principal component analysis).

Plan for Today

- Similarity of matrices
- Diagonalisation of a matrix and how to find it
- Eigendecomposition of a matrix

Outline

- 1 Recap of Last Week & Plan for Today
- 2 Eigendecomposition - Motivation**
- 3 Similarity of Matrices
- 4 Diagonalisation of Matrices
- 5 Eigendecomposition
- 6 Wrapping Things Up

What Is It & Why Does It Matter?

What Is It?

① Decompose matrix A into PDP^{-1} form.

② P - eigenvectors, D - diagonal matrix eigenvalues, e.g., $D = \begin{pmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_3 \end{pmatrix}$

What Is It & Why Does It Matter?

What Is It?

① Decompose matrix A into PDP^{-1} form.

② P - eigenvectors, D - diagonal matrix eigenvalues, e.g., $D = \begin{pmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_3 \end{pmatrix}$

Why Care?

① Faster Self-Matrix Multiplication
 A^k

What Is It & Why Does It Matter?

What Is It?

① Decompose matrix A into PDP^{-1} form.

② P - eigenvectors, D - diagonal matrix eigenvalues, e.g., $D = \begin{pmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_3 \end{pmatrix}$

Why Care?

① Faster Self-Matrix Multiplication

$$A^k = PDP^{-1}PDP^{-1}PDP^{-1}PDP^{-1}\dots k \text{ times}$$

What Is It & Why Does It Matter?

What Is It?

① Decompose matrix A into PDP^{-1} form.

② P - eigenvectors, D - diagonal matrix eigenvalues, e.g., $D = \begin{pmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_3 \end{pmatrix}$

Why Care?

① Faster Self-Matrix Multiplication

$$A^k = PDP^{-1}PDP^{-1}PDP^{-1}PDP^{-1}\dots k \text{ times}$$

$$A^k = PD\text{DDDD}\dots k \text{ times } P^{-1}$$

What Is It & Why Does It Matter?

What Is It?

① Decompose matrix A into PDP^{-1} form.

② P - eigenvectors, D - diagonal matrix eigenvalues, e.g., $D = \begin{pmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_3 \end{pmatrix}$

Why Care?

① Faster Self-Matrix Multiplication

$$A^k = PDP^{-1}PDP^{-1}PDP^{-1}PDP^{-1}\dots k \text{ times}$$

$$A^k = PDDDD\dots k \text{ times } P^{-1}$$

$$A^k = PD^kP^{-1}$$

What Is It & Why Does It Matter?

What Is It?

① Decompose matrix A into PDP^{-1} form.

② P - eigenvectors, D - diagonal matrix eigenvalues, e.g., $D = \begin{pmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_3 \end{pmatrix}$

Why Care?

① Faster Self-Matrix Multiplication

$$A^k = PDP^{-1}PDP^{-1}PDP^{-1}PDP^{-1}\dots k \text{ times}$$

$$A^k = PDDDD\dots k \text{ times } P^{-1}$$

$$A^k = PD^kP^{-1} \text{ (e.g., } D^k = \begin{pmatrix} \lambda_1^k & 0 & 0 \\ 0 & \lambda_2^k & 0 \\ 0 & 0 & \lambda_3^k \end{pmatrix} \text{)}$$

What Is It & Why Does It Matter?

What Is It?

- 1 Decompose matrix A into PDP^{-1} form.

- 2 P - eigenvectors, D - diagonal matrix eigenvalues, e.g., $D = \begin{pmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_3 \end{pmatrix}$

Why Care?

- 1 Faster Self-Matrix Multiplication

$$A^k = PDP^{-1}PDP^{-1}PDP^{-1}PDP^{-1}\dots k \text{ times}$$

$$A^k = PDDDD\dots k \text{ times } P^{-1}$$

$$A^k = PD^kP^{-1} \text{ (e.g., } D^k = \begin{pmatrix} \lambda_1^k & 0 & 0 \\ 0 & \lambda_2^k & 0 \\ 0 & 0 & \lambda_3^k \end{pmatrix} \text{)}$$

- 2 Useful in PCA (covariance matrix)

Plan to Learn About It

- ① Learn useful concept 1: Similarity of matrices
- ② Learn useful concept 2: Diagonalisation of a matrix (what & how)
- ③ Finally get to eigendecomposition (what is it & conditions for it to exist)

Outline

- 1 Recap of Last Week & Plan for Today
- 2 Eigendecomposition - Motivation
- 3 Similarity of Matrices**
- 4 Diagonalisation of Matrices
- 5 Eigendecomposition
- 6 Wrapping Things Up

Similarity of Matrices

Definition

Square matrices A and B are called **similar** if $A = P^{-1}BP$ for some invertible P .

Note that if $A = P^{-1}BP$ then $B = Q^{-1}AQ$ where $Q = P^{-1}$.

Similarity of Matrices

Definition

Square matrices A and B are called **similar** if $A = P^{-1}BP$ for some invertible P .

Note that if $A = P^{-1}BP$ then $B = Q^{-1}AQ$ where $Q = P^{-1}$.

Similar matrices have many features in common, including determinant, trace, rank, nullity, characteristic polynomial, eigenvalues, dimensions of corresponding eigenspaces, etc.

Similarity of Matrices

Definition

Square matrices A and B are called **similar** if $A = P^{-1}BP$ for some invertible P .

Note that if $A = P^{-1}BP$ then $B = Q^{-1}AQ$ where $Q = P^{-1}$.

Similar matrices have many features in common, including determinant, trace, rank, nullity, characteristic polynomial, eigenvalues, dimensions of corresponding eigenspaces, etc.

Lemma

If A and B are similar then $\det(A) = \det(B)$.

Note: Whilst we have proven that A and B being similar implies that they have the same determinant, the converse is not true.

Properties of Similar Matrices

If A and B are similar then:

- ① A is invertible iff B is invertible.
- ② A and B have the same rank.
- ③ A and B have the same characteristic polynomial.
- ④ A and B have the same eigenvalues.

Recap: Representing a Matrix in Terms of a Basis

- 1 Suppose we have a vector $\mathbf{u} = \begin{pmatrix} 5 \\ 2 \end{pmatrix}$ in a coordinate system with the standard basis.

Recap: Representing a Matrix in Terms of a Basis

- 1 Suppose we have a vector $\mathbf{u} = \begin{pmatrix} 5 \\ 2 \end{pmatrix}$ in a coordinate system with the standard basis.
- 2 We want to represent it in terms of the basis $S = \{\mathbf{v}_1, \mathbf{v}_2\}$,
where $\mathbf{v}_1 = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$ and $\mathbf{v}_2 = \begin{pmatrix} 0 \\ 4 \end{pmatrix}$.

Recap: Representing a Matrix in Terms of a Basis

- 1 Suppose we have a vector $\mathbf{u} = \begin{pmatrix} 5 \\ 2 \end{pmatrix}$ in a coordinate system with the standard basis.
- 2 We want to represent it in terms of the basis $S = \{\mathbf{v}_1, \mathbf{v}_2\}$,
where $\mathbf{v}_1 = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$ and $\mathbf{v}_2 = \begin{pmatrix} 0 \\ 4 \end{pmatrix}$.
- 3 In other words, what are a, b such that $\mathbf{u} = a\mathbf{v}_1 + b\mathbf{v}_2$.

Recap: Representing a Matrix in Terms of a Basis

- 1 Suppose we have a vector $\mathbf{u} = \begin{pmatrix} 5 \\ 2 \end{pmatrix}$ in a coordinate system with the standard basis.
- 2 We want to represent it in terms of the basis $S = \{\mathbf{v}_1, \mathbf{v}_2\}$,
where $\mathbf{v}_1 = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$ and $\mathbf{v}_2 = \begin{pmatrix} 0 \\ 4 \end{pmatrix}$.
- 3 In other words, what are a, b such that $\mathbf{u} = a\mathbf{v}_1 + b\mathbf{v}_2$.
- 4 Solving, $a = 5, b = -2$.

Recap: Representing a Matrix in Terms of a Basis

- 1 Suppose we have a vector $\mathbf{u} = \begin{pmatrix} 5 \\ 2 \end{pmatrix}$ in a coordinate system with the standard basis.
- 2 We want to represent it in terms of the basis $S = \{\mathbf{v}_1, \mathbf{v}_2\}$,
where $\mathbf{v}_1 = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$ and $\mathbf{v}_2 = \begin{pmatrix} 0 \\ 4 \end{pmatrix}$.
- 3 In other words, what are a, b such that $\mathbf{u} = a\mathbf{v}_1 + b\mathbf{v}_2$.
- 4 Solving, $a = 5, b = -2$.
- 5 Here, a, b are the coordinates of \mathbf{u} in basis S and the vector $\begin{pmatrix} 5 \\ -2 \end{pmatrix}$ is the **coordinate vector** of vector \mathbf{u} in basis S .

Similarity and Linear Maps

Theorem

Matrices A and B are similar iff they represent the same linear operator $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$, possibly in different bases. (Proof omitted.)

Similarity and Linear Maps

Theorem

Matrices A and B are similar iff they represent the same linear operator $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$, possibly in different bases. (Proof omitted.)

- Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a linear operator and $S = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ a basis in \mathbb{R}^n .
- Let $[T]_S$ be the $n \times n$ matrix $[T]_S = [(T(\mathbf{v}_1))_S | (T(\mathbf{v}_2))_S | \dots | (T(\mathbf{v}_n))_S]$ whose columns are the coordinate vectors of vectors $T(\mathbf{v}_i)$ in basis S .

Similarity and Linear Maps

Theorem

Matrices A and B are similar iff they represent the same linear operator $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$, possibly in different bases. (Proof omitted.)

- Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a linear operator and $S = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ a basis in \mathbb{R}^n .
- Let $[T]_S$ be the $n \times n$ matrix $[T]_S = [(T(\mathbf{v}_1))_S | (T(\mathbf{v}_2))_S | \dots | (T(\mathbf{v}_n))_S]$ whose columns are the coordinate vectors of vectors $T(\mathbf{v}_i)$ in basis S .
This matrix is called the **matrix of T in S** .

Similarity and Linear Maps

Theorem

Matrices A and B are similar iff they represent the same linear operator $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$, possibly in different bases. (Proof omitted.)

- Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a linear operator and $S = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ a basis in \mathbb{R}^n .
- Let $[T]_S$ be the $n \times n$ matrix $[T]_S = [(T(\mathbf{v}_1))_S | (T(\mathbf{v}_2))_S | \dots | (T(\mathbf{v}_n))_S]$ whose columns are the coordinate vectors of vectors $T(\mathbf{v}_i)$ in basis S .
This matrix is called the **matrix of T in S** .
- We also say that the matrix $[T]_S$ **represents** T in basis S .

Similarity and Linear Maps

Theorem

Matrices A and B are similar iff they represent the same linear operator $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$, possibly in different bases. (Proof omitted.)

- Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a linear operator and $S = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ a basis in \mathbb{R}^n .
- Let $[T]_S$ be the $n \times n$ matrix $[T]_S = [(T(\mathbf{v}_1))_S | (T(\mathbf{v}_2))_S | \dots | (T(\mathbf{v}_n))_S]$ whose columns are the coordinate vectors of vectors $T(\mathbf{v}_i)$ in basis S .
This matrix is called the **matrix of T in S** .
- We also say that the matrix $[T]_S$ **represents** T in basis S .
- If S is the standard basis of \mathbb{R}^n then $[T]_S$ is the standard matrix of T .

Similarity and Linear Maps

Theorem

Matrices A and B are similar iff they represent the same linear operator $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$, possibly in different bases. (Proof omitted.)

- Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a linear operator and $S = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ a basis in \mathbb{R}^n .
- Let $[T]_S$ be the $n \times n$ matrix $[T]_S = [(T(\mathbf{v}_1))_S | (T(\mathbf{v}_2))_S | \dots | (T(\mathbf{v}_n))_S]$ whose columns are the coordinate vectors of vectors $T(\mathbf{v}_i)$ in basis S .
This matrix is called the **matrix of T in S** .
- We also say that the matrix $[T]_S$ **represents** T in basis S .
- If S is the standard basis of \mathbb{R}^n then $[T]_S$ is the standard matrix of T .

Theorem: Matrices A and B are similar iff there is a linear operator $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ and two bases S, S' of \mathbb{R}^n such that $A = [T]_S$ and $B = [T]_{S'}$.

Outline

- 1 Recap of Last Week & Plan for Today
- 2 Eigendecomposition - Motivation
- 3 Similarity of Matrices
- 4 Diagonalisation of Matrices**
- 5 Eigendecomposition
- 6 Wrapping Things Up

Diagonalisable Matrices

Definition

A matrix A is called **diagonalisable** if it is similar to a diagonal matrix – in other words, if there exists an invertible matrix P such that $P^{-1}AP$ is diagonal. Then P is said to **diagonalise** A .

Diagonalisable Matrices

Definition

A matrix A is called **diagonalisable** if it is similar to a diagonal matrix – in other words, if there exists an invertible matrix P such that $P^{-1}AP$ is diagonal. Then P is said to **diagonalise** A .

Note that A is diagonalisable if it decomposes as $A = PDP^{-1}$ where P is invertible and $D = \text{diag}(\lambda_1, \dots, \lambda_n)$ is diagonal.

$$D = \text{diag}(\lambda_1, \dots, \lambda_n) \text{ is shorthand for } D = \begin{pmatrix} \lambda_1 & 0 & . & . & . & 0 & 0 \\ 0 & \lambda_2 & . & . & . & 0 & 0 \\ . & . & . & . & . & . & . \\ . & . & . & . & . & . & . \\ . & . & . & . & . & . & . \\ 0 & 0 & . & . & . & \lambda_{n-1} & 0 \\ 0 & 0 & . & . & . & 0 & \lambda_n \end{pmatrix}$$

A Characterisation of Diagonalisable Matrices

Theorem (Thm 14.1)

An $n \times n$ matrix is diagonalisable iff it has n linearly independent eigenvectors.

A Characterisation of Diagonalisable Matrices

Theorem (Thm 14.1)

An $n \times n$ matrix is diagonalisable iff it has n linearly independent eigenvectors.

Proof.

(\Rightarrow). Assume that there is an invertible matrix P and a diagonal matrix $D = \text{diag}(\lambda_1, \dots, \lambda_n)$ such that $D = P^{-1}AP$, or $AP = PD$. Denote the column vectors of P by $\mathbf{p}_1, \dots, \mathbf{p}_n$, so $P = [\mathbf{p}_1 | \dots | \mathbf{p}_n]$. Then

$$AP = A[\mathbf{p}_1 | \dots | \mathbf{p}_n] = [A\mathbf{p}_1 | \dots | A\mathbf{p}_n].$$

On the other hand,

$$PD = [\lambda_1 \mathbf{p}_1 | \dots | \lambda_n \mathbf{p}_n].$$

Since $AP = PD$, we conclude that $A\mathbf{p}_i = \lambda_i \mathbf{p}_i$ for all $1 \leq i \leq n$.

Since P is invertible, its rank is n and so the vectors $\mathbf{p}_1, \dots, \mathbf{p}_n$ are linearly independent. Then none of $\mathbf{p}_1, \dots, \mathbf{p}_n$ is $\mathbf{0}$, so each of them is an eigenvector. □

Proof Cont'd

Theorem (Thm 14.1)

An $n \times n$ matrix is diagonalisable iff it has n linearly independent eigenvectors.

Proof.

(\Leftarrow). Assume that A has n linearly independent eigenvectors $\mathbf{p}_1, \dots, \mathbf{p}_n$ and let $\lambda_1, \dots, \lambda_n$ be the corresponding eigenvalues (not necessarily distinct). Define

$$P = [\mathbf{p}_1 | \dots | \mathbf{p}_n] \quad \text{and} \quad D = \text{diag}(\lambda_1, \dots, \lambda_n).$$

Then

$$AP = A[\mathbf{p}_1 | \dots | \mathbf{p}_n] = [A\mathbf{p}_1 | \dots | A\mathbf{p}_n] = [\lambda_1\mathbf{p}_1 | \dots | \lambda_n\mathbf{p}_n] = PD.$$

The columns of P are linearly independent, so its rank is n and it is invertible.

Finally $AP = PD$ is equivalent to $D = P^{-1}AP$. □

An Algorithm for Diagonalisation

Given a matrix A , this algorithm diagonalises it (or reports that this is impossible):

An Algorithm for Diagonalisation

Given a matrix A , this algorithm diagonalises it (or reports that this is impossible):

- 1 Find the eigenvalues of A (e.g. by solving its characteristic equation).
- 2 Find a basis in each eigenspace of A and merge these bases into one set S .

An Algorithm for Diagonalisation

Given a matrix A , this algorithm diagonalises it (or reports that this is impossible):

- 1 Find the eigenvalues of A (e.g. by solving its characteristic equation).
- 2 Find a basis in each eigenspace of A and merge these bases into one set S .
- 3 If S has fewer than n vectors then A is not diagonalisable.
- 4 Else, form the matrix $P = [\mathbf{p}_1 | \dots | \mathbf{p}_n]$ where $S = \{\mathbf{p}_1, \dots, \mathbf{p}_n\}$.

An Algorithm for Diagonalisation

Given a matrix A , this algorithm diagonalises it (or reports that this is impossible):

- ① Find the eigenvalues of A (e.g. by solving its characteristic equation).
- ② Find a basis in each eigenspace of A and merge these bases into one set S .
- ③ If S has fewer than n vectors then A is not diagonalisable.
- ④ Else, form the matrix $P = [\mathbf{p}_1 | \dots | \mathbf{p}_n]$ where $S = \{\mathbf{p}_1, \dots, \mathbf{p}_n\}$.
 - # The set S contains n vectors and is linearly independent (will prove this),
 - # so S is a basis for \mathbb{R}^n . Hence, P is invertible.

An Algorithm for Diagonalisation

Given a matrix A , this algorithm diagonalises it (or reports that this is impossible):

- ① Find the eigenvalues of A (e.g. by solving its characteristic equation).
- ② Find a basis in each eigenspace of A and merge these bases into one set S .
- ③ If S has fewer than n vectors then A is not diagonalisable.
- ④ Else, form the matrix $P = [\mathbf{p}_1 | \dots | \mathbf{p}_n]$ where $S = \{\mathbf{p}_1, \dots, \mathbf{p}_n\}$.
 - # The set S contains n vectors and is linearly independent (will prove this),
 - # so S is a basis for \mathbb{R}^n . Hence, P is invertible.
- ⑤ The matrix $D = P^{-1}AP$ is diagonal, $D = \text{diag}(\lambda_1, \dots, \lambda_n)$ where, for each i , λ_i is the eigenvalue corresponding to \mathbf{p}_i .

An Algorithm for Diagonalisation

Given a matrix A , this algorithm diagonalises it (or reports that this is impossible):

- ① Find the eigenvalues of A (e.g. by solving its characteristic equation).
- ② Find a basis in each eigenspace of A and merge these bases into one set S .
- ③ If S has fewer than n vectors then A is not diagonalisable.
- ④ Else, form the matrix $P = [\mathbf{p}_1 | \dots | \mathbf{p}_n]$ where $S = \{\mathbf{p}_1, \dots, \mathbf{p}_n\}$.
 - # The set S contains n vectors and is linearly independent (will prove this),
 - # so S is a basis for \mathbb{R}^n . Hence, P is invertible.
- ⑤ The matrix $D = P^{-1}AP$ is diagonal, $D = \text{diag}(\lambda_1, \dots, \lambda_n)$ where, for each i , λ_i is the eigenvalue corresponding to \mathbf{p}_i .

Remark:

- If an $n \times n$ matrix has n distinct eigenvalues then it is diagonalisable. Why? We'll see later on.

Example 14.1

For $k \neq 0$, is the following matrix (corresponding to shear) diagonalisable?

$$A = \begin{pmatrix} 1 & k \\ 0 & 1 \end{pmatrix}$$

Example 14.2

Diagonalise the following matrix, if possible,

$$A = \begin{pmatrix} 0 & 0 & -2 \\ 1 & 2 & 1 \\ 1 & 0 & 3 \end{pmatrix}.$$

Outline

- 1 Recap of Last Week & Plan for Today
- 2 Eigendecomposition - Motivation
- 3 Similarity of Matrices
- 4 Diagonalisation of Matrices
- 5 Eigendecomposition**
- 6 Wrapping Things Up

Eigendecomposition

- Let A be a diagonalisable matrix.

Eigendecomposition

- Let A be a diagonalisable matrix.
- We have $AP = PD$,

Eigendecomposition

- Let A be a diagonalisable matrix.
- We have $AP = PD$,
where P is invertible and its columns $\mathbf{p}_1, \dots, \mathbf{p}_n$ are eigenvectors of A , and $D = \text{diag}(\lambda_1, \dots, \lambda_n)$ and each λ_i is the eigenvalue of A corresponding to \mathbf{p}_i .

Eigendecomposition

- Let A be a diagonalisable matrix.
- We have $AP = PD$,
where P is invertible and its columns $\mathbf{p}_1, \dots, \mathbf{p}_n$ are eigenvectors of A , and $D = \text{diag}(\lambda_1, \dots, \lambda_n)$ and each λ_i is the eigenvalue of A corresponding to \mathbf{p}_i .
- Then we have a decomposition $A = PDP^{-1}$.
Since the factors in this decomposition are made of eigenvectors and eigenvalues, it is called an **eigendecomposition** of A .

Eigendecomposition

- Let A be a diagonalisable matrix.
- We have $AP = PD$,
where P is invertible and its columns $\mathbf{p}_1, \dots, \mathbf{p}_n$ are eigenvectors of A , and $D = \text{diag}(\lambda_1, \dots, \lambda_n)$ and each λ_i is the eigenvalue of A corresponding to \mathbf{p}_i .
- Then we have a decomposition $A = PDP^{-1}$.
Since the factors in this decomposition are made of eigenvectors and eigenvalues, it is called an **eigendecomposition** of A .
(Can use the diagonalisation algorithm to find it.)

Eigendecomposition

- Let A be a diagonalisable matrix.
- We have $AP = PD$,
where P is invertible and its columns $\mathbf{p}_1, \dots, \mathbf{p}_n$ are eigenvectors of A , and $D = \text{diag}(\lambda_1, \dots, \lambda_n)$ and each λ_i is the eigenvalue of A corresponding to \mathbf{p}_i .
- Then we have a decomposition $A = PDP^{-1}$.
Since the factors in this decomposition are made of eigenvectors and eigenvalues, it is called an **eigendecomposition** of A .
(Can use the diagonalisation algorithm to find it.)
- Eigendecomposition of matrix A (of size $n \times n$) exists iff A can be diagonalised.

Eigendecomposition

- Let A be a diagonalisable matrix.
- We have $AP = PD$,
where P is invertible and its columns $\mathbf{p}_1, \dots, \mathbf{p}_n$ are eigenvectors of A , and $D = \text{diag}(\lambda_1, \dots, \lambda_n)$ and each λ_i is the eigenvalue of A corresponding to \mathbf{p}_i .
- Then we have a decomposition $A = PDP^{-1}$.
Since the factors in this decomposition are made of eigenvectors and eigenvalues, it is called an **eigendecomposition** of A .
(Can use the diagonalisation algorithm to find it.)
- Eigendecomposition of matrix A (of size $n \times n$) exists iff A can be diagonalised.
- Eigendecomposition of matrix A iff A has n linearly independent eigenvectors.

The Return of Algebraic Multiplicity of Eigenvalues

Theorem (Thm 14.2)

If vectors $\mathbf{v}_1, \dots, \mathbf{v}_k$ are eigenvectors of a matrix A corresponding to (pairwise) distinct eigenvalues $\lambda_1, \dots, \lambda_k$ then the set $\{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ is linearly independent.

The Return of Algebraic Multiplicity of Eigenvalues

Theorem (Thm 14.2)

If vectors $\mathbf{v}_1, \dots, \mathbf{v}_k$ are eigenvectors of a matrix A corresponding to (pairwise) distinct eigenvalues $\lambda_1, \dots, \lambda_k$ then the set $\{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ is linearly independent.

Proof.

Let $r \leq k$ be the largest number such that $\{\mathbf{v}_1, \dots, \mathbf{v}_r\}$ is linearly independent. Assume for contradiction that $r < k$, so $\{\mathbf{v}_1, \dots, \mathbf{v}_r, \mathbf{v}_{r+1}\}$ is linearly dependent:

$$c_1 \mathbf{v}_1 + \dots + c_r \mathbf{v}_r + c_{r+1} \mathbf{v}_{r+1} = \mathbf{0}$$

where not all c_1, \dots, c_r, c_{r+1} are 0.

Since $\{\mathbf{v}_1, \dots, \mathbf{v}_r\}$ is linearly independent, we conclude that $c_{r+1} \neq 0$.

Since \mathbf{v}_{r+1} is an eigenvector, we conclude that $c_i \neq 0$ for some $i \leq r$.

Continued on next slide ...



Proof Continued

Proof.

We assumed that $\{\mathbf{v}_1, \dots, \mathbf{v}_r\}$ is linearly independent, but $\{\mathbf{v}_1, \dots, \mathbf{v}_r, \mathbf{v}_{r+1}\}$ is not:

$$c_1\mathbf{v}_1 + \dots + c_r\mathbf{v}_r + c_{r+1}\mathbf{v}_{r+1} = \mathbf{0} \quad (1)$$

We derived that $c_{r+1} \neq 0$ and $c_i \neq 0$ for some $i \leq r$.

Left multiply both sides of equation (1) by A and use $A\mathbf{v}_i = \lambda_i\mathbf{v}_i$:

$$c_1\lambda_1\mathbf{v}_1 + \dots + c_r\lambda_r\mathbf{v}_r + c_{r+1}\lambda_{r+1}\mathbf{v}_{r+1} = \mathbf{0}. \quad (2)$$

Now multiply both sides of (1) by λ_{r+1} and subtract that from (2) to obtain

$$c_1(\lambda_1 - \lambda_{r+1})\mathbf{v}_1 + \dots + c_r(\lambda_r - \lambda_{r+1})\mathbf{v}_r = \mathbf{0}. \quad (3)$$

So $c_i(\lambda_i - \lambda_{r+1}) = 0$, and hence $c_i = 0$, for all $i \leq r$, a contradiction. □

The Return of Algebraic Multiplicity of Eigenvalues

- ① From Theorem earlier, we know that $n \times n$ matrix is diagonalisable iff it has n linearly independent eigenvectors.

The Return of Algebraic Multiplicity of Eigenvalues

- ① From Theorem earlier, we know that $n \times n$ matrix is diagonalisable iff it has n linearly independent eigenvectors.
- ② A matrix's eigendecomposition exists iff it is diagonalisable.

The Return of Algebraic Multiplicity of Eigenvalues

- ① From Theorem earlier, we know that $n \times n$ matrix is diagonalisable iff it has n linearly independent eigenvectors.
- ② A matrix's eigendecomposition exists iff it is diagonalisable.
- ③ **Implication:** if we have a matrix of size $n \times n$ that has n distinct eigenvalues, then we know that it has an eigendecomposition.
- ④ **Recall:** n distinct eigenvalues means that algebraic multiplicity of all eigenvalues is 1.

The Return of Algebraic Multiplicity of Eigenvalues

- ① From Theorem earlier, we know that $n \times n$ matrix is diagonalisable iff it has n linearly independent eigenvectors.
- ② A matrix's eigendecomposition exists iff it is diagonalisable.
- ③ **Implication:** if we have a matrix of size $n \times n$ that has n distinct eigenvalues, then we know that it has an eigendecomposition.
- ④ **Recall:** n distinct eigenvalues means that algebraic multiplicity of all eigenvalues is 1.
- ⑤ **Important Note:** This does not mean that a matrix that doesn't have n distinct eigenvalues doesn't have an eigendecomposition.

Example 14.3

Give the Eigendecomposition of A if one exists.

$$A = \begin{pmatrix} -2 & 0 & 3 \\ -8 & 2 & 8 \\ 0 & 0 & 1 \end{pmatrix}.$$

Outline

- 1 Recap of Last Week & Plan for Today
- 2 Eigendecomposition - Motivation
- 3 Similarity of Matrices
- 4 Diagonalisation of Matrices
- 5 Eigendecomposition
- 6 Wrapping Things Up

Example exam question

(d) Consider the matrix $A = \begin{pmatrix} -3 & 1 & 2 \\ 1 & -3 & 2 \\ 2 & 2 & 0 \end{pmatrix}$.

i. Diagonalise A

[6 Marks]

ii. A matrix B has an eigendecomposition of $BP = PD$, where , $B = \begin{pmatrix} -2 & 2 & 1 \\ 2 & -2 & 1 \\ 4 & 4 & -2 \end{pmatrix}$, $P = \begin{pmatrix} 0.5 & -1 & -0.5 \\ 0.5 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix}$, $D = \begin{pmatrix} 2 & 0 & 0 \\ 0 & -4 & 0 \\ 0 & 0 & -4 \end{pmatrix}$.

Are B and A similar, give a justification.

[2 Marks]

What We Learnt Today & Next Week's Plan

Today:

- Similarity of matrices
- Diagonalisable matrices (= similar to a diagonal one)
- A characterisation of diagonalisable matrices
- An algorithm for diagonalisation
- Eigendecomposition of matrices

Next Week:

- Inner product spaces.

Example 14.1

For $k \neq 0$, is the following matrix (corresponding to shear) diagonalisable?

$$A = \begin{pmatrix} 1 & k \\ 0 & 1 \end{pmatrix}$$

Example 14.1

For $k \neq 0$, is the following matrix (corresponding to shear) diagonalisable?

$$A = \begin{pmatrix} 1 & k \\ 0 & 1 \end{pmatrix}$$

Solution. First compute the characteristic polynomial $\det(\lambda I - A)$:

$$\det(\lambda I - A) = \begin{vmatrix} \lambda - 1 & -k \\ 0 & \lambda - 1 \end{vmatrix} = (\lambda - 1)^2.$$

The characteristic equation is $(\lambda - 1)^2 = 0$, so A has only one eigenvalue $\lambda_1 = 1$.

Example 14.1

For $k \neq 0$, is the following matrix (corresponding to shear) diagonalisable?

$$A = \begin{pmatrix} 1 & k \\ 0 & 1 \end{pmatrix}$$

Solution. First compute the characteristic polynomial $\det(\lambda I - A)$:

$$\det(\lambda I - A) = \begin{vmatrix} \lambda - 1 & -k \\ 0 & \lambda - 1 \end{vmatrix} = (\lambda - 1)^2.$$

The characteristic equation is $(\lambda - 1)^2 = 0$, so A has only one eigenvalue $\lambda_1 = 1$.

The corresponding eigenspace is the nullspace of $\lambda_1 I - A = I - A = \begin{pmatrix} 0 & -k \\ 0 & 0 \end{pmatrix}$.

Does it have two linearly independent vectors? (Is $\text{nullity}(I - A) = 2$?)

Example 14.1

For $k \neq 0$, is the following matrix (corresponding to shear) diagonalisable?

$$A = \begin{pmatrix} 1 & k \\ 0 & 1 \end{pmatrix}$$

Solution. First compute the characteristic polynomial $\det(\lambda I - A)$:

$$\det(\lambda I - A) = \begin{vmatrix} \lambda - 1 & -k \\ 0 & \lambda - 1 \end{vmatrix} = (\lambda - 1)^2.$$

The characteristic equation is $(\lambda - 1)^2 = 0$, so A has only one eigenvalue $\lambda_1 = 1$.

The corresponding eigenspace is the nullspace of $\lambda_1 I - A = I - A = \begin{pmatrix} 0 & -k \\ 0 & 0 \end{pmatrix}$.

Does it have two linearly independent vectors? (Is $\text{nullity}(I - A) = 2$?)

Since $\text{rank}(I - A) + \text{nullity}(I - A) = 2$ (by the Dimension Theorem for matrices) and $\text{rank}(I - A) = 1$, we conclude that $\text{nullity}(I - A) = 1$, and therefore A is **not** diagonalisable.

Example 14.2

Diagonalise the following matrix, if possible,

$$A = \begin{pmatrix} 0 & 0 & -2 \\ 1 & 2 & 1 \\ 1 & 0 & 3 \end{pmatrix}.$$

Example 14.2

Diagonalise the following matrix, if possible,

$$A = \begin{pmatrix} 0 & 0 & -2 \\ 1 & 2 & 1 \\ 1 & 0 & 3 \end{pmatrix}.$$

Solution. We have

$$\det(\lambda I - A) = \begin{vmatrix} \lambda & 0 & 2 \\ -1 & \lambda - 2 & -1 \\ -1 & 0 & \lambda - 3 \end{vmatrix} = \lambda^3 - 5\lambda^2 + 8\lambda - 4.$$

The characteristic equation is $\lambda^3 - 5\lambda^2 + 8\lambda - 4 = 0$, can factor it (as in last lecture): $(\lambda - 1)(\lambda - 2)^2 = 0$. The eigenvalues are $\lambda_1 = 1$ and $\lambda_2 = 2$.

Example 14.2

Diagonalise the following matrix, if possible,

$$A = \begin{pmatrix} 0 & 0 & -2 \\ 1 & 2 & 1 \\ 1 & 0 & 3 \end{pmatrix}.$$

Solution. We have

$$\det(\lambda I - A) = \begin{vmatrix} \lambda & 0 & 2 \\ -1 & \lambda - 2 & -1 \\ -1 & 0 & \lambda - 3 \end{vmatrix} = \lambda^3 - 5\lambda^2 + 8\lambda - 4.$$

The characteristic equation is $\lambda^3 - 5\lambda^2 + 8\lambda - 4 = 0$, can factor it (as in last lecture): $(\lambda - 1)(\lambda - 2)^2 = 0$. The eigenvalues are $\lambda_1 = 1$ and $\lambda_2 = 2$.

The corresponding eigenspaces are the nullspaces of $I - A$ and $2I - A$, resp.

Example 14.2

Diagonalise the following matrix, if possible,

$$A = \begin{pmatrix} 0 & 0 & -2 \\ 1 & 2 & 1 \\ 1 & 0 & 3 \end{pmatrix}.$$

Solution. We have

$$\det(\lambda I - A) = \begin{vmatrix} \lambda & 0 & 2 \\ -1 & \lambda - 2 & -1 \\ -1 & 0 & \lambda - 3 \end{vmatrix} = \lambda^3 - 5\lambda^2 + 8\lambda - 4.$$

The characteristic equation is $\lambda^3 - 5\lambda^2 + 8\lambda - 4 = 0$, can factor it (as in last lecture): $(\lambda - 1)(\lambda - 2)^2 = 0$. The eigenvalues are $\lambda_1 = 1$ and $\lambda_2 = 2$.

The corresponding eigenspaces are the nullspaces of $I - A$ and $2I - A$, resp. Using the algorithms for finding a basis in a nullspace, get the following:

$$\lambda_1 = 1 : \mathbf{p}_1 = (-2, 1, 1); \quad \lambda_1 = 2 : \mathbf{p}_2 = (-1, 0, 1) \text{ and } \mathbf{p}_3 = (0, 1, 0).$$

Example 14.2 Cont'd

The eigenvalues of A are $\lambda_1 = 1$ and $\lambda_2 = 2$, and the bases of nullspaces are

$$\lambda_1 = 1 : \mathbf{p}_1 = (-2, 1, 1); \quad \lambda_1 = 2 : \mathbf{p}_2 = (-1, 0, 1) \text{ and } \mathbf{p}_3 = (0, 1, 0).$$

Example 14.2 Cont'd

The eigenvalues of A are $\lambda_1 = 1$ and $\lambda_2 = 2$, and the bases of nullspaces are

$$\lambda_1 = 1 : \mathbf{p}_1 = (-2, 1, 1); \quad \lambda_1 = 2 : \mathbf{p}_2 = (-1, 0, 1) \text{ and } \mathbf{p}_3 = (0, 1, 0).$$

Hence, A is diagonalisable, and one possible matrix that diagonalises it is

$$P = \begin{pmatrix} -2 & -1 & 0 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}.$$

One can check that $P^{-1}AP$ is

$$\begin{pmatrix} -2 & -1 & 0 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}^{-1} \begin{pmatrix} 0 & 0 & -2 \\ 1 & 2 & 1 \\ 1 & 0 & 3 \end{pmatrix} \begin{pmatrix} -2 & -1 & 0 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{pmatrix}$$

Example 14.2 Cont'd

The eigenvalues of A are $\lambda_1 = 1$ and $\lambda_2 = 2$, and the bases of nullspaces are

$$\lambda_1 = 1 : \mathbf{p}_1 = (-2, 1, 1); \quad \lambda_1 = 2 : \mathbf{p}_2 = (-1, 0, 1) \text{ and } \mathbf{p}_3 = (0, 1, 0).$$

Hence, A is diagonalisable, and one possible matrix that diagonalises it is

$$P = \begin{pmatrix} -2 & -1 & 0 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}.$$

One can check that $P^{-1}AP$ is

$$\begin{pmatrix} -2 & -1 & 0 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}^{-1} \begin{pmatrix} 0 & 0 & -2 \\ 1 & 2 & 1 \\ 1 & 0 & 3 \end{pmatrix} \begin{pmatrix} -2 & -1 & 0 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{pmatrix}$$

Remark: The order of the \mathbf{p}_i 's in P can be changed arbitrarily, this will result in changing the order of the λ_i 's in D accordingly.

Example 14.3

Give the Eigendecomposition of A if one exists.

$$A = \begin{pmatrix} -2 & 0 & 3 \\ -8 & 2 & 8 \\ 0 & 0 & 1 \end{pmatrix}.$$

Solution:

- 1) Find eigenvalues: $\lambda_1 = -2, \lambda_2 = 2, \lambda_3 = 1$
- 2) Find eigenvectors: $\mathbf{v}_1 = (1, 2, 0), \mathbf{v}_2 = (0, 1, 0), \mathbf{v}_3 = (1, 0, 1)$
- 3) $D = \begin{pmatrix} -2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad P = \begin{pmatrix} 1 & 0 & 1 \\ 2 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$

Example 14.3 cont

4) Invert P , $P^{-1} = \begin{pmatrix} 1 & 0 & -1 \\ -2 & 1 & 2 \\ 0 & 0 & 1 \end{pmatrix}$

Giving:

$$\begin{pmatrix} -2 & 0 & 3 \\ -8 & 2 & 8 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 1 \\ 2 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} -2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & -1 \\ -2 & 1 & 2 \\ 0 & 0 & 1 \end{pmatrix}$$

The End