

# Lecture 2: Paths, Cycles, Connectivity

Dr. Amitabh Trehan

`amitabh.trehan@durham.ac.uk`

*\*Based on the slides of ADS-21/22 by Dr. George Mertzios*

# Reminder from last lecture

# Reminder from last lecture

- A **graph**  $G$  is a pair  $(V(G), E(G))$ , where
  - $V(G)$  is a **nonempty** set of **vertices** (or **nodes**),
  - $E(G)$  is a set of **unordered pairs**  $uv$  with  $u, v \in V(G)$  and  $u \neq v$ , called the **edges** of  $G$ .

# Reminder from last lecture

- A **graph**  $G$  is a pair  $(V(G), E(G))$ , where
  - $V(G)$  is a **nonempty** set of **vertices** (or **nodes**),
  - $E(G)$  is a set of **unordered pairs**  $uv$  with  $u, v \in V(G)$  and  $u \neq v$ , called the **edges** of  $G$ .
- When  $G$  is clear from the context, we write simply  $V$  and  $E$ .

# Reminder from last lecture

- A **graph**  $G$  is a pair  $(V(G), E(G))$ , where
  - $V(G)$  is a **nonempty** set of **vertices** (or **nodes**),
  - $E(G)$  is a set of **unordered pairs**  $uv$  with  $u, v \in V(G)$  and  $u \neq v$ , called the **edges** of  $G$ .
- When  $G$  is clear from the context, we write simply  $V$  and  $E$ .
- Graphs can be directed, multi-, pseudo-, weighted. By default they are not.

# Reminder from last lecture

- A **graph**  $G$  is a pair  $(V(G), E(G))$ , where
  - $V(G)$  is a **nonempty** set of **vertices** (or **nodes**),
  - $E(G)$  is a set of **unordered pairs**  $uv$  with  $u, v \in V(G)$  and  $u \neq v$ , called the **edges** of  $G$ .
- When  $G$  is clear from the context, we write simply  $V$  and  $E$ .
- Graphs can be directed, multi-, pseudo-, weighted. By default they are not.
- Many problems can be modelled using graphs.

# Reminder from last lecture

- A **graph**  $G$  is a pair  $(V(G), E(G))$ , where
  - $V(G)$  is a **nonempty** set of **vertices** (or **nodes**),
  - $E(G)$  is a set of **unordered pairs**  $uv$  with  $u, v \in V(G)$  and  $u \neq v$ , called the **edges** of  $G$ .
- When  $G$  is clear from the context, we write simply  $V$  and  $E$ .
- Graphs can be directed, multi-, pseudo-, weighted. By default they are not.
- Many problems can be modelled using graphs.
- We can draw graphs:
  - represent each vertex  $u$  by a point, and
  - each edge  $uv$  by a line or curve connecting  $u$  and  $v$ .

# Reminder from last lecture

- A **graph**  $G$  is a pair  $(V(G), E(G))$ , where
  - $V(G)$  is a **nonempty** set of **vertices** (or **nodes**),
  - $E(G)$  is a set of **unordered pairs**  $uv$  with  $u, v \in V(G)$  and  $u \neq v$ , called the **edges** of  $G$ .
- When  $G$  is clear from the context, we write simply  $V$  and  $E$ .
- Graphs can be directed, multi-, pseudo-, weighted. By default they are not.
- Many problems can be modelled using graphs.
- We can draw graphs:
  - represent each vertex  $u$  by a point, and
  - each edge  $uv$  by a line or curve connecting  $u$  and  $v$ .
- Paths, cycles, bipartite graphs, complete graphs, hypercubes



# Contents for today's lecture

- Paths and directed paths;
- The shortest path problem;
- Connectivity and connected components;
- Eulerian and Hamiltonian cycles;
- Examples and exercises.

# Walks, paths, cycles, and distances

# Walks, paths, cycles, and distances

- A **walk** in a graph  $G$  is a sequence of edges  $v_0v_1, v_1v_2, v_2v_3, \dots, v_{n-1}v_n$ . In this case we also say that  $v_0, v_1, \dots, v_n$  is a walk in  $G$ .
- A walk  $v_0, v_1, \dots, v_n$  in  $G$  is a **path** if all  $v_i$ 's are distinct. In this case we also say that  $v_0, v_1, \dots, v_n$  is a path in  $G$ .

# Walks, paths, cycles, and distances

- A **walk** in a graph  $G$  is a sequence of edges  $v_0v_1, v_1v_2, v_2v_3, \dots, v_{n-1}v_n$ . In this case we also say that  $v_0, v_1, \dots, v_n$  is a walk in  $G$ .
- A walk  $v_0, v_1, \dots, v_n$  in  $G$  is a **path** if all  $v_i$ 's are distinct. In this case we also say that  $v_0, v_1, \dots, v_n$  is a path in  $G$ .
- A walk  $v_0, v_1, \dots, v_n$  with  $v_0 = v_n$  is called a **circuit** or **closed walk**.

# Walks, paths, cycles, and distances

- A **walk** in a graph  $G$  is a sequence of edges  $v_0v_1, v_1v_2, v_2v_3, \dots, v_{n-1}v_n$ . In this case we also say that  $v_0, v_1, \dots, v_n$  is a walk in  $G$ .
- A walk  $v_0, v_1, \dots, v_n$  in  $G$  is a **path** if all  $v_i$ 's are distinct. In this case we also say that  $v_0, v_1, \dots, v_n$  is a path in  $G$ .
- A walk  $v_0, v_1, \dots, v_n$  with  $v_0 = v_n$  is called a **circuit** or **closed walk**.
- A closed walk is a **cycle** (or **simple circuit**) if all  $v_i$ 's in it are distinct except  $v_0 = v_n$ .

# Walks, paths, cycles, and distances

- A **walk** in a graph  $G$  is a sequence of edges  $v_0v_1, v_1v_2, v_2v_3, \dots, v_{n-1}v_n$ . In this case we also say that  $v_0, v_1, \dots, v_n$  is a walk in  $G$ .
- A walk  $v_0, v_1, \dots, v_n$  in  $G$  is a **path** if all  $v_i$ 's are distinct. In this case we also say that  $v_0, v_1, \dots, v_n$  is a path in  $G$ .
- A walk  $v_0, v_1, \dots, v_n$  with  $v_0 = v_n$  is called a **circuit** or **closed walk**.
- A closed walk is a **cycle** (or **simple circuit**) if all  $v_i$ 's in it are distinct except  $v_0 = v_n$ .
- If  $G$  is a directed graph then the **directed paths** and **directed cycles** are defined in a natural way, with each edge being directed from  $v_i$  to  $v_{i+1}$ .

# Walks, paths, cycles, and distances

- A **walk** in a graph  $G$  is a sequence of edges  $v_0v_1, v_1v_2, v_2v_3, \dots, v_{n-1}v_n$ . In this case we also say that  $v_0, v_1, \dots, v_n$  is a walk in  $G$ .
- A walk  $v_0, v_1, \dots, v_n$  in  $G$  is a **path** if all  $v_i$ 's are distinct. In this case we also say that  $v_0, v_1, \dots, v_n$  is a path in  $G$ .
- A walk  $v_0, v_1, \dots, v_n$  with  $v_0 = v_n$  is called a **circuit** or **closed walk**.
- A closed walk is a **cycle** (or **simple circuit**) if all  $v_i$ 's in it are distinct except  $v_0 = v_n$ .
- If  $G$  is a directed graph then the **directed paths** and **directed cycles** are defined in a natural way, with each edge being directed from  $v_i$  to  $v_{i+1}$ .
- The **length** of a path or a cycle is the number of edges in it.

# Walks, paths, cycles, and distances


- A **walk** in a graph  $G$  is a sequence of edges  $v_0 v_1, v_1 v_2, v_2 v_3, \dots, v_{n-1} v_n$ . In this case we also say that  $v_0, v_1, \dots, v_n$  is a walk in  $G$ .
- A walk  $v_0, v_1, \dots, v_n$  in  $G$  is a **path** if all  $v_i$ 's are distinct. In this case we also say that  $v_0, v_1, \dots, v_n$  is a path in  $G$ .
- A walk  $v_0, v_1, \dots, v_n$  with  $v_0 = v_n$  is called a **circuit** or **closed walk**.
- A closed walk is a **cycle** (or **simple circuit**) if all  $v_i$ 's in it are distinct except  $v_0 = v_n$ .
- If  $G$  is a directed graph then the **directed paths** and **directed cycles** are defined in a natural way, with each edge being directed from  $v_i$  to  $v_{i+1}$ .
- The **length** of a path or a cycle is the number of edges in it.
- The **distance** between vertices  $u$  and  $v$  in a graph, denoted  $dist(u, v)$ , is the length of a shortest path from  $u$  to  $v$  if such a path exists, and  $\infty$  otherwise.



# Walks, paths, cycles, and distances

- A **walk** in a graph  $G$  is a sequence of edges  $v_0v_1, v_1v_2, v_2v_3, \dots, v_{n-1}v_n$ . In this case we also say that  $v_0, v_1, \dots, v_n$  is a walk in  $G$ .
- A walk  $v_0, v_1, \dots, v_n$  in  $G$  is a **path** if all  $v_i$ 's are distinct. In this case we also say that  $v_0, v_1, \dots, v_n$  is a path in  $G$ .
- A walk  $v_0, v_1, \dots, v_n$  with  $v_0 = v_n$  is called a **circuit** or **closed walk**.
- A closed walk is a **cycle** (or **simple circuit**) if all  $v_i$ 's in it are distinct except  $v_0 = v_n$ .
- If  $G$  is a directed graph then the **directed paths** and **directed cycles** are defined in a natural way, with each edge being directed from  $v_i$  to  $v_{i+1}$ .
- The **length** of a path or a cycle is the number of edges in it.
- The **distance** between vertices  $u$  and  $v$  in a graph, denoted  $dist(u, v)$ , is the length of a shortest path from  $u$  to  $v$  if such a path exists, and  $\infty$  otherwise.
- The **diameter** of a graph is the largest distance between two vertices in it


# Exercise



- Does this graph contain a path of length 7?
- Does it contain a cycle of length 7?
- What is the distance from 2 to 6?
- What is the diameter of this graph?
- Answer at [PolIeV.com/amitabhTREHAN005](https://pollev.com/amitabhTREHAN005)

## Exercise: Solutions


- Does this graph contain a path of length 7?



- Does it contain a cycle of length 7?
- What is the distance from 2 to 6?
- What is the diameter of this graph?

## Exercise: Solutions


- Does this graph contain a path of length 7?
- Does it contain a cycle of length 7?



- What is the distance from 2 to 6?
- What is the diameter of this graph?


## Exercise: Solutions

- Does this graph contain a path of length 7?
- Does it contain a cycle of length 7?
- What is the distance from 2 to 6?



- What is the diameter of this graph?


## Exercise: Solutions



- Does this graph contain a path of length 7?
- Does it contain a cycle of length 7?
- What is the distance from 2 to 6?
- What is the diameter of this graph?

The diameter is 3. Any way to prove this?

## Exercise: Solutions



- Does this graph contain a path of length 7?
- Does it contain a cycle of length 7?
- What is the distance from 2 to 6?
- What is the diameter of this graph?

The diameter is 3. Any way to prove this?

Proof: there is a DOMINATING SET  $\{1,5,7\}$  which is a TRIANGLE

# The acquaintance graph and six degrees of separation

The **acquaintance graph**:

- The vertices are all people
- There is an edge between two of them if they are acquainted





# The acquaintance graph and six degrees of separation

The **acquaintance graph**:

- The vertices are all people
- There is an edge between two of them if they are acquainted



- Many sociologists conjectured/observed that this graph has diameter 6, after removing a tiny fraction of vertices.

# The acquaintance graph and six degrees of separation

The **acquaintance graph**:

- The vertices are all people
- There is an edge between two of them if they are acquainted



- Many sociologists conjectured/observed that this graph has diameter 6, after removing a tiny fraction of vertices. (Can you put this in plain words?)

# The acquaintance graph and six degrees of separation

The **acquaintance graph**:

- The vertices are all people
- There is an edge between two of them if they are acquainted



- Many sociologists conjectured/observed that this graph has diameter 6, after removing a tiny fraction of vertices. (Can you put this in plain words?)
- This is known as the “small world phenomenon”.

# The acquaintance graph and six degrees of separation

The acquaintance graph:

- The vertices are all people
- There is an edge between two of them if they are acquainted



- Many sociologists conjectured/observed that this graph has diameter 6, after removing a tiny fraction of vertices. (Can you put this in plain words?)
- This is known as the “small world phenomenon”.
- There is a popular play (and a film) based on this, called “Six degrees of separation”.

# The collaboration graph and the Erdős number

The **collaboration graph**:

- The vertices are all people (or only scientists)
- There is an edge between two of them if they have written a joint paper

# The collaboration graph and the Erdős number

The **collaboration graph**:

- The vertices are all people (or only scientists)
- There is an edge between two of them if they have written a joint paper



- Paul Erdős (1913-1996) was a great and extremely prolific mathematician.

# The collaboration graph and the Erdős number

The **collaboration graph**:

- The vertices are all people (or only scientists)
- There is an edge between two of them if they have written a joint paper



- Paul Erdős (1913-1996) was a great and extremely prolific mathematician.
- The Erdős number of a person is the distance to Paul Erdős in the collaboration graph.

# The collaboration graph and the Erdős number

The **collaboration graph**:

- The vertices are all people (or only scientists)
- There is an edge between two of them if they have written a joint paper



- Paul Erdős (1913-1996) was a great and extremely prolific mathematician.
- The Erdős number of a person is the distance to Paul Erdős in the collaboration graph.
- Btw, my Erdős number is 3. (Can you put this in plain words?)



# The Hollywood graph and the Bacon number

The **Hollywood graph**:

- The vertices are all people (or only actors).
- There is an edge between two of them if they appeared in the same film.

# The Hollywood graph and the Bacon number

The **Hollywood graph**:

- The vertices are all people (or only actors).
- There is an edge between two of them if they appeared in the same film.



- Kevin Bacon is a famous Hollywood actor.

# The Hollywood graph and the Bacon number

The **Hollywood graph**:

- The vertices are all people (or only actors).
- There is an edge between two of them if they appeared in the same film.



- Kevin Bacon is a famous Hollywood actor.
- The Bacon number of a person is the distance to Kevin Bacon in the Hollywood graph.

# The Hollywood graph and the Bacon number

The **Hollywood graph**:

- The vertices are all people (or only actors).
- There is an edge between two of them if they appeared in the same film.



- Kevin Bacon is a famous Hollywood actor.
- The Bacon number of a person is the distance to Kevin Bacon in the Hollywood graph.
- About 90% of actors have a Bacon number (i.e. the distance is not  $\infty$ )

# The Erdős-Bacon number

## Definition

For any person, their Erdős-Bacon number is the sum of their Erdős number and their Bacon number.

# The Erdős-Bacon number

## Definition

For any person, their Erdős-Bacon number is the sum of their Erdős number and their Bacon number.

There are not many people with a small Erdős-Bacon number.

# The Erdős-Bacon number

## Definition

For any person, their Erdős-Bacon number is the sum of their Erdős number and their Bacon number.

There are not many people with a small Erdős-Bacon number.

For example, the following people have Erdős-Bacon number 7 or less.

# The Erdős-Bacon number

## Definition

For any person, their Erdős-Bacon number is the sum of their Erdős number and their Bacon number.

There are not many people with a small Erdős-Bacon number.

For example, the following people have Erdős-Bacon number 7 or less.





# The Erdős-Bacon number

## Definition

For any person, their Erdős-Bacon number is the sum of their Erdős number and their Bacon number.

There are not many people with a small Erdős-Bacon number.

For example, the following people have Erdős-Bacon number 7 or less.



# The Erdős-Bacon number

## Definition

For any person, their Erdős-Bacon number is the sum of their Erdős number and their Bacon number.

There are not many people with a small Erdős-Bacon number.

For example, the following people have Erdős-Bacon number 7 or less.



- Btw, my Erdős-Bacon number is  $\infty$ , but I have a colleague of a colleague who has a co-author (Hubie Chen) with Erdős-Bacon number 5 ( $3+2$ )

# Shortest-path problems

In a graph (possibly with **edge weights**), the problem of computing a path from a given vertex  $u$  (“source”) to a given vertex  $v$  (“target”) with the smallest total length (or weight) is known as the **shortest-path problem**.

# Shortest-path problems


In a graph (possibly with **edge weights**), the problem of computing a path from a given vertex  $u$  (“source”) to a given vertex  $v$  (“target”) with the smallest total length (or weight) is known as the **shortest-path problem**.

We all often (use software applications that) compute such paths. Example?

# Shortest-path problems

In a graph (possibly with **edge weights**, the problem of computing a path from a given vertex  $u$  (“source”) to a given vertex  $v$  (“target”) with the smallest total length (or weight) is known as the **shortest-path problem**.


We all often (use software applications that) compute such paths. Example?



# Shortest-path problems

In a graph (possibly with **edge weights**, the problem of computing a path from a given vertex  $u$  (“source”) to a given vertex  $v$  (“target”) with the smallest total length (or weight) is known as the **shortest-path problem**.

We all often (use software applications that) compute such paths. Example?




We will learn about algorithms for the (unweighted) problem in a few lectures.

# Connectivity

## Definition

A graph  $G = (V, E)$  is called **connected** if, between every pair of vertices  $u, v$ , there exists at least one path in  $G$ .

A **connected component** of  $G$  is a **maximal** connected subgraph of  $G$ .




- Is this graph connected?

# Connectivity

## Definition

A graph  $G = (V, E)$  is called **connected** if, between every pair of vertices  $u, v$ , there exists at least one path in  $G$ .

A **connected component** of  $G$  is a **maximal** connected subgraph of  $G$ .



- What about this graph?




# Connectivity

## Definition

A graph  $G = (V, E)$  is called **connected** if, between every pair of vertices  $u, v$ , there exists at least one path in  $G$ .

A **connected component** of  $G$  is a **maximal** connected subgraph of  $G$ .




- What about this graph?

# Connectivity

## Definition

A graph  $G = (V, E)$  is called **connected** if, between every pair of vertices  $u, v$ , there exists at least one path in  $G$ .

A **connected component** of  $G$  is a **maximal** connected subgraph of  $G$ .



- How many connected components does this graph have?

# Connected components

## Theorem

*Every graph  $G = (V, E)$  contains at least  $|V| - |E|$  connected components.*

# Connected components

## Theorem

*Every graph  $G = (V, E)$  contains at least  $|V| - |E|$  connected components.*

## Proof.

By induction over  $m = |E|$ .



# Connected components

## Theorem

*Every graph  $G = (V, E)$  contains at least  $|V| - |E|$  connected components.*

## Proof.

By induction over  $m = |E|$ .

**Induction basis:**  $m = 0$ . No edges, so only isolated vertices, i.e.  $|V|$  connected components.



# Connected components

## Theorem

*Every graph  $G = (V, E)$  contains at least  $|V| - |E|$  connected components.*

## Proof.

By induction over  $m = |E|$ .

**Induction basis:**  $m = 0$ . No edges, so only isolated vertices, i.e.  $|V|$  connected components.

**Induction step:** Let  $G = (V, E)$  with  $|E| = m + 1$ . Consider an arbitrary  $e \in E$ , and define  $E' = E \setminus \{e\}$ . By induction hypothesis:  $G' = (V, E')$  has at least  $|V| - |E'| = |V| - m$  connected components. Two cases:

- ① With  $e$ , the number of connected components does not change.
- ② With  $e$ , the number of connected components ???



# Connected components

## Theorem

*Every graph  $G = (V, E)$  contains at least  $|V| - |E|$  connected components.*

## Proof.

By induction over  $m = |E|$ .

**Induction basis:**  $m = 0$ . No edges, so only isolated vertices, i.e.  $|V|$  connected components.

**Induction step:** Let  $G = (V, E)$  with  $|E| = m + 1$ . Consider an arbitrary  $e \in E$ , and define  $E' = E \setminus \{e\}$ . By induction hypothesis:  $G' = (V, E')$  has at least  $|V| - |E'| = |V| - m$  connected components. Two cases:

- 1 With  $e$ , the number of connected components does not change.
- 2 With  $e$ , the number of connected components decreases by 1.



# Connected components

## Theorem

Every graph  $G = (V, E)$  contains at least  $|V| - |E|$  connected components.

## Proof.

By induction over  $m = |E|$ .

**Induction basis:**  $m = 0$ . No edges, so only isolated vertices, i.e.  $|V|$  connected components.

**Induction step:** Let  $G = (V, E)$  with  $|E| = m + 1$ . Consider an arbitrary  $e \in E$ , and define  $E' = E \setminus \{e\}$ . By induction hypothesis:  $G' = (V, E')$  has at least  $|V| - |E'| = |V| - m$  connected components. Two cases:

- 1 With  $e$ , the number of connected components does not change.
- 2 With  $e$ , the number of connected components decreases by 1.

In both cases,  $G$  has at least  $|V| - m - 1 = |V| - |E|$  connected components.  $\square$



# Connected components

## Theorem

Every graph  $G = (V, E)$  contains at least  $|V| - |E|$  connected components.

## Proof.

By induction over  $m = |E|$ .

**Induction basis:**  $m = 0$ . No edges, so only isolated vertices, i.e.  $|V|$  connected components.

**Induction step:** Let  $G = (V, E)$  with  $|E| = m + 1$ . Consider an arbitrary  $e \in E$ , and define  $E' = E \setminus \{e\}$ . By induction hypothesis:  $G' = (V, E')$  has at least  $|V| - |E'| = |V| - m$  connected components. Two cases:

- 1 With  $e$ , the number of connected components does not change.
- 2 With  $e$ , the number of connected components decreases by 1.

In both cases,  $G$  has at least  $|V| - m - 1 = |V| - |E|$  connected components.  $\square$

## Corollary (useful in various algorithmic proofs)

If  $G = (V, E)$  is connected then  $|E| \geq |V| - 1$ .

# Exercise

**Exercise 1:** Prove that if  $G$  is a graph on  $n$  vertices and  $\delta(G) \geq (n-1)/2$  then  $G$  is connected.

# Exercise

**Exercise 1:** Prove that if  $G$  is a graph on  $n$  vertices and  $\delta(G) \geq (n-1)/2$  then  $G$  is connected.

- Proof **by contradiction**: Assume  $G$  is not connected, derive a contradiction.

# Exercise

**Exercise 1:** Prove that if  $G$  is a graph on  $n$  vertices and  $\delta(G) \geq (n-1)/2$  then  $G$  is connected.

- Proof **by contradiction**: Assume  $G$  is not connected, derive a contradiction.
- Take two vertices  $u$  and  $v$  in different connected components.

# Exercise

**Exercise 1:** Prove that if  $G$  is a graph on  $n$  vertices and  $\delta(G) \geq (n-1)/2$  then  $G$  is connected.

- Proof **by contradiction**: Assume  $G$  is not connected, derive a contradiction.
- Take two vertices  $u$  and  $v$  in different connected components.
- We have that  $\deg(u) \geq (n-1)/2$  and  $\deg(v) \geq (n-1)/2$ .

# Exercise

**Exercise 1:** Prove that if  $G$  is a graph on  $n$  vertices and  $\delta(G) \geq (n-1)/2$  then  $G$  is connected.

- Proof **by contradiction**: Assume  $G$  is not connected, derive a contradiction.
- Take two vertices  $u$  and  $v$  in different connected components.
- We have that  $\deg(u) \geq (n-1)/2$  and  $\deg(v) \geq (n-1)/2$ .
- The set  $N(u)$  contains none of the neighbours of  $v$  nor  $v$  itself.

# Exercise

**Exercise 1:** Prove that if  $G$  is a graph on  $n$  vertices and  $\delta(G) \geq (n-1)/2$  then  $G$  is connected.

- Proof **by contradiction**: Assume  $G$  is not connected, derive a contradiction.
- Take two vertices  $u$  and  $v$  in different connected components.
- We have that  $\deg(u) \geq (n-1)/2$  and  $\deg(v) \geq (n-1)/2$ .
- The set  $N(u)$  contains none of the neighbours of  $v$  nor  $v$  itself.
- Similarly,  $N(v)$  contains none of the neighbours of  $u$  nor  $u$  itself.

# Exercise

**Exercise 1:** Prove that if  $G$  is a graph on  $n$  vertices and  $\delta(G) \geq (n-1)/2$  then  $G$  is connected.

- Proof **by contradiction**: Assume  $G$  is not connected, derive a contradiction.
- Take two vertices  $u$  and  $v$  in different connected components.
- We have that  $\deg(u) \geq (n-1)/2$  and  $\deg(v) \geq (n-1)/2$ .
- The set  $N(u)$  contains none of the neighbours of  $v$  nor  $v$  itself.
- Similarly,  $N(v)$  contains none of the neighbours of  $u$  nor  $u$  itself.
- Hence  $G$  contains at least  $\frac{n-1}{2} + 1 + \frac{n-1}{2} + 1 = n + 1$  vertices, a contradiction.



# Exercise

**Exercise 1:** Prove that if  $G$  is a graph on  $n$  vertices and  $\delta(G) \geq (n-1)/2$  then  $G$  is connected.

- Proof **by contradiction**: Assume  $G$  is not connected, derive a contradiction.
- Take two vertices  $u$  and  $v$  in different connected components.
- We have that  $\deg(u) \geq (n-1)/2$  and  $\deg(v) \geq (n-1)/2$ .
- The set  $N(u)$  contains none of the neighbours of  $v$  nor  $v$  itself.
- Similarly,  $N(v)$  contains none of the neighbours of  $u$  nor  $u$  itself.
- Hence  $G$  contains at least  $\frac{n-1}{2} + 1 + \frac{n-1}{2} + 1 = n + 1$  vertices, a contradiction.
- In fact, we even proved more: any two vertices in  $G$  are at distance at most 2 (so the diameter of  $G$  is at most 2).


# Strong connectivity

## Definition

A directed graph  $G$  is called (weakly) connected if the graph obtained from  $G$  by forgetting directions is connected.

A directed graph is called strongly connected if any two distinct vertices are connected by directed paths in both directions.

A strongly connected component (or simply strong component) of a digraph  $G$  is a maximal strongly connected subgraph of  $G$ .



- Is this graph strongly connected?


# Strong connectivity

## Definition

A directed graph  $G$  is called (weakly) connected if the graph obtained from  $G$  by forgetting directions is connected.

A directed graph is called strongly connected if any two distinct vertices are connected by directed paths in both directions.

A strongly connected component (or simply strong component) of a digraph  $G$  is a maximal strongly connected subgraph of  $G$ .



- Is this graph strongly connected?

# Connectivity properties of commercially important graphs

Important large graphs tend to have similar connectivity properties:

See the Stanford Large Networks Dataset Collection:

<https://snap.stanford.edu/data/index.html>

# Connectivity properties of commercially important graphs

Important large graphs tend to have similar connectivity properties:

See the Stanford Large Networks Dataset Collection:

<https://snap.stanford.edu/data/index.html>

Key: WCC = weakly connected component, SCC = strongly ...

# Connectivity properties of commercially important graphs

Important large graphs tend to have similar connectivity properties:

See the Stanford Large Networks Dataset Collection:

<https://snap.stanford.edu/data/index.html>

Key: WCC = weakly connected component, SCC = strongly ...

- Web graph: 50% SCC, 91% WCC

# Connectivity properties of commercially important graphs

Important large graphs tend to have similar connectivity properties:

See the Stanford Large Networks Dataset Collection:

<https://snap.stanford.edu/data/index.html>

Key: WCC = weakly connected component, SCC = strongly ...

- Web graph: 50% SCC, 91% WCC
- Amazon co-purchase graph: 91% SCC, 100% WCC

# Connectivity properties of commercially important graphs

Important large graphs tend to have similar connectivity properties:

See the Stanford Large Networks Dataset Collection:

<https://snap.stanford.edu/data/index.html>

Key: WCC = weakly connected component, SCC = strongly ...

- Web graph: 50% SCC, 91% WCC
- Amazon co-purchase graph: 91% SCC, 100% WCC
- Wikipedia talk graph: 48% SCC, 100% WCC



# Connectivity properties of commercially important graphs

Important large graphs tend to have similar connectivity properties:

See the Stanford Large Networks Dataset Collection:

<https://snap.stanford.edu/data/index.html>

Key: WCC = weakly connected component, SCC = strongly ...

- Web graph: 50% SCC, 91% WCC
- Amazon co-purchase graph: 91% SCC, 100% WCC
- Wikipedia talk graph: 48% SCC, 100% WCC
- Facebook graph: 100% WCC

# Connectivity properties of commercially important graphs

Important large graphs tend to have similar connectivity properties:

See the Stanford Large Networks Dataset Collection:

<https://snap.stanford.edu/data/index.html>

Key: WCC = weakly connected component, SCC = strongly ...

- Web graph: 50% SCC, 91% WCC
- Amazon co-purchase graph: 91% SCC, 100% WCC
- Wikipedia talk graph: 48% SCC, 100% WCC
- Facebook graph: 100% WCC
- Google+ graph: 61% SCC, 100% WCC
- ...

# Special circuits/cycles in graphs


- Can we travel along the edges of a given graph  $G$  so that we start and finish at the same vertex and traverse **each edge exactly once**?

# Special circuits/cycles in graphs

- Can we travel along the edges of a given graph  $G$  so that we start and finish at the same vertex and traverse **each edge exactly once**?
  - Such a circuit in  $G$  is called a **Eulerian** circuit, after Leonhard Euler (1707-83).


# Special circuits/cycles in graphs

- Can we travel along the edges of a given graph  $G$  so that we start and finish at the same vertex and traverse **each edge exactly once**?
  - Such a circuit in  $G$  is called a **Eulerian** circuit, after Leonhard Euler (1707-83).



# Special circuits/cycles in graphs


- Can we travel along the edges of a given graph  $G$  so that we start and finish at the same vertex and traverse **each edge exactly once**?
  - Such a circuit in  $G$  is called a **Eulerian** circuit, after Leonhard Euler (1707-83).



- Can we travel along the edges of a given graph so that we start and finish at the same vertex and visit **each vertex exactly once**?

# Special circuits/cycles in graphs


- Can we travel along the edges of a given graph  $G$  so that we start and finish at the same vertex and traverse **each edge exactly once**?
  - Such a circuit in  $G$  is called a **Eulerian** circuit, after Leonhard Euler (1707-83).




- Can we travel along the edges of a given graph so that we start and finish at the same vertex and visit **each vertex exactly once**?
  - Such a cycle is called a **Hamiltonian** cycle, after William Hamilton (1805-65).

# Special circuits/cycles in graphs

- Can we travel along the edges of a given graph  $G$  so that we start and finish at the same vertex and traverse **each edge exactly once**?
  - Such a circuit in  $G$  is called a **Eulerian** circuit, after Leonhard Euler (1707-83).




- Can we travel along the edges of a given graph so that we start and finish at the same vertex and visit **each vertex exactly once**?
  - Such a cycle is called a **Hamiltonian** cycle, after William Hamilton (1805-65).






# Special circuits/cycles in graphs

- Can we travel along the edges of a given graph  $G$  so that we start and finish at the same vertex and traverse **each edge exactly once**?
  - Such a circuit in  $G$  is called a **Eulerian** circuit, after Leonhard Euler (1707-83).



- Can we travel along the edges of a given graph so that we start and finish at the same vertex and visit **each vertex exactly once**?
  - Such a cycle is called a **Hamiltonian** cycle, after William Hamilton (1805-65).



- Detecting one of these two types of circuits is easy, while detecting the other is not easy at all. Which is which?

# Eulerian circuits

# Eulerian circuits

## Theorem (Rosen p.634-636)

*A connected graph with at least two vertices has an Eulerian circuit iff each of its vertices has even degree.*

# Eulerian circuits

## Theorem (Rosen p.634-636)

*A connected graph with at least two vertices has an Eulerian circuit iff each of its vertices has even degree.*

Idea of proof:

# Eulerian circuits

## Theorem (Rosen p.634-636)

*A connected graph with at least two vertices has an Eulerian circuit iff each of its vertices has even degree.*

Idea of proof:

Necessity ( $\Rightarrow$ ): each time this circuit passes through a vertex  $v$ , it contributes 2 to  $\deg(v)$ . Since each edge is used exactly once,  $\deg(v)$  must be even.

# Eulerian circuits

## Theorem (Rosen p.634-636)

*A connected graph with at least two vertices has an Eulerian circuit iff each of its vertices has even degree.*

Idea of proof:

Necessity ( $\Rightarrow$ ): each time this circuit passes through a vertex  $v$ , it contributes 2 to  $\deg(v)$ . Since each edge is used exactly once,  $\deg(v)$  must be even.

Sufficiency ( $\Leftarrow$ ): Induction on the number of vertices in  $G$ .

# Eulerian circuits

## Theorem (Rosen p.634-636)

*A connected graph with at least two vertices has an Eulerian circuit iff each of its vertices has even degree.*

Idea of proof:

Necessity ( $\Rightarrow$ ): each time this circuit passes through a vertex  $v$ , it contributes 2 to  $\deg(v)$ . Since each edge is used exactly once,  $\deg(v)$  must be even.

Sufficiency ( $\Leftarrow$ ): Induction on the number of vertices in  $G$ .

Induction base:  $G = K_3$ , the claim is obvious. Induction step:

# Eulerian circuits

## Theorem (Rosen p.634-636)

*A connected graph with at least two vertices has an Eulerian circuit iff each of its vertices has even degree.*

Idea of proof:

Necessity ( $\Rightarrow$ ): each time this circuit passes through a vertex  $v$ , it contributes 2 to  $\deg(v)$ . Since each edge is used exactly once,  $\deg(v)$  must be even.

Sufficiency ( $\Leftarrow$ ): Induction on the number of vertices in  $G$ .

Induction base:  $G = K_3$ , the claim is obvious. Induction step:

- Start walking from any vertex  $u$  along the untraversed edges, marking off the traversed edges.



# Eulerian circuits

## Theorem (Rosen p.634-636)

*A connected graph with at least two vertices has an Eulerian circuit iff each of its vertices has even degree.*

Idea of proof:

Necessity ( $\Rightarrow$ ): each time this circuit passes through a vertex  $v$ , it contributes 2 to  $\deg(v)$ . Since each edge is used exactly once,  $\deg(v)$  must be even.

Sufficiency ( $\Leftarrow$ ): Induction on the number of vertices in  $G$ .

Induction base:  $G = K_3$ , the claim is obvious. Induction step:

- Start walking from any vertex  $u$  along the untraversed edges, marking off the traversed edges.
- Stop when you arrive at a vertex where you can't continue (all edges leading from it are already traversed). This vertex must be  $u$  again.

# Eulerian circuits

## Theorem (Rosen p.634-636)

*A connected graph with at least two vertices has an Eulerian circuit iff each of its vertices has even degree.*

Idea of proof:

Necessity ( $\Rightarrow$ ): each time this circuit passes through a vertex  $v$ , it contributes 2 to  $\deg(v)$ . Since each edge is used exactly once,  $\deg(v)$  must be even.

Sufficiency ( $\Leftarrow$ ): Induction on the number of vertices in  $G$ .

Induction base:  $G = K_3$ , the claim is obvious. Induction step:

- Start walking from any vertex  $u$  along the untraversed edges, marking off the traversed edges.
- Stop when you arrive at a vertex where you can't continue (all edges leading from it are already traversed). This vertex must be  $u$  again.
- Hence we have a circuit  $C$ . Delete it from  $G$  to obtain a smaller graph  $H$  in which all degrees are also even.

# Eulerian circuits

## Theorem (Rosen p.634-636)

*A connected graph with at least two vertices has an Eulerian circuit iff each of its vertices has even degree.*

Idea of proof:

Necessity ( $\Rightarrow$ ): each time this circuit passes through a vertex  $v$ , it contributes 2 to  $\deg(v)$ . Since each edge is used exactly once,  $\deg(v)$  must be even.

Sufficiency ( $\Leftarrow$ ): Induction on the number of vertices in  $G$ .

Induction base:  $G = K_3$ , the claim is obvious. Induction step:

- Start walking from any vertex  $u$  along the untraversed edges, marking off the traversed edges.
- Stop when you arrive at a vertex where you can't continue (all edges leading from it are already traversed). This vertex must be  $u$  again.
- Hence we have a circuit  $C$ . Delete it from  $G$  to obtain a smaller graph  $H$  in which all degrees are also even.
- By induction hypothesis, each connected component of  $H$  has an Eulerian circuit.

# Eulerian circuits

## Theorem (Rosen p.634-636)

*A connected graph with at least two vertices has an Eulerian circuit iff each of its vertices has even degree.*

Idea of proof:


Necessity ( $\Rightarrow$ ): each time this circuit passes through a vertex  $v$ , it contributes 2 to  $\deg(v)$ . Since each edge is used exactly once,  $\deg(v)$  must be even.

Sufficiency ( $\Leftarrow$ ): Induction on the number of vertices in  $G$ .

Induction base:  $G = K_3$ , the claim is obvious. Induction step:


- Start walking from any vertex  $u$  along the untraversed edges, marking off the traversed edges.
- Stop when you arrive at a vertex where you can't continue (all edges leading from it are already traversed). This vertex must be  $u$  again.
- Hence we have a circuit  $C$ . Delete it from  $G$  to obtain a smaller graph  $H$  in which all degrees are also even.
- By induction hypothesis, each connected component of  $H$  has an Eulerian circuit.
- Combine  $C$  and these circuits to obtain the required circuit for  $G$ .

# Hamiltonian cycles



- Does this graph have a Eulerian circuit?
- Does it have a Hamiltonian cycle?
- Answer at [PolLEV.com/amitabhtrehan005](https://pollev.com/amitabhtrehan005)

# Hamiltonian cycles



- Does this graph have a Eulerian circuit?
- Does it have a Hamiltonian cycle?
- Answer at [PolIIEV.com/amitabhrehan005](http://PolIIEV.com/amitabhrehan005)

- Detecting Eulerian circuits algorithmically is easy. (How?)
- Detecting Hamiltonian cycles is hard (NP-complete).

# Travelling Salesman Problem (TSP)



# Travelling Salesman Problem (TSP)

The (famous) TSP is the following problem:

# Travelling Salesman Problem (TSP)

The (famous) TSP is the following problem:

- A salesman should visit cities  $c_1, c_2, \dots, c_n$  in some order, visiting each city exactly once and returning to the starting point

# Travelling Salesman Problem (TSP)

The (famous) TSP is the following problem:

- A salesman should visit cities  $c_1, c_2, \dots, c_n$  in some order, visiting each city exactly once and returning to the starting point
- A (positive integer) cost  $d(i, j)$  of travel between each pair  $(c_i, c_j)$  is known.

# Travelling Salesman Problem (TSP)

The (famous) TSP is the following problem:

- A salesman should visit cities  $c_1, c_2, \dots, c_n$  in some order, visiting each city exactly once and returning to the starting point
- A (positive integer) cost  $d(i, j)$  of travel between each pair  $(c_i, c_j)$  is known.
- Goal: find an optimal (i.e. cheapest) route for the salesman.

# Travelling Salesman Problem (TSP)

The (famous) TSP is the following problem:

- A salesman should visit cities  $c_1, c_2, \dots, c_n$  in some order, visiting each city exactly once and returning to the starting point
- A (positive integer) cost  $d(i, j)$  of travel between each pair  $(c_i, c_j)$  is known.
- Goal: find an optimal (i.e. cheapest) route for the salesman.

Given a graph  $G$  with set  $V$  of vertices ( $|V| = n$ ) and set  $E$  of edges,

- for each vertex  $v$ , create a city  $c_v$ ;
- for each pair of distinct  $u, v \in V$ , set  $d(c_u, c_v) = 1$  if  $uv \in E$  and  $d(c_u, c_v) = 2$  otherwise.

# Travelling Salesman Problem (TSP)

The (famous) TSP is the following problem:

- A salesman should visit cities  $c_1, c_2, \dots, c_n$  in some order, visiting each city exactly once and returning to the starting point
- A (positive integer) cost  $d(i, j)$  of travel between each pair  $(c_i, c_j)$  is known.
- Goal: find an optimal (i.e. cheapest) route for the salesman.

Given a graph  $G$  with set  $V$  of vertices ( $|V| = n$ ) and set  $E$  of edges,

- for each vertex  $v$ , create a city  $c_v$ ;
- for each pair of distinct  $u, v \in V$ , set  $d(c_u, c_v) = 1$  if  $uv \in E$  and  $d(c_u, c_v) = 2$  otherwise.

Then detecting a Hamiltonian cycle in  $G$  can be viewed as TSP:

- if  $G$  has a Hamiltonian cycle then the cycle is a route of cost exactly  $n$ .

# Travelling Salesman Problem (TSP)

The (famous) TSP is the following problem:

- A salesman should visit cities  $c_1, c_2, \dots, c_n$  in some order, visiting each city exactly once and returning to the starting point
- A (positive integer) cost  $d(i, j)$  of travel between each pair  $(c_i, c_j)$  is known.
- Goal: find an optimal (i.e. cheapest) route for the salesman.

Given a graph  $G$  with set  $V$  of vertices ( $|V| = n$ ) and set  $E$  of edges,

- for each vertex  $v$ , create a city  $c_v$ ;
- for each pair of distinct  $u, v \in V$ , set  $d(c_u, c_v) = 1$  if  $uv \in E$  and  $d(c_u, c_v) = 2$  otherwise.

Then detecting a Hamiltonian cycle in  $G$  can be viewed as TSP:

- if  $G$  has a Hamiltonian cycle then the cycle is a route of cost exactly  $n$ .
- if there is a route of cost  $n$  then it can't use pairs with cost 2 (why?) and so goes through edges of  $G$  and hence is a Hamiltonian cycle.

# Travelling Salesman Problem (TSP)

The (famous) TSP is the following problem:

- A salesman should visit cities  $c_1, c_2, \dots, c_n$  in some order, visiting each city exactly once and returning to the starting point
- A (positive integer) cost  $d(i, j)$  of travel between each pair  $(c_i, c_j)$  is known.
- Goal: find an optimal (i.e. cheapest) route for the salesman.

Given a graph  $G$  with set  $V$  of vertices ( $|V| = n$ ) and set  $E$  of edges,

- for each vertex  $v$ , create a city  $c_v$ ;
- for each pair of distinct  $u, v \in V$ , set  $d(c_u, c_v) = 1$  if  $uv \in E$  and  $d(c_u, c_v) = 2$  otherwise.

Then detecting a Hamiltonian cycle in  $G$  can be viewed as TSP:

- if  $G$  has a Hamiltonian cycle then the cycle is a route of cost exactly  $n$ .
- if there is a route of cost  $n$  then it can't use pairs with cost 2 (why?) and so goes through edges of  $G$  and hence is a Hamiltonian cycle.

You have just witnessed a **Reduction!** (from one problem to another).



# Thank You!