

Visualizing all valid witness pairs

Assume we want to show: $5x + 10$ is $O(x)$

Visualizing all valid witness pairs

Assume we want to show: $5x + 10$ is $O(x)$

For any $C > 5$: $5x + 10 \leq C \cdot x$ for $x \geq \frac{10}{C-5}$

Visualizing all valid witness pairs

Assume we want to show: $5x + 10$ is $O(x)$

For any $C > 5$: $5x + 10 \leq C \cdot x$ for $x \geq \frac{10}{C-5}$


The valid witness pairs are $\{(C, k) \mid C > 5, k \geq \frac{10}{C-5}\}$.

Visualizing all valid witness pairs

Assume we want to show: $5x + 10$ is $O(x)$

For any $C > 5$: $5x + 10 \leq C \cdot x$ for $x \geq \frac{10}{C-5}$

The valid witness pairs are $\{(C, k) \mid C > 5, k \geq \frac{10}{C-5}\}$.



Another example

Assume we want to show: $x^2 + 2x + 1$ is $O(x^2)$

Another example

Assume we want to show: $x^2 + 2x + 1$ is $O(x^2)$

For any $C > 1$: $x^2 + 2x + 1 \leq C \cdot x^2$ for $x \geq \frac{2+2\sqrt{C}}{2(C-1)}$

Another example

Assume we want to show: $x^2 + 2x + 1$ is $O(x^2)$

For any $C > 1$: $x^2 + 2x + 1 \leq C \cdot x^2$ for $x \geq \frac{2+2\sqrt{C}}{2(C-1)}$

The valid witness pairs are $\{(C, k) \mid C > 1, k \geq \frac{2+2\sqrt{C}}{2(C-1)}\}$.

Another example

Assume we want to show: $x^2 + 2x + 1$ is $O(x^2)$

For any $C > 1$: $x^2 + 2x + 1 \leq C \cdot x^2$ for $x \geq \frac{2+2\sqrt{C}}{2(C-1)}$

The valid witness pairs are $\{(C, k) \mid C > 1, k \geq \frac{2+2\sqrt{C}}{2(C-1)}\}$.

