

Mathematics for Computer Science
Linear Algebra (Part 2)
Eigen Values & Eigen Vectors

Karl Southern

Durham University

January 20th, 2025

Thanks to Andrei Krokhin and Billy Moses for use of some slides.

Outline

- 1 Recap & Plan for Today
- 2 Understanding eigenvalues and eigenvectors
- 3 Finding eigenvalues and eigenvectors
- 4 Principal Component Analysis (PCA)
- 5 Wrapping Things Up

Name of slide

Last Week

- 1 LU decomposition - what is it/how to use it

This Week

- 1 eigenvalues
- 2 eigenvectors
- 3 Characteristic equation/polynomial
- 4 Principal Component Analysis

Outline

- 1 Recap & Plan for Today
- 2 Understanding eigenvalues and eigenvectors
- 3 Finding eigenvalues and eigenvectors
- 4 Principal Component Analysis (PCA)
- 5 Wrapping Things Up

Intuition

- ① eigen - proper, characteristic, own
- ② Consider some linear mapping A
- ③ When applied to a vector \mathbf{v} , it might cause rotation and scaling
- ④ All those vectors \mathbf{v} that are only scaled: eigenvectors
- ⑤ How much a eigenvector \mathbf{v} is scaled by: corresponding eigenvalue

Eigenvalues and eigenvectors

Definition

Let A be an $n \times n$ matrix. A **non-zero** vector $\mathbf{x} \in \mathbb{R}^n$ is called an **eigenvector** of A (or, equivalently, of the operator $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^n$) if, for some scalar λ ,

$$A\mathbf{x} = \lambda\mathbf{x} \quad (\text{or, equivalently, } T_A(\mathbf{x}) = \lambda\mathbf{x}.)$$

Eigenvalues and eigenvectors

Definition

Let A be an $n \times n$ matrix. A **non-zero** vector $\mathbf{x} \in \mathbb{R}^n$ is called an **eigenvector** of A (or, equivalently, of the operator $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^n$) if, for some scalar λ ,

$$A\mathbf{x} = \lambda\mathbf{x} \quad (\text{or, equivalently, } T_A(\mathbf{x}) = \lambda\mathbf{x}.)$$

In this case, λ is called an **eigenvalue** of A (and of T_A), and \mathbf{x} is an **eigenvector corresponding to λ** .

Eigenvalues and eigenvectors

Definition

Let A be an $n \times n$ matrix. A **non-zero** vector $\mathbf{x} \in \mathbb{R}^n$ is called an **eigenvector** of A (or, equivalently, of the operator $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^n$) if, for some scalar λ ,

$$A\mathbf{x} = \lambda\mathbf{x} \quad (\text{or, equivalently, } T_A(\mathbf{x}) = \lambda\mathbf{x}.)$$

In this case, λ is called an **eigenvalue** of A (and of T_A), and \mathbf{x} is an **eigenvector corresponding to λ** .

- The assumption $\mathbf{x} \neq \mathbf{0}$ is necessary to avoid the case $A\mathbf{0} = \lambda\mathbf{0}$ which always holds.
- The meaning of the notion is that T_A does not change the direction of \mathbf{x} (up to reversal), it only scales \mathbf{x} by λ .

Eigenvalues and eigenvectors

Definition

Let A be an $n \times n$ matrix. A **non-zero** vector $\mathbf{x} \in \mathbb{R}^n$ is called an **eigenvector** of A (or, equivalently, of the operator $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^n$) if, for some scalar λ ,

$$A\mathbf{x} = \lambda\mathbf{x} \quad (\text{or, equivalently, } T_A(\mathbf{x}) = \lambda\mathbf{x}.)$$

Eigenvalues and eigenvectors

Definition

Let A be an $n \times n$ matrix. A **non-zero** vector $\mathbf{x} \in \mathbb{R}^n$ is called an **eigenvector** of A (or, equivalently, of the operator $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^n$) if, for some scalar λ ,

$$A\mathbf{x} = \lambda\mathbf{x} \quad (\text{or, equivalently, } T_A(\mathbf{x}) = \lambda\mathbf{x}.)$$

In this case, λ is called an **eigenvalue** of A (and of T_A), and \mathbf{x} is an **eigenvector corresponding to λ** .

Eigenvalues and eigenvectors

Definition

Let A be an $n \times n$ matrix. A **non-zero** vector $\mathbf{x} \in \mathbb{R}^n$ is called an **eigenvector** of A (or, equivalently, of the operator $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^n$) if, for some scalar λ ,

$$A\mathbf{x} = \lambda\mathbf{x} \quad (\text{or, equivalently, } T_A(\mathbf{x}) = \lambda\mathbf{x}.)$$

In this case, λ is called an **eigenvalue** of A (and of T_A), and \mathbf{x} is an **eigenvector corresponding to λ** .

Example: vector $\mathbf{x} = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$ is an eigenvector of $A = \begin{pmatrix} 3 & 0 \\ 8 & -1 \end{pmatrix}$ corr. to eigenvalue 3.

Indeed,


$$A\mathbf{x} = \begin{pmatrix} 3 & 0 \\ 8 & -1 \end{pmatrix} \begin{pmatrix} 1 \\ 2 \end{pmatrix} = \begin{pmatrix} 3 \\ 6 \end{pmatrix} = 3\mathbf{x}.$$

Example in \mathbb{R}^2

Consider linear operators T_A on \mathbb{R}^2 where A is one of the following matrices:

$$\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

They correspond to **reflections** of \mathbb{R}^2 about y -axis, x -axis, and line $x = y$, resp.




Example in \mathbb{R}^2

Consider linear operators T_A on \mathbb{R}^2 where A is one of the following matrices:

$$\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

They correspond to **reflections** of \mathbb{R}^2 about y -axis, x -axis, and line $x = y$, resp.




- 1 Eigenvectors: all non-zero vectors $(x, 0)$ and $(0, y)$, corr. to eigenvalues -1 and 1 , resp.
- 2 Eigenvectors: all non-zero vectors $(x, 0)$ and $(0, y)$, corr. to eigenvalues 1 and -1 , resp.

Example in \mathbb{R}^2

Consider linear operators T_A on \mathbb{R}^2 where A is one of the following matrices:

$$\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

They correspond to **reflections** of \mathbb{R}^2 about y -axis, x -axis, and line $x = y$, resp.




- 1 Eigenvectors: all non-zero vectors $(x, 0)$ and $(0, y)$, corr. to eigenvalues -1 and 1 , resp.
- 2 Eigenvectors: all non-zero vectors $(x, 0)$ and $(0, y)$, corr. to eigenvalues 1 and -1 , resp.
- 3 Eigenvectors: all non-zero vectors (x, x) and $(-x, x)$, corr. to eigenvalues 1 and -1 , resp.

Example in \mathbb{R}^2

Consider linear operators T_A on \mathbb{R}^2 where A is one of the following matrices:

$$\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}.$$

They correspond to **orthogonal projections** of \mathbb{R}^2 onto x -axis and y -axis, resp.




Example in \mathbb{R}^2

Consider linear operators T_A on \mathbb{R}^2 where A is one of the following matrices:

$$\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}.$$

They correspond to **orthogonal projections** of \mathbb{R}^2 onto x -axis and y -axis, resp.



- 1 Eigenvectors: all non-zero vectors $(x, 0)$ and $(0, y)$, corr. to eigenvalues 1 and 0, resp.
- 2 Eigenvectors: all non-zero vectors $(x, 0)$ and $(0, y)$, corr. to eigenvalues 0 and 1, resp.

Example in \mathbb{R}^2


Consider the linear operator T_A on \mathbb{R}^2 where A is the following matrix:

$$\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}.$$

The corresponding linear map T_A satisfies

$$T_A(x, y) = (x', y') = (x \cos \theta - y \sin \theta, x \sin \theta + y \cos \theta).$$

This corresponds to the **rotation** of \mathbb{R}^2 by angle θ counterclock-wise.



Example in \mathbb{R}^2


Consider the linear operator T_A on \mathbb{R}^2 where A is the following matrix:

$$\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}.$$

The corresponding linear map T_A satisfies

$$T_A(x, y) = (x', y') = (x \cos \theta - y \sin \theta, x \sin \theta + y \cos \theta).$$

This corresponds to the **rotation** of \mathbb{R}^2 by angle θ counterclock-wise.




This linear map has no eigenvectors for any $0 < \theta < 180^\circ$.

Example in \mathbb{R}^2

Consider linear operators T_A on \mathbb{R}^2 where A is the following matrix:

$$\begin{pmatrix} k & 0 \\ 0 & k \end{pmatrix}.$$

This is **contraction** (if $k < 1$) or **dilation** (if $k > 1$) of \mathbb{R}^2 .




Example in \mathbb{R}^2

Consider linear operators T_A on \mathbb{R}^2 where A is the following matrix:

$$\begin{pmatrix} k & 0 \\ 0 & k \end{pmatrix}.$$

This is **contraction** (if $k < 1$) or **dilation** (if $k > 1$) of \mathbb{R}^2 .




The eigenvectors are all non-zero vectors, corresponding to eigenvalue k .

Example in \mathbb{R}^2

Consider linear operators T_A on \mathbb{R}^2 where A is the following matrix:

$$\begin{pmatrix} k & 0 \\ 0 & 1 \end{pmatrix}.$$

They correspond to **compressions** (if $k < 1$) and **expansions** (if $k > 1$) of \mathbb{R}^2 along x-axis.




Example in \mathbb{R}^2

Consider linear operators T_A on \mathbb{R}^2 where A is the following matrix:

$$\begin{pmatrix} k & 0 \\ 0 & 1 \end{pmatrix}.$$

They correspond to **compressions** (if $k < 1$) and **expansions** (if $k > 1$) of \mathbb{R}^2 along x-axis.




The eigenvectors are all non-zero vectors $(x, 0)$ and $(0, y)$, corresponding to eigenvalues k and 1, respectively.

Example in \mathbb{R}^2

Consider the transformation T_A on \mathbb{R}^2 where A is the following matrix:

$$\begin{pmatrix} 1 & k \\ 0 & 1 \end{pmatrix}.$$

The transformation T_A satisfies $T_A(x, y) = (x + ky, y)$. For $k \neq 0$, it corresponds to **shear** of \mathbb{R}^2 in the x -direction with factor k .




Example in \mathbb{R}^2

Consider the transformation T_A on \mathbb{R}^2 where A is the following matrix:

$$\begin{pmatrix} 1 & k \\ 0 & 1 \end{pmatrix}.$$

The transformation T_A satisfies $T_A(x, y) = (x + ky, y)$. For $k \neq 0$, it corresponds to **shear** of \mathbb{R}^2 in the x -direction with factor k .



The eigenvectors are all non-**0** vectors $(x, 0)$, corresponding to eigenvalue 1.

Outline

- 1 Recap & Plan for Today
- 2 Understanding eigenvalues and eigenvectors
- 3 Finding eigenvalues and eigenvectors**
- 4 Principal Component Analysis (PCA)
- 5 Wrapping Things Up

Characteristic equation of a matrix

Theorem

If A is an $n \times n$ matrix then λ is an eigenvalue of A iff it satisfies $\det(\lambda I - A) = 0$.

Characteristic equation of a matrix

Theorem

If A is an $n \times n$ matrix then λ is an eigenvalue of A iff it satisfies $\det(\lambda I - A) = 0$.

The equation $\det(\lambda I - A) = 0$ is called the **characteristic equation** of A .

Characteristic equation of a matrix

Theorem

If A is an $n \times n$ matrix then λ is an eigenvalue of A iff it satisfies $\det(\lambda I - A) = 0$.

The equation $\det(\lambda I - A) = 0$ is called the **characteristic equation** of A .

Proof.

By definition, λ is an eigenvalue of A iff $A\mathbf{x} = \lambda\mathbf{x}$ for some $\mathbf{x} \neq \mathbf{0}$. We have

$$A\mathbf{x} = \lambda\mathbf{x} \quad \Leftrightarrow \quad A\mathbf{x} = \lambda I\mathbf{x} \quad \Leftrightarrow \quad (\lambda I - A)\mathbf{x} = \mathbf{0}.$$

Characteristic equation of a matrix

Theorem

If A is an $n \times n$ matrix then λ is an eigenvalue of A iff it satisfies $\det(\lambda I - A) = 0$.

The equation $\det(\lambda I - A) = 0$ is called the **characteristic equation** of A .

Proof.

By definition, λ is an eigenvalue of A iff $A\mathbf{x} = \lambda\mathbf{x}$ for some $\mathbf{x} \neq \mathbf{0}$. We have

$$A\mathbf{x} = \lambda\mathbf{x} \quad \Leftrightarrow \quad A\mathbf{x} = \lambda I\mathbf{x} \quad \Leftrightarrow \quad (\lambda I - A)\mathbf{x} = \mathbf{0}.$$

By theorem about invertible matrices, the last equation has a solution $\mathbf{x} \neq \mathbf{0}$ iff $\det(\lambda I - A) = 0$. □

Example 12.1

Example: find eigenvalues of the matrix $A = \begin{pmatrix} 2 & -1 \\ 10 & -9 \end{pmatrix}$.

Characteristic polynomial of a matrix

- In general, the expression $\det(\lambda I - A)$ is a polynomial

$$p(\lambda) = \lambda^n + c_1\lambda^{n-1} + \dots + c_{n-1}\lambda + c_n.$$

where n is the order of A . It is called the **characteristic polynomial** of A .

Characteristic polynomial of a matrix

- In general, the expression $\det(\lambda I - A)$ is a polynomial

$$p(\lambda) = \lambda^n + c_1\lambda^{n-1} + \dots + c_{n-1}\lambda + c_n.$$

where n is the order of A . It is called the **characteristic polynomial** of A .

- Solving the equation $p(\lambda) = 0$ is difficult in general — there is no closed formula or exact algorithm.
- There are numerical algorithms for computing eigenvalues approximately.

Characteristic polynomial of a matrix

- In general, the expression $\det(\lambda I - A)$ is a polynomial

$$p(\lambda) = \lambda^n + c_1\lambda^{n-1} + \dots + c_{n-1}\lambda + c_n.$$

where n is the order of A . It is called the **characteristic polynomial** of A .

- Solving the equation $p(\lambda) = 0$ is difficult in general — there is no closed formula or exact algorithm.
- There are numerical algorithms for computing eigenvalues approximately.
- If all coefficients of $p(\lambda)$ are integers and the equation $p(\lambda) = 0$ has an integer solution $\lambda = k$ then $k|c_n$. This can be used to find some eigenvalues.

Example 12.2

Example: find eigenvalues of $A = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 4 & -17 & 8 \end{pmatrix}$.

Eigenspaces and their bases

- Let λ_0 be an eigenvalue of A and consider the equation $(\lambda_0 I - A)\mathbf{x} = \mathbf{0}$.
- The null space of $\lambda_0 I - A$ is called the eigenspace of A corresponding to λ_0 .

Eigenspaces and their bases

- Let λ_0 be an eigenvalue of A and consider the equation $(\lambda_0 I - A)\mathbf{x} = \mathbf{0}$.
- The null space of $\lambda_0 I - A$ is called the eigenspace of A corresponding to λ_0 .
- The non- $\mathbf{0}$ vectors in this space are the eigenvectors of A corresponding to λ_0 .

Eigenspaces and their bases

- Let λ_0 be an eigenvalue of A and consider the equation $(\lambda_0 I - A)\mathbf{x} = \mathbf{0}$.
- The null space of $\lambda_0 I - A$ is called the eigenspace of A corresponding to λ_0 .
- The non- $\mathbf{0}$ vectors in this space are the eigenvectors of A corresponding to λ_0 .
- To find a basis in this subspace, use the algorithm for finding a basis in the null space of a matrix.

Eigenspaces and their bases

- Let λ_0 be an eigenvalue of A and consider the equation $(\lambda_0 I - A)\mathbf{x} = \mathbf{0}$.
- The null space of $\lambda_0 I - A$ is called the eigenspace of A corresponding to λ_0 .
- The non- $\mathbf{0}$ vectors in this space are the eigenvectors of A corresponding to λ_0 .
- To find a basis in this subspace, use the algorithm for finding a basis in the null space of a matrix.

Find (a basis of) the eigenspace of $A = \begin{pmatrix} 2 & -1 \\ 10 & -9 \end{pmatrix}$ corresponding to $\lambda = -8$.

Eigenspaces and their bases

- Let λ_0 be an eigenvalue of A and consider the equation $(\lambda_0 I - A)\mathbf{x} = \mathbf{0}$.
- The null space of $\lambda_0 I - A$ is called the eigenspace of A corresponding to λ_0 .
- The non- $\mathbf{0}$ vectors in this space are the eigenvectors of A corresponding to λ_0 .
- To find a basis in this subspace, use the algorithm for finding a basis in the null space of a matrix.

Find (a basis of) the eigenspace of $A = \begin{pmatrix} 2 & -1 \\ 10 & -9 \end{pmatrix}$ corresponding to $\lambda = -8$.

Solution. Form the equation $(-8I - A)\mathbf{x} = \mathbf{0}$, or

$$\begin{pmatrix} -10 & 1 \\ -10 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad \text{or} \quad \begin{array}{rcl} -10x_1 + x_2 & = & 0 \\ -10x_1 + x_2 & = & 0 \end{array}$$

Eigenspaces and their bases

- Let λ_0 be an eigenvalue of A and consider the equation $(\lambda_0 I - A)\mathbf{x} = \mathbf{0}$.
- The null space of $\lambda_0 I - A$ is called the eigenspace of A corresponding to λ_0 .
- The non- $\mathbf{0}$ vectors in this space are the eigenvectors of A corresponding to λ_0 .
- To find a basis in this subspace, use the algorithm for finding a basis in the null space of a matrix.

Find (a basis of) the eigenspace of $A = \begin{pmatrix} 2 & -1 \\ 10 & -9 \end{pmatrix}$ corresponding to $\lambda = -8$.

Solution. Form the equation $(-8I - A)\mathbf{x} = \mathbf{0}$, or

$$\begin{pmatrix} -10 & 1 \\ -10 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad \text{or} \quad \begin{array}{rcl} -10x_1 + x_2 & = & 0 \\ -10x_1 + x_2 & = & 0 \end{array}$$

The subspace consists of all vectors of the form $(x, 10x)$. One basis is $\{(1, 10)\}$.

Eigenspaces and their bases

- Let λ_0 be an eigenvalue of A and consider the equation $(\lambda_0 I - A)\mathbf{x} = \mathbf{0}$.
- The null space of $\lambda_0 I - A$ is called the eigenspace of A corresponding to λ_0 .
- The non- $\mathbf{0}$ vectors in this space are the eigenvectors of A corresponding to λ_0 .
- To find a basis in this subspace, use the algorithm for finding a basis in the null space of a matrix.

Find (a basis of) the eigenspace of $A = \begin{pmatrix} 2 & -1 \\ 10 & -9 \end{pmatrix}$ corresponding to $\lambda = -8$.

Solution. Form the equation $(-8I - A)\mathbf{x} = \mathbf{0}$, or

$$\begin{pmatrix} -10 & 1 \\ -10 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad \text{or} \quad \begin{array}{rcl} -10x_1 + x_2 & = & 0 \\ -10x_1 + x_2 & = & 0 \end{array}$$

The subspace consists of all vectors of the form $(x, 10x)$. One basis is $\{(1, 10)\}$.

Exercise: Find the eigenspace of A corresponding to eigenvalue $\lambda = 1$.

Multiplicities of an eigenvalue

Let λ_0 be an eigenvalue of a matrix A .

- The **algebraic multiplicity** of λ_0 is the power k with which $(\lambda - \lambda_0)$ appears as a factor of $\det(\lambda I - A)$ - the characteristic polynomial of A .
 - ▶ E.g. if $\det(\lambda I - A) = (\lambda - 2)^3 \cdot (\lambda + 5)^2 \cdots$, then it's 3 for 2 and 2 for -5

Multiplicities of an eigenvalue

Let λ_0 be an eigenvalue of a matrix A .

- The **algebraic multiplicity** of λ_0 is the power k with which $(\lambda - \lambda_0)$ appears as a factor of $\det(\lambda I - A)$ - the characteristic polynomial of A .
 - ▶ E.g. if $\det(\lambda I - A) = (\lambda - 2)^3 \cdot (\lambda + 5)^2 \cdots$, then it's 3 for 2 and 2 for -5
- The **geometric multiplicity** of λ_0 is the dimension of the eigenspace corresponding to λ_0 .

Multiplicities of an eigenvalue

Let λ_0 be an eigenvalue of a matrix A .

- The **algebraic multiplicity** of λ_0 is the power k with which $(\lambda - \lambda_0)$ appears as a factor of $\det(\lambda I - A)$ - the characteristic polynomial of A .
 - ▶ E.g. if $\det(\lambda I - A) = (\lambda - 2)^3 \cdot (\lambda + 5)^2 \cdots$, then it's 3 for 2 and 2 for -5
- The **geometric multiplicity** of λ_0 is the dimension of the eigenspace corresponding to λ_0 .

Theorem

Let A be any square matrix. For every eigenvalue of A , its algebraic multiplicity is greater than or equal to its geometric multiplicity. (Proof omitted)

Multiplicities of an eigenvalue

Let λ_0 be an eigenvalue of a matrix A .

- The **algebraic multiplicity** of λ_0 is the power k with which $(\lambda - \lambda_0)$ appears as a factor of $\det(\lambda I - A)$ - the characteristic polynomial of A .
 - ▶ E.g. if $\det(\lambda I - A) = (\lambda - 2)^3 \cdot (\lambda + 5)^2 \cdots$, then it's 3 for 2 and 2 for -5
- The **geometric multiplicity** of λ_0 is the dimension of the eigenspace corresponding to λ_0 .

Theorem

Let A be any square matrix. For every eigenvalue of A , its algebraic multiplicity is greater than or equal to its geometric multiplicity. (Proof omitted)

Exercise: Find an example of A and its eigenvalue where the inequality in the theorem is strict.

Outline

- 1 Recap & Plan for Today
- 2 Understanding eigenvalues and eigenvectors
- 3 Finding eigenvalues and eigenvectors
- 4 Principal Component Analysis (PCA)**
- 5 Wrapping Things Up

Purpose of This Section

- ① PCA - Why study it?
 - ① Useful applications - data compression
 - ② Uses eigenvalues and eigenvectors

Purpose of This Section

- ① PCA - Why study it?
 - ① Useful applications - data compression
 - ② Uses eigenvalues and eigenvectors
- ② What we care about:
 - ① General idea of PCA
 - ② Understanding how eigenvalues and eigenvectors are used

Purpose of This Section

- ① PCA - Why study it?
 - ① Useful applications - data compression
 - ② Uses eigenvalues and eigenvectors
- ② What we care about:
 - ① General idea of PCA
 - ② Understanding how eigenvalues and eigenvectors are used
- ③ What we don't care about:
 - ① Completely understanding PCA
 - ② Learning concepts used in PCA that are beyond the scope of this module

PCA - what is it?

① Principal Component Analysis

PCA - what is it?

- ① Principal Component Analysis
- ② eigen meaning: characteristic

PCA - what is it?

- ① Principal Component Analysis
- ② eigen meaning: characteristic
- ③ So...finding the principal (most important) components (eigenvectors)...of some data set.

PCA - what is it?

- ① Principal Component Analysis
- ② eigen meaning: characteristic
- ③ So...finding the principal (most important) components (eigenvectors)...of some data set.
- ④ Data set? But couldn't we only find eigenvectors for square matrices?

PCA - what is it?

- ① Principal Component Analysis
- ② eigen meaning: characteristic
- ③ So...finding the principal (most important) components (eigenvectors)...of some data set.
- ④ Data set? But couldn't we only find eigenvectors for square matrices?
- ⑤ Concept beyond scope of module: **covariance matrix**

PCA - what is it?


- ① Principal Component Analysis
- ② eigen meaning: characteristic
- ③ So...finding the principal (most important) components (eigenvectors)...of some data set.
- ④ Data set? But couldn't we only find eigenvectors for square matrices?
- ⑤ Concept beyond scope of module: **covariance matrix**
- ⑥ What does it do?

PCA - what is it?

- ① Principal Component Analysis
- ② eigen meaning: characteristic
- ③ So...finding the principal (most important) components (eigenvectors)...of some data set.
- ④ Data set? But couldn't we only find eigenvectors for square matrices?
- ⑤ Concept beyond scope of module: **covariance matrix**
- ⑥ What does it do?
 - ▶ **Input:** data set in n dimensions (possibly m data points, $m \neq n$)
 - ▶ **Output:** $n \times n$ matrix, element e_{ij} shows how dimension i data varies with dimension j data.
 - ▶ Captures the **variance** of one dimension of the data with another.

Visual representation of principal components of data set (eigenvectors of its covariance matrix)

- Data set: m points of 2 dimensions (x & y); Covariance matrix $COV_{2 \times 2}$
- Eigenvectors of $COV_{2 \times 2}$ (aka principal axes) - axes which best capture variance of data
- Eigenvalues of $COV_{2 \times 2}$ (aka explained variance) - amount of variance along eigenvectors



Given an $n \times n$ covariance matrix, how to find principal components

- 1 Pretend we have some covariance matrix $COV_{n \times n}$.

Given an $n \times n$ covariance matrix, how to find principal components

- 1 Pretend we have some covariance matrix $COV_{n \times n}$.
- 2 Find the n eigenvalues & corresponding eigenvectors of $COV_{n \times n}$

Given an $n \times n$ covariance matrix, how to find principal components

- 1 Pretend we have some covariance matrix $COV_{n \times n}$.
- 2 Find the n eigenvalues & corresponding eigenvectors of $COV_{n \times n}$
- 3 Order eigenvalues in decreasing order.

How much does each eigenvalue contribute to total explained variance/how important is each eigenvalue?

Given an $n \times n$ covariance matrix, how to find principal components

- 1 Pretend we have some covariance matrix $COV_{n \times n}$.
- 2 Find the n eigenvalues & corresponding eigenvectors of $COV_{n \times n}$
- 3 Order eigenvalues in decreasing order.
How much does each eigenvalue contribute to total explained variance/how important is each eigenvalue?
- 4 E.g., eigenvalues 6, 4, 2.
6 contributes $6/(6 + 4 + 2) = .5$, 4 contributes .33, and 2 contributes .17.

Given an $n \times n$ covariance matrix, how to find principal components

- 1 Pretend we have some covariance matrix $COV_{n \times n}$.
- 2 Find the n eigenvalues & corresponding eigenvectors of $COV_{n \times n}$
- 3 Order eigenvalues in decreasing order.
How much does each eigenvalue contribute to total explained variance/how important is each eigenvalue?
- 4 E.g., eigenvalues 6, 4, 2.
6 contributes $6/(6 + 4 + 2) = .5$, 4 contributes .33, and 2 contributes .17.
- 5 If capturing 80% of the “explained variance” is good enough, then using just first two eigenvalues is enough (totally .83)

Given an $n \times n$ covariance matrix, how to find principal components

- 1 Pretend we have some covariance matrix $COV_{n \times n}$.
- 2 Find the n eigenvalues & corresponding eigenvectors of $COV_{n \times n}$
- 3 Order eigenvalues in decreasing order.
How much does each eigenvalue contribute to total explained variance/how important is each eigenvalue?
- 4 E.g., eigenvalues 6, 4, 2.
6 contributes $6/(6 + 4 + 2) = .5$, 4 contributes .33, and 2 contributes .17.
- 5 If capturing 80% of the “explained variance” is good enough, then using just first two eigenvalues is enough (totally .83)
- 6 Let $P_{n \times k}$ be matrix with k chosen eigenvectors as column vectors

Given an $n \times n$ covariance matrix, how to find principal components

- 1 Pretend we have some covariance matrix $COV_{n \times n}$.
- 2 Find the n eigenvalues & corresponding eigenvectors of $COV_{n \times n}$
- 3 Order eigenvalues in decreasing order.
How much does each eigenvalue contribute to total explained variance/how important is each eigenvalue?
- 4 E.g., eigenvalues 6, 4, 2.
6 contributes $6/(6 + 4 + 2) = .5$, 4 contributes .33, and 2 contributes .17.
- 5 If capturing 80% of the “explained variance” is good enough, then using just first two eigenvalues is enough (totally .83)
- 6 Let $P_{n \times k}$ be matrix with k chosen eigenvectors as column vectors
- 7 Now we can represent original data (n dimensions) using less data (k dimensions) while possibly losing some information

PCA - high level

- 1 Preprocessing step: put data set (m rows of n dimensions) into “good form”, call $A_{m \times n}$

PCA - high level

- 1 Preprocessing step: put data set (m rows of n dimensions) into “good form”, call $A_{m \times n}$
- 2 For $A_{m \times n}$, find its covariance matrix $COV_{n \times n}$

PCA - high level

- 1 Preprocessing step: put data set (m rows of n dimensions) into “good form”, call $A_{m \times n}$
- 2 For $A_{m \times n}$, find its covariance matrix $COV_{n \times n}$
- 3 Find the n eigenvalues & corresponding eigenvectors of $COV_{n \times n}$ and choose k principal components

PCA - high level

- 1 Preprocessing step: put data set (m rows of n dimensions) into “good form”, call $A_{m \times n}$
- 2 For $A_{m \times n}$, find its covariance matrix $COV_{n \times n}$
- 3 Find the n eigenvalues & corresponding eigenvectors of $COV_{n \times n}$ and choose k principal components
- 4 Let $P_{n \times k}$ be matrix with k eigenvectors as column vectors

PCA - high level

- 1 Preprocessing step: put data set (m rows of n dimensions) into “good form”, call $A_{m \times n}$
- 2 For $A_{m \times n}$, find its covariance matrix $COV_{n \times n}$
- 3 Find the n eigenvalues & corresponding eigenvectors of $COV_{n \times n}$ and choose k principal components
- 4 Let $P_{n \times k}$ be matrix with k eigenvectors as column vectors
- 5 Put data into new form: $A_{m \times n} P_{n \times k} = B_{m \times k}$

PCA - high level

- 1 Preprocessing step: put data set (m rows of n dimensions) into “good form”, call $A_{m \times n}$
- 2 For $A_{m \times n}$, find its covariance matrix $COV_{n \times n}$
- 3 Find the n eigenvalues & corresponding eigenvectors of $COV_{n \times n}$ and choose k principal components
- 4 Let $P_{n \times k}$ be matrix with k eigenvectors as column vectors
- 5 Put data into new form: $A_{m \times n} P_{n \times k} = B_{m \times k}$
- 6 Can retrieve old data (possibly with some loss):
 $A'_{m \times n} = B_{m \times k} P_{k \times n}^T$, then “un-preprocess A' ”

Outline

- 1 Recap & Plan for Today
- 2 Understanding eigenvalues and eigenvectors
- 3 Finding eigenvalues and eigenvectors
- 4 Principal Component Analysis (PCA)
- 5 Wrapping Things Up

What we learnt today

- Eigenvalues and eigenvectors of matrices
- Examples in \mathbb{R}^2
- Characteristic equation of a matrix – how to find eigenvalues
- Eigenspaces and how to find their bases
- Principal Component Analysis (PCA)

Next Week:

- Complex vector spaces.
- If you're unfamiliar with complex numbers, read up on them on this [Wikipedia page](#) and/or watch this [video lecture](#) by 3Blue1Brown in advance of the lecture.

Example 12.1

Example: find eigenvalues of the matrix $A = \begin{pmatrix} 2 & -1 \\ 10 & -9 \end{pmatrix}$.

Example 12.1

Example: find eigenvalues of the matrix $A = \begin{pmatrix} 2 & -1 \\ 10 & -9 \end{pmatrix}$. We have

$$\det(\lambda I - A) = \begin{vmatrix} \lambda - 2 & 1 \\ -10 & \lambda + 9 \end{vmatrix} = (\lambda - 2) \cdot (\lambda + 9) - 1 \cdot (-10) = \lambda^2 + 7\lambda - 8.$$

Example 12.1

Example: find eigenvalues of the matrix $A = \begin{pmatrix} 2 & -1 \\ 10 & -9 \end{pmatrix}$. We have

$$\det(\lambda I - A) = \begin{vmatrix} \lambda - 2 & 1 \\ -10 & \lambda + 9 \end{vmatrix} = (\lambda - 2) \cdot (\lambda + 9) - 1 \cdot (-10) = \lambda^2 + 7\lambda - 8.$$

So, the characteristic equation of A is $\lambda^2 + 7\lambda - 8 = 0$.

Example 12.1

Example: find eigenvalues of the matrix $A = \begin{pmatrix} 2 & -1 \\ 10 & -9 \end{pmatrix}$. We have

$$\det(\lambda I - A) = \begin{vmatrix} \lambda - 2 & 1 \\ -10 & \lambda + 9 \end{vmatrix} = (\lambda - 2) \cdot (\lambda + 9) - 1 \cdot (-10) = \lambda^2 + 7\lambda - 8.$$

So, the characteristic equation of A is $\lambda^2 + 7\lambda - 8 = 0$.

Its solutions $\lambda_1 = 1$ and $\lambda_2 = -8$ are the eigenvalues of A .

Example 12.1

Example: find eigenvalues of the matrix $A = \begin{pmatrix} 2 & -1 \\ 10 & -9 \end{pmatrix}$. We have

$$\det(\lambda I - A) = \begin{vmatrix} \lambda - 2 & 1 \\ -10 & \lambda + 9 \end{vmatrix} = (\lambda - 2) \cdot (\lambda + 9) - 1 \cdot (-10) = \lambda^2 + 7\lambda - 8.$$

So, the characteristic equation of A is $\lambda^2 + 7\lambda - 8 = 0$.

Its solutions $\lambda_1 = 1$ and $\lambda_2 = -8$ are the eigenvalues of A .

Example: find eigenvalues of the matrix $B = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$.

Example 12.1

Example: find eigenvalues of the matrix $A = \begin{pmatrix} 2 & -1 \\ 10 & -9 \end{pmatrix}$. We have

$$\det(\lambda I - A) = \begin{vmatrix} \lambda - 2 & 1 \\ -10 & \lambda + 9 \end{vmatrix} = (\lambda - 2) \cdot (\lambda + 9) - 1 \cdot (-10) = \lambda^2 + 7\lambda - 8.$$

So, the characteristic equation of A is $\lambda^2 + 7\lambda - 8 = 0$.

Its solutions $\lambda_1 = 1$ and $\lambda_2 = -8$ are the eigenvalues of A .

Example: find eigenvalues of the matrix $B = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. We have

$$\det(\lambda I - B) = \begin{vmatrix} \lambda & 1 \\ -1 & \lambda \end{vmatrix} = \lambda^2 + 1.$$

Example 12.1

Example: find eigenvalues of the matrix $A = \begin{pmatrix} 2 & -1 \\ 10 & -9 \end{pmatrix}$. We have

$$\det(\lambda I - A) = \begin{vmatrix} \lambda - 2 & 1 \\ -10 & \lambda + 9 \end{vmatrix} = (\lambda - 2) \cdot (\lambda + 9) - 1 \cdot (-10) = \lambda^2 + 7\lambda - 8.$$

So, the characteristic equation of A is $\lambda^2 + 7\lambda - 8 = 0$.

Its solutions $\lambda_1 = 1$ and $\lambda_2 = -8$ are the eigenvalues of A .

Example: find eigenvalues of the matrix $B = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. We have

$$\det(\lambda I - B) = \begin{vmatrix} \lambda & 1 \\ -1 & \lambda \end{vmatrix} = \lambda^2 + 1.$$

The characteristic equation of B is $\lambda^2 + 1 = 0$, so B has no (real) eigenvalues.

Example 12.2

Example: find eigenvalues of $A = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 4 & -17 & 8 \end{pmatrix}$.

Example 12.2

Example: find eigenvalues of $A = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 4 & -17 & 8 \end{pmatrix}$. We have

$$\det(\lambda I - A) = \begin{vmatrix} \lambda & -1 & 0 \\ 0 & \lambda & -1 \\ -4 & 17 & \lambda - 8 \end{vmatrix} = \lambda^3 - 8\lambda^2 + 17\lambda - 4 = 0.$$

Example 12.2

Example: find eigenvalues of $A = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 4 & -17 & 8 \end{pmatrix}$. We have

$$\det(\lambda I - A) = \begin{vmatrix} \lambda & -1 & 0 \\ 0 & \lambda & -1 \\ -4 & 17 & \lambda - 8 \end{vmatrix} = \lambda^3 - 8\lambda^2 + 17\lambda - 4 = 0.$$

If A has an integer eigenvalue then it is a divisor of -4 , i.e., $\pm 1, \pm 2, \pm 4$.

Example 12.2

Example: find eigenvalues of $A = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 4 & -17 & 8 \end{pmatrix}$. We have

$$\det(\lambda I - A) = \begin{vmatrix} \lambda & -1 & 0 \\ 0 & \lambda & -1 \\ -4 & 17 & \lambda - 8 \end{vmatrix} = \lambda^3 - 8\lambda^2 + 17\lambda - 4 = 0.$$

If A has an integer eigenvalue then it is a divisor of -4 , i.e., $\pm 1, \pm 2, \pm 4$. Checking these numbers in turn, we find that $\lambda = 4$ is a solution.

Example 12.2

Example: find eigenvalues of $A = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 4 & -17 & 8 \end{pmatrix}$. We have

$$\det(\lambda I - A) = \begin{vmatrix} \lambda & -1 & 0 \\ 0 & \lambda & -1 \\ -4 & 17 & \lambda - 8 \end{vmatrix} = \lambda^3 - 8\lambda^2 + 17\lambda - 4 = 0.$$

If A has an integer eigenvalue then it is a divisor of -4 , i.e., $\pm 1, \pm 2, \pm 4$. Checking these numbers in turn, we find that $\lambda = 4$ is a solution.

Divide $\lambda^3 - 8\lambda^2 + 17\lambda - 4$ by $\lambda - 4$ to get

$$\lambda^3 - 8\lambda^2 + 17\lambda - 4 = (\lambda - 4)(\lambda^2 - 4\lambda + 1).$$

Solving the equation $\lambda^2 - 4\lambda + 1 = 0$, we get that the eigenvalues of A are

$$\lambda_1 = 4, \lambda_2 = 2 + \sqrt{3}, \text{ and } \lambda_3 = 2 - \sqrt{3}.$$

The End