## **Quantum Schrödinger Functions**

by Sven Nilsen, 2020

*In this paper I formalize Quantum Schrödinger Functions and introduce the Phi quantum function.* 

The time-dependent Schrödinger equation for a  $\dot{\psi}$  wavefunction over a Hilbert space:

$$i\hbar\psi'(t)=H\psi(t)$$
 time-dependent Schrödinger equation  $\psi: time \rightarrow \mathbb{H}$ 

The type of complex numbers is a Hilbert space with the inner product:

```
 \langle a, b \rangle <=> ab^* 
 \langle \mathbb{C}, \operatorname{mul}_{\mathbb{C}} \cdot (\operatorname{id}, \operatorname{conj})) : [\operatorname{hilbert\_space}] \text{ true} 
 \langle \operatorname{conj}(x) = \operatorname{re}(x) - \operatorname{im}(x)i
```

A quantum Schrödinger function `f` is a higher order quantum function depending on time:

```
f: time \to T where `T` is usually `Bn` or `Rn` and `T^0 <=> ()` \exists_{cp} f(t) <=> \psi(t) \psi: time \to T \to \mathbb{C} `\psi` satisfies time-dependent Schrödinger for all `T`
```

The Phi quantum function is a basic building block for quantum Schrödinger functions:

```
phi : frequency \rightarrow time \rightarrow ()

\exists_{cp}phi(f)(t) <=> \varphi(f)(t)

\varphi(f)(t) = \sin(2\pi ft) + i\cos(2\pi ft)

\varphi(f)'(t) = (2\pi f)(\cos(2\pi ft) - i\sin(2\pi ft))
```

Observed alone, `phi` always returns, but nevertheless it is has a complex probability amplitude. The function  $\varphi(f)(t)$  simply rotates around in a unit circle at frequency `f`.

One can derive the analogue of the Schrödinger equation for  $\hat{\varphi}(t)$ :

```
∴ hφ(f)'(t) = (hf)(cos(2πft) - isin(2πft))
∴ ihφ(f)'(t) = (hf)(isin(2πft) - i²cos(2πft))
∴ ihφ(f)'(t) = Eφ(f)(t)

∴ h = h / 2π reduced Planck's constant
∴ E = hf Planck-Einstein relation
```

The Hamiltonian `H` is a measure of the total energy, so Phi is a quantum Schrödinger function.