HLA−II PTM figures: supplementary set

Caron Lab / Yale · JY cell line, DDA

These are extra panels beyond the set you asked for (1A−B, 2, 4A−B, 6E are in the first−pass PDF). Same data and same pipeline, in case any of them are useful for the talk or the paper.

- Built from the same processed tables that already match your embedded summaries (my Strong+Weak equals your All count for every PTM).
- 6A is the one exception: I went back to the raw Phospho sheet for it, because the pipeline only keeps the first phospho site per peptide and 6A needs all of them. The phospho count there (818) still matches the pipeline.
- Length 10−25 mer, sequence variants kept separate, same as the first pass.
- 14 HLA−II allele pairs (DP/DQ/DR), 14 PTMs including Nitrosylation.

One caveat on 5A: the HLA−I version only had 3 alleles to spread peptides over; here there are 14, and each peptide only counts toward its best allele, so the per−cell numbers get thin. I dropped any cell with fewer than 10 peptides (left grey). The by−PTM panels pool the residues together, so the signal still comes through where the by−residue panels get sparse.

Panel	Figure
5A	Strong binders by PTM x allele (heatmap)
5A	Weak binders by PTM x allele (heatmap)
5A	Strong binders by PTM+residue x allele (heatmap)
5A	Weak binders by PTM+residue x allele (heatmap)
5B	EL−rank distribution, modified vs background (14 PTMs)
6A	Doubly+ phosphorylated peptides: site position distribution
6B	Ubiquitination G vs GG
6C	Oxidation: biological vs artifact
6D	Biological oxidation: binding by oxidized residue
6D2	Biological oxidation: %Strong by residue x allele (heatmap)

Goes with the first−pass PDF. Tell me what you want changed. − Peter