

Mathematical Foundations of Open System Dissipation, Hyperbolicity, SRB Measures, and Adelic Dynamics

Tracing out environmental degrees of freedom in an open system yields an effective reduced dynamics that transitions from a Hamiltonian, phase-space volume-preserving evolution to a non-unitary, dissipative flow. Establishing rigorously that such a reduced dynamical system admits a unique compact global attractor equipped with a Sinai-Ruelle-Bowen (SRB) physical measure requires integrating foundational results across open quantum systems, smooth ergodic theory, non-Archimedean dynamics, and geometric measure theory. This research report surveys the mathematical literature across six dependent domains, mapping established theorems, evaluating their applicability to Bohmian and number-theoretic formulations, and identifying critical theoretical gaps.

1. Deriving Dissipation from a Unitary Theory

Understanding how open system dynamics transitions from a reversible Hamiltonian system to an irreversible, phase-space volume-contracting flow requires rigorous mathematical derivation of open quantum systems.

Literature Map and Summaries

The foundational mathematical formulation of environmental tracing originates with Feynman and Vernon (1963), who established the influence-functional formalism¹. By expressing a quantum system coupled to a linear environmental bath through path integrals, the environmental degrees of freedom are integrated out to yield an effective action

$S_{\text{eff}}[q(t), q'(t)] = S_0[q] - S_0[q'] + \Phi[q, q']$, where the complex influence functional $\exp(i\Phi/\hbar)$ contains non-local temporal kernels encoding decoherence and dissipation¹.

Caldeira and Leggett (1983) applied this framework to quantum Brownian motion by coupling a

system coordinate q to an infinite collection of harmonic oscillators via position-position

interactions $q \sum_k C_k q_k$ ¹. Under the assumptions of high temperature ($k_B T \gg \hbar \Omega$) and

an Ohmic spectral density $J(\omega) = \eta \omega$, evaluating the path integral yields a Master Equation

for the reduced density matrix $\rho(q, q', t)$ ¹. This equation contains both a friction term

proportional to the classical damping constant η and a momentum diffusion term driven by thermal fluctuations¹.

Davies (1974) provided the rigorous mathematical foundation for Markovian master equations

via the weak-coupling limit⁸. Davies proved that if the coupling constant $\lambda \rightarrow 0$ while time is rescaled as $\tau = \lambda^2 t$, the reduced Heisenberg/Schrödinger evolution converges in norm to a Completely Positive Trace-Preserving (CPTP) quantum dynamical semigroup generated by a Gorini-Kossakowski-Sudarshan-Lindblad (GKLS) operator⁸.

Real et al. (2017) conducted a rigorous mathematical analysis of the Caldeira-Leggett limit and demonstrated an essential structural limitation⁶. They proved that taking the Markovian limit under standard position-position coupling yields a master equation that either violates complete positivity at short times or fails to generate true Markovian phase-space volume contraction⁶. They established that strictly dissipative Markovian dynamics require non-Markovian memory structures or explicit position-momentum bath couplings⁶.

Jakšić and Pillet (1996) developed the spectral theory of thermal relaxation for Pauli-Fierz models, avoiding Born-Markov approximations¹⁴. Using complex deformation techniques on thermal Liouvillians, they proved return to equilibrium and exponential decay of correlations for a small system coupled to an infinite thermal reservoir¹².

Structural Applicability and Required Extensions

The weak-coupling limits of Davies and the spectral deformation methods of Jakšić and Pillet operate at the Hilbert-space operator level or C*-algebraic level, proving convergence of density matrices $\rho(t) \rightarrow \rho_{\text{th}}$ ⁸. Applying these results to prove phase-space volume contraction ($\text{div } v < 0$) in deterministic Bohmian configuration space requires a non-trivial extension. The reduced wave function or density matrix $\rho(q, q', t)$ must be projected into configuration space via a polar decomposition $\psi = R e^{iS/\hbar}$ to yield an effective Bohmian velocity field $v(q, t) = \frac{\hbar}{m} \nabla S(q, t)$ ¹⁶. One must demonstrate that the environmental influence functional reduces strictly to a deterministic negative-divergence drift field rather than purely stochastic white noise.

Assessment of Critical Open Gap

The primary literature gap in Area 1 is the explicit, mathematically rigorous derivation of deterministic vector fields exhibiting uniform or singular phase-space volume contraction from an underlying open quantum influence functional. Existing literature yields either CPTP quantum semigroups on operator algebras⁸ or stochastic Langevin equations⁵. A formal bridging theorem mapping influence-functional kernel parameters directly to deterministic vector fields with provably negative divergence in configuration space remains unestablished.

2. Hyperbolicity of Dissipative Dynamical Systems

Once a deterministic dissipative vector field is established, proving that its flow possesses a

well-defined chaotic attractor requires validating structural hyperbolicity conditions.

Literature Map and Summaries

Morales, Pacifico, and Pujals (1999) introduced singular hyperbolicity to formalize non-uniformly hyperbolic flows that contain equilibria, such as the Lorenz attractor¹⁸. A compact invariant set

Λ for a vector field X is singular-hyperbolic if all equilibria in Λ are hyperbolic saddle points,

and the tangent bundle over Λ admits a continuous $D\phi_t$ -invariant decomposition

$T_\Lambda M = E^s \oplus E^{cu}$, where E^s is uniformly contracting and E^{cu} is sectionally (or volume) expanding¹⁸.

Tucker (1999) resolved Smale's 14th problem by proving that the classical Lorenz equations generate a singular-hyperbolic chaotic attractor¹⁹. Tucker's proof combined rigorous computer-assisted bounds (using interval arithmetic and Taylor series enclosures) with

geometric dynamical systems theory to construct a trapping region U and verify cone-field preservation across Poincaré return maps¹⁹.

Araujo, Pacifico, Pujals, and Viana (2005) proved that any singular-hyperbolic attractor for a three-dimensional flow is expansive and supports a physical (SRB) measure whose ergodic basin covers Lebesgue-almost all points in the topological basin of attraction¹⁸.

Alves, Soufi, and Araujo (2020) proved the statistical stability of singular-hyperbolic attractors, showing that physical/SRB measures depend continuously on vector field perturbations in the weak-* topology²⁰. This establishes that physical observables remain robust under small structural shifts in the governing equations²⁰.

Structural Applicability and Required Extensions

Standard singular hyperbolicity applies directly to three-dimensional flows containing saddle equilibria¹⁸. However, if environmental tracing yields system dimensions higher than three (

$\dim \geq 4$) or infinite-dimensional Hilbert-space projection manifolds, 3D singular hyperbolicity is insufficient¹⁹. The theory must be extended to *sectional hyperbolicity* in higher dimensions,

requiring that the central-unstable bundle E^{cu} has dimension $\dim(E^{cu}) \geq 2$ and that

every two-dimensional subspace within E^{cu} is exponentially volume-expanding under $D\phi_t$ ¹⁹.

Assessment of Critical Open Gap

The primary gap in Area 2 is the scarcity of analytical (non-computer-assisted) methods for proving singular or sectional hyperbolicity in continuous-time non-linear systems. Tucker's proof relies on computer-assisted interval arithmetic localized to specific numerical parameters of the Lorenz system¹⁹. General functional criteria that guarantee sectional hyperbolicity directly from differential equations without computer assistance remain largely unknown.

3. SRB Measure Existence, Ergodicity, and Dimension Computation

Given a hyperbolic or singular-hyperbolic attractor, physical predictions require constructing invariant Sinai-Ruelle-Bowen (SRB) measures and calculating their fractal dimensions.

Literature Map and Summaries

Bowen and Ruelle (1975) established the existence, uniqueness, and ergodic properties of SRB measures for C^2 Axiom A flows¹⁸. They proved that an invariant probability measure μ is an SRB measure if it possesses smooth conditional measures along local unstable manifolds, guaranteeing that time averages of continuous observables equal spatial averages for Lebesgue-almost all initial states in the topological basin¹⁸.

Ledrappier and Young (1985) established the relation between metric entropy $h_\mu(f)$, Lyapunov exponents $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_d$, and the pointwise dimension of invariant measures for C^2 diffeomorphisms²⁶. They proved that a measure μ is an SRB measure if and only if it satisfies the Pesin entropy formula $h_\mu(f) = \sum_{\lambda_i > 0} \lambda_i$, and that its Hausdorff dimension along unstable manifolds equals the sum of partial dimensions²⁷.

Kaplan and Yorke (1979) conjectured that the Hausdorff dimension of a chaotic attractor is

bounded by and typically equal to the Lyapunov dimension $D_{KY} = j + \frac{\sum_{i=1}^j \lambda_i}{|\lambda_{j+1}|}$, where j

is the largest integer such that $\sum_{i=1}^j \lambda_i \geq 0$ ²³.

Lu and Young (1999) extended SRB measure theory to infinite-dimensional state spaces (e.g., evolutionary PDEs in Hilbert spaces)²⁶. They proved the existence of SRB measures for infinite-dimensional dynamical systems possessing finite positive Lyapunov exponents and finite-dimensional unstable manifolds, provided the linear evolution satisfies a squeezing property²⁶.

Structural Applicability and Required Extensions

The Ledrappier-Young dimension formula and Lu-Young infinite-dimensional framework apply

directly if the system is smooth ($C^{1+\alpha}$) and possesses a spectral gap with finite positive Lyapunov exponents²⁶. However, if the configuration space incorporates non-Archimedean variables or singular continuous distributions, the smooth differential topology on tangent spaces breaks down³⁰. The Ledrappier-Young theory must then be generalized from smooth Riemannian manifolds to non-Archimedean metric measure spaces.

Assessment of Critical Open Gap

The primary literature gap in Area 3 is the calculation of Lyapunov exponents and verification of the Kaplan-Yorke dimension formula for non-uniformly hyperbolic, singular-hyperbolic, or non-smooth systems. While verified for Axiom A systems and specific 3D flows, general theoretical frameworks connecting positive Lyapunov exponents to exact fractal dimension bounds in high-dimensional open flows are lacking²³.

4. Ergodic Theory and Dynamics on p -Adic and Adelic Spaces

When physical space or environment-bath states are modeled over number-theoretic fields, dynamical systems must be constructed on p -adic fields $\mathbb{Q}_p$ or the ring of adeles $\mathbb{A}_{\mathbb{Q}}$.

Literature Map and Summaries

Mukhamedov and Khrennikov (2012) analyzed the local phase space dynamics of rational functions over the field of p -adic complex numbers $\mathbb{C}_p$ ³¹. They classified attracting, repelling, and indifferent fixed points, mapped Siegel disk structures, and proved that non-Archimedean rational dynamics are non-ergodic on unit spheres with respect to the Haar measure due to ultrametric isometric rigidity³¹.

Benedetto (2010) developed Fatou and Julia set theory in non-Archimedean dynamics,

establishing wandering domain theorems and p -adic potential theory over Berkovich spaces, resolving top-level structural distinctions between Archimedean and ultrametric Julia sets.

Schmidt (1995) established ergodic theory for algebraic endomorphisms on compact abelian groups, demonstrating that solenoids (such as $\Lambda = \bigcap_{n=0}^{\infty} f^n(T^3)$) act as compact adelic attractors with uniform mixing and Bernoullicity under diagonal endomorphisms³².

Einsiedler, Katok, and Lindenstrauss (2006) analyzed measure rigidity for higher-rank diagonal actions on arithmetic homogeneous spaces over the adeles $\mathbb{A}_{\mathbb{Q}}$ ³³. They proved that invariant probability measures under multi-parameter algebraic actions must be algebraic, exhibiting strong rigidity properties across places³³.

Structural Applicability and Required Extensions

Schmidt's solenoid formulations and Einsiedler-Katok-Lindenstrauss homogenous dynamics apply directly to algebraic actions on arithmetic quotients³². However, for non-linear dissipative

flows defined on continuous product spaces $\mathbb{A}_{\mathbb{Q}}^d = (\mathbb{R} \times \prod_p \mathbb{Q}_p)^d$, standard p -adic

dynamics fails to generate volume contraction because local p -adic maps are

non-Archimedean isometries ($|x + y|_p \leq \max(|x|_p, |y|_p)$)³¹. Standard p -adic dynamics lacks phase-space contraction in the classical sense³¹. Constructing non-linear dissipative flows on adelic spaces requires a non-trivial extension combining Archimedean volume contraction with non-Archimedean symbolic/tree-like dynamics³⁰.

Assessment of Critical Open Gap

Area 4 contains the most significant gap in the literature. There is virtually no research establishing Sinai-Ruelle-Bowen (SRB) physical measures, transfer operators, or hyperbolic attractors specifically on restricted product adelic spaces $\mathbb{A}_{\mathbb{Q}}$. While algebraic homogeneous adelic dynamics exists³³ and isolated p -adic discrete iterations are well-mapped³¹, non-linear dissipative flows on adelic manifolds remain unexplored.

5. Factorization and Correlation in Coupled Dynamical Systems

In multi-component or environment-coupled systems, establishing whether an invariant measure factorizes $\rho(\lambda|\text{settings}) \approx \rho(\lambda)$ determines whether subsystem variables maintain statistical independence.

Literature Map and Summaries

Bricmont and Kupiainen (1995) developed cluster expansion techniques for transfer operators of weakly Coupled Map Lattices (CML). They proved that for weakly coupled hyperbolic maps, the coupled system possesses a unique SRB measure μ , and space-time correlations decay exponentially.

Keller and Liverani (2006) proved the uniqueness and spectral gap properties of SRB measures for CMLs in arbitrary spatial dimensions under weak coupling $\epsilon > 0$, establishing statistical stability under small inter-component perturbations.

Cowieson and Young (2005) extended SRB measure theory to high- and infinite-dimensional coupled map lattices with local interaction kernels, establishing absolute continuity of conditional measures on unstable manifolds.

Viana (1997) surveyed dynamics in CMLs, proving that fully factorized independent product

states $\mu = \mu_1 \otimes \mu_2$ are generically non-invariant under non-zero coupling ($\epsilon > 0$).

Inter-component coupling distorts the joint invariant measure, generating spatial correlations

$$\text{Cov}_{\mu}(\phi_1, \phi_2) = O(\epsilon)$$

Structural Applicability and Required Extensions

These results demonstrate that an absolute independence identity

$\rho(\lambda_1, \lambda_2) = \rho(\lambda_1)\rho(\lambda_2)$ is generically false on coupled chaotic attractors. The coupling distorts the joint SRB measure away from a product measure. However, for weak coupling $\epsilon \ll 1$ and local interaction kernels, the literature proves approximate factorizability:

$$\|\mu_{\text{coupled}} - \mu_1 \otimes \mu_2\|_{\text{TV}} \leq C\epsilon$$

To apply this to open quantum or Bohmian settings, these discrete-time CML spectral bounds must be translated to continuous-time flows driven by nonlocal influence kernels.

Assessment of Critical Open Gap

The open gap in Area 5 lies in quantifying correlation bounds for strongly non-linear continuous-time coupled flows. Existing proofs rely on small coupling expansions ($\epsilon \ll 1$) or discrete-time expanding map lattices. Rigorous upper bounds on conditional correlation deviations for non-weakly coupled continuous dissipative flows do not currently exist.

6. Geometric Measure Theory and Transport on Singular/Fractal Sets

Because chaotic attractors are supported on sets of zero Lebesgue volume with fractal geometry, smooth differential equations and continuity equations must be generalized using geometric measure theory.

Literature Map and Summaries

Federer and Fleming (1960) developed the theory of normal currents, extending differential forms and integration over smooth manifolds to measure-theoretic chains¹⁸. They defined

spaces of currents $T \in \mathcal{N}_k(M)$ equipped with norm structures that permit the generalized

Stokes' theorem $\partial T(\omega) = T(d\omega)$ on nonsmooth, rectifiable sets¹⁸.

Harrison (2004) established chainlet geometry, providing a generalized calculus that integrates non-smooth vector fields across fractal boundaries. Harrison proved a generalized continuity equation and Stokes' theorem for domains with fractal boundaries whose Hausdorff dimension

exceeds $d - 1$, extending fluid-flow conservation laws to singular currents.

Kigami (2001) formulated analysis on post-critically finite (PCF) self-similar fractals by defining weak Laplacians Δ_μ as limits of discrete graph Laplacians, enabling Sobolev space theory (H^1) on non-manifold spaces.

Strichartz (2006) systematized differential equations on fractals, proving existence, uniqueness, and spectral properties for wave, heat, and advection operators defined directly on fractal

measure supports.

Structural Applicability and Required Extensions

Harrison's chainlet geometry and Federer-Fleming currents apply directly to continuity equations $\partial_t \rho + \nabla \cdot (\rho v) = 0$ defined on fractal attractors supported in $\mathbb{R}^n$ ¹⁸. They allow velocity fields to transport singular probability measures while rigorously conserving total mass. However, if the underlying space combines smooth manifolds with non-Archimedean spaces ($\mathbb{R}^n \times \mathbb{Q}_p^m$), standard real current theory breaks down³⁰. It must be synthesized with p -adic integration and Berkovich analytic spaces.

Assessment of Critical Open Gap

The key gap in Area 6 is the formulation of advection-dispersion operator calculus on multi-place product spaces (Archimedean $\times$ non-Archimedean). Existing fractal analysis operates on Euclidean fractals or abstract metric measure spaces, but lacks differential operator tools tailored to adelic spaces.

Comparative Assessment of Literature Across Domains

The following matrix summarizes the foundational theorems, structural applicability, relative maturity, and primary literature gaps across the six research domains.

Research Area	Foundational Mathematical Results	Direct Applicability to Open System Attractors	Relative Literature Maturity	Primary Open Literature Gap
1. Open System Dissipation	Feynman-Vern on Influence Functional ¹ , Davies Weak-Coupling Limit ⁸ , Jakšić-Pillet Spectral Deformation ¹⁴	Partial: Operator semigroups need mapping to deterministic phase-space vector fields ⁶ .	High (Operator level) / Moderate (Trajectory level)	Absence of rigorous proof deriving deterministic phase-space volume contraction ($\text{div } v < 0$) from open quantum density

				matrices without stochastic noise ⁶ .
2. Hyperbolicity of Dissipative Flows	Morales-Pacifico-Pujals Singular Hyperbolicity ¹⁸ , Tucker's Lorenz Attractor Proof ²⁰	High: Direct model for low-dimensional chaotic attractors with saddle points ¹⁸ .	High (3D flows) / Low (High-D flows)	Scarcity of non-computer-assisted analytical proofs of singular/sectional hyperbolicity for continuous-time non-linear flows ¹⁹ .
3. SRB Measure Existence & Computation	Bowen-Ruelle SRB Existence ¹⁸ , Ledrappier-Young Dimension Formula ²⁶ , Kaplan-Yorke Conjecture ²³	High: Provides explicit formulas for physical measures and fractal dimensions ²⁶ .	High (Smooth systems) / Moderate (Infinite-D)	Lack of theoretical frameworks verifying Kaplan-Yorke dimension equality for non-smooth, non-uniformly hyperbolic high-dimensional flows ²³ .
4. Adelic & p-Adic Dynamics	Mukhamedov-Khrennikov Rational Dynamics ³¹ , Schmidt Algebraic Solenoids ³² , Einsiedler-Katok-Lindenstrauss Rigidity ³³	Low: Local p -adic flows are isometric; lack volume contraction ³¹ .	Very Low (Adelic SRB) / High (Homogeneous)	CRITICAL GAP: Absence of Sinai-Ruelle-Bowen (SRB) measure theory and hyperbolic volume contraction on restricted

				product adeles $\mathbb{A}_{\mathbb{Q}}^{31}$
5. Factorization in Coupled Systems	Bricmont-Kupia inen Cluster Expansion, Keller-Liverani CML Uniqueness, Viana Coupling Theorems	Moderate: Factorization holds up to $O(\epsilon)$ weak coupling errors; false for strong coupling.	High (Discrete maps) / Moderate (Continuous flows)	Absence of rigorous upper bounds on conditional correlation deviations for continuous-tim e non-weakly coupled dissipative flows.
6. GMT on Singular/Fractal Sets	Federer-Flemin g Normal Currents ¹⁸ , Harrison Chainlet Geometry, Kigami Analysis on Fractals	High: Generalizes continuity equations and Stokes' theorem to fractal sets.	High (Euclidean fractals) / Low (Adelic fractals)	Lack of advection-disp ersion operator calculus formulated on multi-place product spaces (Archimedean $\times$ non-Archimede an).

Synthesis and Inter-Domain Dependencies

Establishing that an environment-traced open dynamical system admits a unique compact global attractor with an SRB measure requires bridging two major theoretical gaps across the literature pipeline:

- The Mapping Gap (Area 1 $\rightarrow$ Area 2):** Tracing out bath degrees of freedom via Feynman-Vernon influence functionals yields non-unitary reduced state dynamics¹. However, proving that the resulting Bohmian trajectory field produces a deterministic vector field with uniformly negative phase-space divergence ($\text{div } v < 0$) without inserting artificial Brownian noise remains an open challenge⁶.
- The Adelic SRB Gap (Area 4):** This represents the most significant deficit in the literature. While ergodic theory on arithmetic homogeneous spaces

(Einsiedler-Katok-Lindenstrauss)³³ and local rational P -adic dynamics (Mukhamedov-Khrennikov)³¹ are well-developed, there is virtually no formal mathematical framework for Sinai-Ruelle-Bowen measures, physical attractors, or volume-contracting

hyperbolic flows on restricted product adelic spaces $\mathbb{A}_{\mathbb{Q}}$.

Resolving these bottlenecks requires synthesizing operator-algebraic open system derivations with higher-dimensional sectional hyperbolicity, geometric measure theory, and non-Archimedean analysis.

Works cited

1. Caldeira-Leggett model - Scholarpedia,
http://www.scholarpedia.org/article/Caldeira-Leggett_model
2. A Rigorous Approach to the Feynman-Vernon Influence Functional,
https://www.researchgate.net/publication/30530888_A_Rigorous_Approach_to_the_Feynman-Vernon_Influence_Functional_and_its_Applications_I
3. Feynman-Vernon Influence Functional Theories,
https://www1.itp.tu-berlin.de/brandes/public_html/publications/notes/node67.html
4. Feynman-Vernon influence functional approach to quantum,
<https://epub.uni-regensburg.de/52240/1/PhysRevB.105.125417.pdf>
5. Open quantum systems with particle and bath driven by time ... - arXiv,
<https://arxiv.org/pdf/2505.05348>
6. (PDF) Dissipation in open quantum systems - ResearchGate,
https://www.researchgate.net/publication/312521113_Dissipation_in_open_quantum_systems
7. arXiv:2206.08636v2 [quant-ph] 13 Oct 2022, <https://arxiv.org/pdf/2206.08636>
8. THE WEAK COUPLING LIMIT AS A QUANTUM FUNCTIONAL,
https://art.torvergata.it/bitstream/2108/74968/1/AcFriLu90_The%20weak%20coupling%20limit.pdf
9. Weak-Coupling Limit. I A Contraction Semigroup for Infinite ... - arXiv,
<https://arxiv.org/html/0905.1012v1>
10. On the Derivation of the GKLS Equation for Weakly Coupled Systems,
https://www.researchgate.net/publication/319928829_On_the_Derivation_of_the_GKLS_Equation_for_Weakly_Coupled_Systems
11. CHARACTERIZATION AND SIMULATION OF MULTIPARTITE,
<https://helda.helsinki.fi/bitstreams/e7ad1ba4-ccc2-461a-bc2e-2504c36c28a7/download>
12. arXiv:1601.00509v1 [math-ph] 4 Jan 2016, <https://arxiv.org/pdf/1601.00509>
13. [1701.05024] Dissipation in the Caldeira-Leggett model - arXiv,
<https://arxiv.org/abs/1701.05024>
14. (PDF) Spectral Theory of Thermal Relaxation - ResearchGate,
https://www.researchgate.net/publication/258068743_Spectral_Theory_of_Thermal_Relaxation
15. Reviews in Mathematical Physics - World Scientific Publishing,

- <https://www.worldscientific.com/doi/10.1142/S0129055X24300097>
16. Scaled Caldeira-Leggett Dynamics: Deterministic Trajectories and, <https://arxiv.org/abs/2608.02054>
 17. Quantum thermodynamics of the Caldeira-Leggett model with non, <https://arxiv.org/abs/2405.00215>
 18. arXiv:math/0511352v3 [math.DS] 22 Mar 2007, <https://arxiv.org/pdf/math/0511352>
 19. arXiv:2204.10084v1 [math.DS] 21 Apr 2022, <https://arxiv.org/pdf/2204.10084>
 20. Statistical stability of geometric Lorenz attractors - ResearchGate, https://www.researchgate.net/publication/231337866_Statistical_stability_of_geometric_Lorenz_attractors
 21. University of Groningen Extreme value laws in dynamical systems, <https://pure.rug.nl/ws/files/2424964/2012PhysDHolland.pdf>
 22. Extreme value laws in dynamical systems under physical ... - SciSpace, <https://scispace.com/pdf/extreme-value-laws-in-dynamical-systems-under-physical-4oz8wawlfs.pdf>
 23. Robust Chaos and Its Applications 9814374075, 9789814374071, <https://dokumen.pub/robust-chaos-and-its-applications-9814374075-9789814374071.html>
 24. CSG Rds Final | PDF | Stochastic Process | Attractor - Scribd, <https://www.scribd.com/document/1056985683/Csg-Rds-Final>
 25. arXiv:2006.12157v2 [math.DS] 3 Mar 2021, <https://arxiv.org/pdf/2006.12157>
 26. SRB Measures and the Ledrappier-Young Dimension Formula, <https://www.authorea.com/doi/pdf/10.22541/au.174664030.00126829>
 27. Predicting Neural Scaling Laws without Training: A Data Manifold, <https://openreview.net/attachment?id=j2nPdWdRLs&name=pdf>
 28. Multifractal Analysis, Livšic Rigidity, and Fluctuation Theorems for, <https://arxiv.org/pdf/2604.18931>
 29. The Pesin Formula and the Gallavotti-Cohen Symmetry - arXiv, <https://arxiv.org/html/2604.18931v2>
 30. Primacohedron: A p-Adic String & Random-Matrix Framework for, https://www.preprints.org/frontend/manuscript/b9c473e9eb7a9e04cd885802ba0bd4d8/download_pub
 31. arXiv:1111.6725v1 [math.DS] 29 Nov 2011, <https://arxiv.org/pdf/1111.6725>
 32. ra.rings and algebras - How to understand a solenoid? - MathOverflow, <https://mathoverflow.net/questions/141049/how-to-understand-a-solenoid>
 33. arXiv:2101.05797v4 [math.DS] 10 Nov 2022, <https://arxiv.org/pdf/2101.05797>
 34. open problems in dynamics and related fields, https://www.imath.kiev.ua/~skolyada/Gorodnik_OP.pdf