component
fold


zoom: densest subfamily
zoom: most studied subfamily
zoom: IsPETase subfamily

Kubu-P
Mipa-P
TfCut_2
Cut190
LCC
ISPETase
HiC
PhaZ1
PueA
NylC
NylA
NylB
PLAase I


c00
c03
c04
c09
c11
c07
c08
c01
c02
c05
c06

Alpha/beta hydrolase (309 seq, 466 rec)
Ntn amidase (NylC) (45 seq, 53 rec)
Amidase signature (19 seq, 35 rec)
Oxygenase MpaB (Lcp) (10 seq, 15 rec)
Beta-lactamase / DD-peptidase (5 seq, 6 rec)
Trypsin-like protease (5 seq, 6 rec)
Subtilisin S8/S53 (5 seq, 12 rec)
Cytochrome c (RoxA) (3 seq, 5 rec)
Cupredoxin (laccase) (2 seq, 2 rec)
SGNH hydrolase (2 seq, 2 rec)
Peroxidase (MnP) (1 seq, 1 rec)
Cytochrome P450 (1 seq, 1 rec)
Acid protease (1 seq, 1 rec)
Glycoside hydrolase/deacetylase (1 seq, 2 rec)
Purple acid phosphatase (1 seq, 1 rec)
Beta-lactamase, unconfirmed (1 seq, 1 rec)

share of the 27,272 alignable pairs, per bin (%)




## Pairwise amino acid identity (%)




not alignable: the local alignment covers less than half of the shorter sequence

## How a cell gets its colour

**1 one pair**
row sequence i, column sequence j

**2 align**
Smith-Waterman, BLOSUM62, gap 11/1 (BLAST defaults)

**3 coverage**
aligned columns / length of the shorter

**4 identity**
identical columns / aligned columns

under 50%: grey, no identity

50% or more: binned identity, scale above

411 centroids, 84,255 pairs; 32% alignable, 68% not.
Outlined triangles: one fold each, in the fold colour used by the tree.
Rows run in the order of the circular tree.