

Analysis: Burger Optimization

Burger Optimization: Analysis

Small dataset

This dataset can be solved using complete search. Since $K \leq 8$, there are only at most $8! = 40320$ different orderings of ingredients.

Therefore, we can try all $K!$ different orderings of ingredients and find which ordering of ingredients has the minimum sum of squared differences between each ingredient's optimal and actual distance-to-bun values. Since we can calculate the sum of the squared differences between each ingredient's optimal and actual distance-to-bun values for one particular ordering of ingredients in $O(K)$ time, this solution runs in $O(K! \times K)$ time.

Large dataset

We need an observation to solve this dataset. Suppose we have two ingredients A and B to be placed at two positions X and Y. Let ingredients A and B have optimal distance-to-bun values of D_A and D_B , respectively, and let positions X and Y have actual distance-to-bun values of D_X and D_Y , respectively. Suppose $D_A \leq D_B$ and $D_X \leq D_Y$. We claim that it is no less optimal to put ingredient A at X and to put ingredient B at Y.

Why? Let us assume that $D_B = D_A + K_1$ and $D_Y = D_X + K_2$ for some non-negative integers K_1 and K_2 . Therefore, the cost of putting ingredient A at X and putting ingredient B at Y is $(D_A - D_X)^2 + (D_B - D_Y)^2$, and the cost of putting ingredient A at Y and putting ingredient B at X is $(D_A - D_Y)^2 + (D_B - D_X)^2$.

We can derive that

$$\begin{aligned} & [(D_A - D_X)^2] + [(D_B - D_Y)^2] \\ &= [D_A^2 - 2D_AD_X + D_X^2] + [D_B^2 - 2D_BD_Y + D_Y^2] \\ &= [D_A^2 - 2D_AD_X + D_X^2] + [D_B^2 - 2(D_A + K_1)(D_X + K_2) + D_Y^2] \\ &= [D_A^2 - 2D_AD_X + D_X^2] + [D_B^2 - 2D_AD_X - 2D_AK_2 - 2K_1D_X - 2K_1K_2 + D_Y^2] \\ &= [D_A^2 - 2D_AD_X - 2D_AK_2 + D_Y^2] + [D_B^2 - 2D_AD_X - 2K_1D_X + D_X^2] - [2K_1K_2] \\ &= [D_A^2 - 2D_A(D_X + K_2) + D_Y^2] + [D_B^2 - 2(D_A + K_1)D_X + D_X^2] - [2K_1K_2] \\ &= [D_A^2 - 2D_AD_Y + D_Y^2] + [D_B^2 - 2D_BD_X + D_X^2] - [2K_1K_2] \\ &= [(D_A - D_Y)^2] + [(D_B - D_X)^2] - [2K_1K_2] \end{aligned}$$

Therefore, $(D_A - D_X)^2 + (D_B - D_Y)^2 \leq (D_A - D_Y)^2 + (D_B - D_X)^2$, since $2K_1K_2$ is a non-negative integer. Therefore, we have proved that it is no less optimal to put ingredient A at X and to put ingredient B at Y. In other words, we want to put an ingredient which has a smaller optimal distance-to-bun value in a position which has a smaller actual distance-to-bun value.

Based on the above observation, the solution to this problem becomes simple. Let D' be the list of actual distance-to-bun values for the ingredients sorted in non-decreasing order (e.g. if K is 7, then $D' = \{0, 0, 1, 1, 2, 2, 3\}$), and let D be the list of optimal distance-to-bun values for the

ingredients sorted in non-decreasing order. Using the observation above, it is optimal to put the i -th ingredient on the list of D to a position which has an actual distance-to-bun value of $D'[i]$. Therefore, the answer is simply $\sum (D'[i] - D[i])^2$.

We can sort the ingredients in $O(K \log(K))$ time and calculate the sum of the squared differences between each ingredient's optimal and actual distance-to-bun values in $O(K)$ time. Therefore, this solution runs in $O(K \log(K))$ time.

(For a similar approach, you might enjoy reading about the [rearrangement inequality](#)).