



[SYLLABUS](#)
[Previous: 5.3.1 The Vasicek model](#)
[Up: 5.3 Methods for bonds](#)
[Next: 5.4 Computer quiz](#)

5.3.2 Extensions for derivatives ♠

The material in this section is intended for students at a more advanced level than your profile

Looking at a swap as being a bond paying a coupon $(r - X)dt$ over an infinitesimally short time interval, the differential change in the value of a bond (3.5#eq.6) can simply be supplemented with the corresponding increment. This immediately leads to the partial differential equation for a swap

$$\frac{\partial P}{\partial t} + \frac{1}{2}\sigma_s^2 \frac{\partial^2 P}{\partial r^2} + (\mu_s - \lambda\sigma_s) \frac{\partial P}{\partial r} = rP - (r - X) \quad (5.3.2\#eq.1)$$

and can again be solved using the finite element method. The new term does not involve any unknown and therefore appears on the right hand side of the linear problem (5.3.1#eq.4) with a contribution that can be integrated analytically

$$\int_{r-}^{r+} dr e_i(r) \Delta t (r - X) = \Delta t h (ih - X) \quad (5.3.2\#eq.2)$$

It has been coded into [FEMSolution.java](#) with an increment of the right hand side vector

```

c=b.dot(f);
for (int i=1; i<n; i++)
    c[i]+= timeStep*h*(i*h-X);
c0=(f[0]+(x[0]-X)*timeStep)*Math.exp(timeStep*X); // Boundary conditions
a.setL(0, 0.); a.setD(0, 1.); a.setR(0, 0.); c[0]=c0; //left: Dirichlet
dPdyn=(f[n]-f[n-1])/h; cn=c[n-1]-2*dx[0]*a.getL(n-1)*dPdyn;
a.setL(n,aln); a.setD(n,ann); a.setR(n, 0.); c[n]=cn; //right: Neuman
    
```

The Dirichlet boundary condition has been modified to account for the compounded interest from the fixed swap rate $-X$, but the rest remains the same as for the pricing of a bond.

Similar considerations are valid for [caplets](#) and [floorlets](#): viewed as a bond paying a coupon $\max(r - X, 0)dt$ for the caplet and $\max(X - r, 0)dt$ for the floorlet, this yields the modified Vasicek equation for a caplet

$$\frac{\partial P}{\partial t} + \frac{1}{2}\sigma_s^2 \frac{\partial^2 P}{\partial r^2} + (\mu_s - \lambda\sigma_s) \frac{\partial P}{\partial r} = rP - \max(r - X, 0) \quad (5.3.2\#eq.3)$$

and, by replacing $\max(r - X, 0)$ with $\max(X - r, 0)$, the counterpart for the floorlet. Both have been implemented into [FEMSolution.java](#) and the scheme for the caplet reads


```

c=b.dot(f);
for (int i=1; i<n; i++)
    if (i*h<X) c[i]+=0;
    else      c[i]+=timeStep*h*(i*h-X);
a.setL(0, 0.);a.setD(0, 1.);a.setR(0, 0.);c[0]=0.;//left: Dirichlet
dPdyn=0; cn=c[n-1]-2*dx[0]*a.getL(n-1)*dPdyn;
a.setL(n,aln);a.setD(n,ann);a.setR(n, 0.);c[n]=cn;//right:Neuman
// Construct the problem
// A*fp=B*f=c as before
```

Having calculated the fair price for a bond, a swap, cap or floor, it is relatively easy to calculate the value of derivatives such as bond options, swaptions, captions and floortions: their value depends on the same random variable and therefore satisfies the same equation as the underlying. For example, after calculating the value of the bond by solving the Vasicek equation (3.5#eq.6) backwards in time $T_B \rightarrow T$, the terminal bond option payoff (2.2.4#eq.1) can be used to integrate backwards further $T \rightarrow t$ until the present value of the bond option is found.

SYLLABUS Previous: 5.3.1 The Vasicek model Up: 5.3 Methods for bonds Next: 5.4 Computer quiz

Bond Pricing Independent valuations on over 2.5 million fixed income bonds www.standardandpoors.com	Bond Calculations in .NET Software components for MS .NET to calculate yield,etc.US & Intl bonds www.FTLabs.com	4.20%AER from HiSAVE Guarantee a high savings rate with our 2 Year Fixed Rate Account www.ICICIukPromotions.com	Top Savings Bonds Compare the Top Savings Accounts Online to Find the Best Interest lovemoney.com/bond-accounts
--	--	---	---



Copyright © [Lifelong-learners](#) at 11:29:14, April 24th, 2009

Ads by Google