# Know Your Weapon

Dr. Espen Gaarder Haug

1

# · Bronzin 1988 1900

### Market Formula (Bachelier-Thorp)

$$c = Se^{(b-r)T}N(d_1) - Xe^{-rT}N(d_2)$$
$$p = Xe^{-rT}N(-d_2) - Se^{(b-r)T}N(-d_1)$$

Where:

$$d_1 = \frac{\ln(S/X) + (b + \sigma_{X,T}^2/2)T}{\sigma_{X,T}\sqrt{T}}$$

$$d_1 = d_1 - \sigma_{X,T}\sqrt{T}$$

1104 Nelson

S = Asset price

X = Strike

T =Years to maturity

r = risk - free - rate

 $b = \cos t - \operatorname{of} - \operatorname{carry}$ 

 $\sigma_{xx}$  = volatility that can be different for each strike and maturity

See Haug 2007 "Derivatives Models on Models" chapter 2

2

### Black-Scholes-Merton

$$c = Se^{(b-r)T}N(d_1) - Xe^{-rT}N(d_2)$$
$$p = Xe^{-rT}N(-d_2) - Se^{(b-r)T}N(-d_1)$$

Where:

$$d_1 = \frac{\ln(S/X) + (b + \sigma^2/2)T}{\sigma\sqrt{T}}$$
$$d_2 = d_1 - \sigma\sqrt{T}$$

S =Asset price

X = Strike

T =Years to maturity

r = risk - free - rate

 $b = \cos t - \operatorname{of} - \operatorname{carry}$ 

 $\sigma$  = volatility

3


# Delta Greeks

- •Delta
- •Delta mirror strikes
- •Strike from delta
- •Elasticity

Delta higher than one

$$\Delta_{call} = \frac{\partial c}{\partial S} = e^{(b-r)T} N(d_1)$$

$$\Delta_{put} = \frac{\partial p}{\partial S} = -e^{(b-r)T} N(-d_1)$$



### Delta Mirror Strikes

 $X_{p} = \frac{S^{2}}{X_{C}} \left( e^{(2b+\sigma^{2})T}, \quad X_{C} = \frac{S^{2}}{X_{p}} e^{(2b+\sigma^{2})T} \right)$ 

Special case delta symmetric straddle (Wystrup(1999)):

$$X_C = X_P = Se^{(b+\sigma^2/2)T}$$

Delta symmetric asset:  $S = Xe^{(-b-\sigma^2/2)T}$ 

At this strike the delta is 
$$\Delta_C = \frac{e^{(b-r)T}}{2}$$
,  $\Delta_P = -\frac{e^{(b-r)T}}{2}$ 

$$c = \frac{Se^{(b-r)T}}{2} - X^{-rT}N(-\sigma\sqrt{T}), \quad p = X^{-rT}N(\sigma\sqrt{T}) - \frac{Se^{(b-r)T}}{2}$$

At this strike the delta is 
$$\Delta_C = \frac{e^{(b-r)T}}{2}$$
,  $\Delta_P = -\frac{e^{(b-r)T}}{2}$ 

$$C = \frac{Se^{(b-r)T}}{2} - X^{-rT}N(-\sigma\sqrt{T}), \quad p = X^{-rT}N(\sigma\sqrt{T}) - \frac{Se^{(b-r)T}}{2}$$

X-5

12/56/3 12,56/3 Strikes from delta

4=25% T=3/12

Wystrup(1999):

$$X_C = S \exp[N^{-1}(\Delta_C e^{(r-b)T})\sigma\sqrt{T} + (b+\sigma^2/2)T]$$

$$X_P = S \exp[N^{-1}(-\Delta_P e^{(r-b)T})\sigma\sqrt{T} + (b+\sigma^2/2)T]$$

Robust and accurate approximation of inverse cumulative normal distribution needed, Moro(1995).

(AW) FX +/

GRAB
At 09:16 Op 12.1000 H

Range 172/702 - Upper Chart: 2 Candle C



### DdeltaDvol

$$\frac{\partial c}{\partial S \partial \sigma} = \frac{\partial p}{\partial S \partial \sigma} = \frac{-e^{(b-r)T} d_2}{\sigma} n(d_1)$$

# m0

#### Maximal value at

$$S_L = Xe^{-bT - \sigma\sqrt{T}\sqrt{4 + T\sigma^2}/2}$$

Minimal value at


$$S_U = X e^{-bT + \sigma\sqrt{T}\sqrt{4 + T\sigma^2}/2}$$

Minimal value at

$$X_L = S e^{bT - \sigma \sqrt{T} \sqrt{4 + T\sigma^2}/2}$$

Maximal value at

$$X_U = Se^{bT + \sigma\sqrt{T}\sqrt{4 + T\sigma^2}/2}$$



# **Useful Tools**

- A library
- Paper and pencil
- Mathematica
- Maple
- Matlab (?)
- Others?

Implementation:

VBA, VB, C/C++, Java....

you name it

# Elasticity

$$\Lambda_{call} = \Delta_{call} \frac{S}{call}, \quad \Lambda_{put} = \Delta_{put} \frac{S}{put}$$

Option volatility:  $\sigma_0 \approx \sigma |\Lambda|$  Compound options

Option Beta, expected return satisfy the CAPM equation (Merton-71):

$$E[return] = r + E[r_m - r]\beta_i$$

$$\beta_C = \frac{S}{call} \Delta_C \beta_S = \Lambda_C \beta_S, \quad \beta_P = \frac{S}{put} \Delta_P \beta_S = \Lambda_P \beta_S$$

Option Sharp ratios


$$\frac{\mu_O - r}{\sigma_O} = \frac{\mu_S - r}{\sigma}$$

Smile?

Charm

$$\frac{\partial \Delta_C}{\partial T} = -e^{(b-r)T} \left[ n(d_1) \left( \frac{b}{\sigma \sqrt{T}} - \frac{d_2}{2T} \right) + (b-r)N(d_1) \right]$$

$$\frac{\partial \Delta_P}{\partial T} = -e^{(b-r)T} \left[ n(d_1) \left( \frac{b}{\sigma \sqrt{T}} - \frac{d_2}{2T} \right) - (b-r)N(-d_1) \right]$$



# Gamma Greeks

- •Gamma
- •Saddle gamma
- •GammaP
- •Gamma symmetry
- •DGammaDVol
- ${\color{red}\bullet} DGammaDspot$
- •DGammaDTime

13


#### Saddle Gamma

Alexander Adamchuk www.wilmott.com

$$T_{\Gamma} = \frac{1}{2(\sigma^2 + b)}$$

$$S_{\Gamma} = Xe^{(-b-3\sigma^2/2)T_S}$$

$$\Gamma_{S} = \frac{e^{(b-r)T} \sqrt{\frac{e}{\pi}} \sqrt{\frac{b}{\sigma^2} + 1}}{X}$$




#### GammaP

$$\Gamma_P = \Gamma \frac{S}{100}$$

Max GammaP at

$$S = Xe^{(-b-\sigma^2/2)T}$$

$$X = Se^{(b+\sigma^2/2)T}$$




15

#### Saddle GammaP

•Spot gamma

•Forward gamma



F= SobT

# Gamma-symmetry


Put-call symmetry Bates(1991) and Carr and Bowie (1994):

$$c(S, X, T, r, b, \sigma) = \frac{X}{Se^{bT}} p(S(Se^{bT})^2)^2, T, r, b, \sigma)$$

Gamma-symmetry

$$\Gamma(S, X, T, r, b, \sigma) = \frac{X}{Se^{bT}} \Gamma(S, \frac{(Se^{bT})^2}{X}, T, r, b, \sigma)$$

Also gives vega and cost-of-carry symmetry



17

DgammaDvol

$$\frac{\partial \Gamma}{\partial \sigma} = \Gamma \left( \frac{d_1 d_2 - 1}{\sigma} \right)$$


$$\frac{\partial \Gamma}{\partial \sigma} = \Gamma \left( \frac{d_1 d_2 - 1}{\sigma} \right) \qquad \frac{\partial \Gamma_P}{\partial \sigma} = \Gamma_P \left( \frac{d_1 d_2 - 1}{\sigma} \right)$$


Positive outside interval

$$S_L = Xe^{-bT - \sigma\sqrt{T}\sqrt{4 + T\sigma^2}/2}$$

$$S_U = Xe^{-bT + \sigma\sqrt{T}\sqrt{4 + T\sigma^2}/2}$$

0=30%,
0=40%,
0=20%






19

#### Speed (DgammaDspot)


$$\frac{\partial^{3} c}{\partial S^{3}} = -\frac{\Gamma\left(1 + \frac{d_{1}}{\sigma\sqrt{T}}\right)}{S}$$

$$SpeedP = -\Gamma \frac{d_1}{S}$$

Speed is used by Fouque, Papanicolaou, and Sircar (2000) as part of stochastic vol model



$$\begin{split} \frac{\partial \Gamma}{\partial T} &= \Gamma \bigg( r - b + \frac{b d_1}{\sigma \sqrt{T}} + \frac{1 - d_1 d_2}{2T} \bigg) \\ \frac{\partial \Gamma_P}{\partial T} &= \Gamma_P \bigg( r - b + \frac{b d_1}{\sigma \sqrt{T}} + \frac{1 - d_1 d_2}{2T} \bigg) \end{split}$$



21

### Numerical Greeks

5=100\$

- •More robust (?)
- •Model independent
- •Faster to implement (?)

Two-sided finite difference

$$\Delta_C \approx \frac{c(S + \Delta S, X, T, r, b, \sigma) - c(S - \Delta S, X, T, r, b, \sigma)}{2\Delta S}$$

Backward derivative.

$$\Theta \approx \frac{c(S,X,T,r,b,\sigma) - c(S,X,T - \Delta T,r,b,\sigma)}{\Delta T}$$



Numerical Greeks
$$\Delta_{C} \approx \frac{c(S + \Delta S, X, T, r, b, \sigma_{1}) - c(S - \Delta S, X, T, r, b, \sigma_{2})}{2\Delta S}$$

$$\Theta \approx \frac{c(S, X, T, r, b, \sigma_1) - c(S, X, T - \Delta T, r, b, \sigma_2)}{\Delta T}$$

Gamma and other second derivatives, central finite difference

$$\Gamma \approx \frac{c(S + \Delta S,...) - 2c(S,...) + c(S - \Delta S,...)}{\Delta S^2}$$

Speed and other third order derivatives, central finite difference

$$Speed \approx \frac{1}{\Delta S^3} [c(S + 2\Delta S,...) - 3c(S + \Delta S,...) + 3c(S,...) - c(S - \Delta S,...)]$$

23

# Know Your Weapon Part 2

### **Numerical Greeks**

What about mixed derivatives? For example DdeltaDvol and Charm

$$\begin{aligned} & D delta D vol \approx \frac{1}{4 \Delta S \Delta \sigma} [c(S + \Delta S, ..., \sigma + \Delta \sigma) - c(S + \Delta S, ..., \sigma - \Delta \sigma) \\ & - c(S - \Delta S, ..., \sigma + \Delta \sigma) + c(S - \Delta S, ..., \sigma - \Delta \sigma)] \end{aligned}$$

25

# Vega "Greeks"

- •Vega
- •Vega maximum
- •VegaP
- •Vega symmetry
- •Vega Leverage
- •DVegaDvol
- •DVegaDtime

Vega 
$$\frac{\partial c}{\partial \sigma} = 3$$

 $\frac{\partial c}{\partial \sigma} = Se^{(b-r)T} n(d_1) \sqrt{T} \quad P = C \approx 0.45 \cdot \sigma = 7$ 

Vega local max

$$S = Xe^{(-b+\sigma^2/2)T}$$
$$X = Se^{(b+\sigma^2/2)T}$$


#### Global maximum

$$T_{V} = \frac{1}{2r}$$

$$S_{V} = Xe^{(-b+\sigma^{2}/2)T_{V}}$$

$$= Xe^{\frac{-b+\sigma^{2}/2}{2r}}$$


$$Vega(S_{\overline{V}}, T_{\overline{V}}) = \frac{X}{2\sqrt{re\pi}}$$



27

Why the Yega top?

Discounting at some point will dominate over volations (Vega).



# Vega-symmetry

Put-call symmetry Bates(1991) and Carr and Bowie (1994):

$$c(S, X, T, r, b, \sigma) = \frac{X}{Se^{bT}} p(S, \frac{(Se^{bT})^2}{X}, T, r, b, \sigma)$$

Vega-symmetry

$$Vega(S, X, T, r, b, \sigma) = \frac{X}{Se^{bT}} Vega(S, \frac{(Se^{bT})^2}{X}, T, r, b, \sigma)$$

Also gives gamma and cost-of-carry symmetry

29

Vega-gamma relationship

Taleb(1997):

$$Vega = \Gamma \sigma S^2 T$$

Vega from delta

$$Vega = Se^{(b-r)T} \sqrt{T} n[N^{-1}(e^{(r-b)T} \mid \Delta \mid)]$$

Gamma from delta

$$\Gamma = \frac{e^{(b-r)T} n[N^{-1}(e^{(r-b)T} \mid \Delta \mid)]}{S\sigma\sqrt{T}}$$

# VegaP

Vega gives dollar change in option value for one percent point change in implied volatility. VegaP gives dollar change in option value for percentage move in volatility.

$$VegaP = \frac{\sigma}{10} Se^{(b-r)T} n(d_1) \sqrt{T}$$

VegaP makes much more sense when comparing sensitivity to changes in Implied volatility.


31

If you want to speculate on an increase in implied volatility what type of options offers the most bang for the bucks?

#### Vega leverage

Percent change in option value for percent point change in implied volatility.

$$Vega \frac{\sigma}{call}$$
,  $Vega \frac{\sigma}{put}$ 



33

#### DvegaDvol Vomma/Volga

$$\frac{\partial^2 c}{\partial \sigma^2} = Vega\left(\frac{d_1 d_2}{\sigma}\right)$$

#### Positive outside


$$S_L = Xe^{(-b-\sigma^2/2)T}$$

$$S_U = Xe^{(-b+\sigma^2/2)T}$$

#### Positive outside

$$X_L = Se^{(b-\sigma^2/2)T}$$


$$S_U = Se^{(b+\sigma^2/2)T}$$



VouV.L

#### DvegaDtime


$$\frac{\partial^{2} c}{\partial \sigma \partial T} = Vega \left( r - b + \frac{bd_{1}}{\sigma \sqrt{T}} - \frac{1 + d_{1}d_{2}}{2T} \right)$$



35

DvegaDtime

$$\frac{\partial^{2} c}{\partial \sigma \partial T} Vega \left( r - b + \frac{bd_{1}}{\sigma \sqrt{T}} - \frac{1 + d_{1}d_{2}}{2T} \right)$$



### Theta

$$\begin{split} \Theta_C &= -\frac{\partial c}{\partial T} = -\frac{Se^{(b-r)T}n(d_1)}{2\sqrt{T}} - (b-r)Se^{(b-r)T}N(d_1) - rXe^{-rT}N(d_2) \\ \Theta_C &= -\frac{\partial c}{\partial T} = -\frac{Se^{(b-r)T}n(d_1)\sigma}{2\sqrt{T}} + (b-r)Se^{(b-r)T}N(-d_1) + rXe^{-rT}N(-d_2) \end{split}$$

Drift-less theta

$$\theta_C = \theta_P = -\frac{Sn(d_1)}{2\sqrt{T}}$$

Theta symmetry

$$\theta(S, X, T, 0, 0, \sigma) = \frac{X}{S}\theta(S, \frac{S^2}{X}, T, 0, 0, \sigma)$$

Bleed-offset volatility

37

#### Rho

$$\rho_C = \frac{\partial c}{\partial r} = TXe^{-rT}N(d_2), \quad \rho_P = \frac{\partial p}{\partial r} = -TXe^{-rT}N(-d_2)$$

In case of options on futures (b=0)

$$\rho_C = \frac{\partial c}{\partial r} = -Tc, \quad \rho_P = \frac{\partial p}{\partial r} = -Tp$$

# Probability "Greeks"

> Black Scholes

Risk neutral probability of ending up in-the-money

$$\zeta_C = N(d_2) > 0, \quad \zeta_P = N(-d_2) > 0$$

Strike-delta

$$\frac{\partial c}{\partial X} = -e^{-rT}N(d_2), \quad \frac{\partial p}{\partial X} = e^{-rT}N(-d_2)$$

Probability mirror strikes

$$X_P = \frac{S^2}{X_C} \, e^{(2b - \sigma^2)T}, \quad X_C = \frac{S^2}{X_P} \, e^{(2b - \sigma^2)T}$$

Probability neutral straddle

$$X_C = X_P = Se^{(b-\sigma^2/2)T}$$

39

# Probability "Greeks"

Strikes from probability

$$X_C = S \exp[-N^{-1}(p_i)\sigma\sqrt{T} + (b - \sigma^2/2)T]$$

$$X_P = S \exp[N^{-1}(-p_i)\sigma\sqrt{T} + (b - \sigma^2/2)T]$$

Risk neutral probability density

$$RND = \frac{\partial^{2} c}{\partial X^{2}} = \frac{\partial^{2} p}{\partial X^{2}} = \frac{n(d_{2})e^{-rT}}{X\sigma\sqrt{T}}$$

Probability neutral straddle


$$X_C = X_P = Se^{(b-\sigma^2/2)T}$$

Risk neutral probability density

$$RND = \frac{\partial^2 c}{\partial X^2} = \frac{\partial^2 p}{\partial X^2} = \frac{n(d_2)e^{-rT}}{X\sigma\sqrt{T}}$$

Breeden and Litzenberger (1978)

1994-199404-1994-



esponhaug 6 mac. com

# Probability "Greeks"

Risk neutral probability of ever being in-the-money

$$p_C = (X/S)^{\mu+\lambda} N(-z) + (X/S)^{\mu-\lambda} N(-z + 2\lambda\sigma\sqrt{T})$$

$$p_p = (X/S)^{\mu+\lambda} N(z) + (X/S)^{\mu-\lambda} N(z - 2\lambda\sigma\sqrt{T})$$

where

$$z = \frac{\ln(X/S)}{\sigma\sqrt{T}} + \lambda\sigma\sqrt{T}, \quad \mu = \frac{b - \sigma^2/2}{\sigma^2}, \quad \lambda = \sqrt{\mu^2 + \frac{2r}{\sigma^2}}$$