

Consider a 1 + 1D massless Dirac fermion with Lagrangian

$$\mathcal{L} = i\bar{\psi}\gamma^\mu\partial_\mu\psi.$$

on a circle of perimeter L . Recall in the class that, we can choose the 2D Gamma matrices as $\gamma^\mu = (\sigma_2, i\sigma_1)$, therefore the 2D Dirac fermion is a two-component spinor. Further, for fermions, we need to impose anti-periodic boundary condition as

$$\psi(t, x + L) = -\psi(t, x).$$

Now answer the following questions:

- a. Show that the solution to the Dirac equation can be written as

$$\psi(t, x) = \frac{1}{\sqrt{L}} \sum_{n=0}^{+\infty} \begin{pmatrix} c_n e^{-i\omega_n(t-x)} \\ d_n^\dagger e^{i\omega_n(t+x)} \end{pmatrix},$$

and specify the value of ω_n . Further use canonical quantization to diagonalize the Hamilton \hat{H} in terms of c_n and d_n . Can you interpret the Dirac spinor ψ as a 2D electron-positron system? Explain your reason;

- b. The Lagrangian admits a $U(1)$ -symmetry respect to $\psi \rightarrow e^{i\alpha}\psi$, with α a constant phase factor. The associated $U(1)$ -charge \hat{N} is called fermionic number. Find the fermionic number \hat{N} in terms of c_n and d_n , and show that it can be interpreted as the difference between the numbers of positrons and electrons;
- c. Define $q = e^{-\omega/T}$ and $z = e^{\mu/T}$ for the energy and the fermionic number fugacities, then the grand canonical partition function is defined as

$$Z(q, z) = \text{Tr} \left(e^{-\hat{H}/T + \mu \hat{N}/T} \right) = \sum_n \left\langle n \left| e^{-\hat{H}/T + \mu \hat{N}/T} \right| n \right\rangle.$$

Show that the partition function is given by

$$Z(q, z) = \prod_{n=1}^{\infty} \left(1 + z q^{n-\frac{1}{2}} \right) \left(1 + z^{-1} q^{n-\frac{1}{2}} \right);$$

- d. From an old perspective of “Dirac sea” (The vacuum is the state in which all negative energy states are filled. Therefore an electron is a state created above the vacuum, while a positron is a “hole” state where all negative energy states are occupied except one), the partition function can be recast as

$$Z(q, z) = \sum_{\substack{\text{fermion} \\ \text{occupations}}} e^{-E/T + \mu N/T} = \sum_{N=-\infty}^{+\infty} z^N Z_N(q),$$

where $Z_N(q)$ is the partition function counting states for a given fermionic number N . Show that

$$Z_0(q) = \prod_{n=1}^{\infty} \frac{1}{1 - q^n},$$

and thus

$$Z_N(q) = q^{\frac{N^2}{2}} Z_0(q).$$

Now combining the result of (c), you can give a physical proof of the famous Jacobi triple identity

$$\prod_{n=1}^{\infty} (1 - q^n) \left(1 + z q^{n-\frac{1}{2}}\right) \left(1 + z^{-1} q^{n-\frac{1}{2}}\right) = \sum_{N=-\infty}^{+\infty} z^N q^{\frac{N^2}{2}}.$$