

# **A Study of Various Activating Mutations and Conditions of the Epithelial Sodium Channel**

Richard Posert

2023-07-21



Table of contents

Acknowledgements 1

I. Introduction 3

1. ENaC in the Body 5

1.1. ENaC in the kidney . . . . . 5

References 7



# Acknowledgements

We'll get here.



## **Part I.**

# **Introduction**





# 1. ENaC in the Body

## 1.1. ENaC in the kidney

Blood pressure must be maintained within a narrow window of acceptable values. Too low, and vital organs do not receive sufficient oxygen and nutrients to function, but blood vessels sustain damage when pressure is too high. It is not surprising, then, that the human body has evolved several mechanisms for responding to changes in blood pressure, each with their own timescale<sup>1</sup>. ENaC is the essential mechanism of the longest-term control, kidney excretion.

When the kidney is not receiving enough salt, it initiates a cascade which ends with the mineralocorticoid hormone aldosterone augmenting the activity of ENaC in the distal nephron. Apical sodium permeability is increased both by synthesis of new channels as well as trafficking of an existing pool of channels to the cell surface<sup>2,3</sup>. Despite its expression relatively late in the nephron, ENaC is the rate-limiting step of sodium reabsorption in the kidney. Sodium in the principal cells is transported across the basolateral membrane by the  $\text{Na}^+/\text{K}^+$  ATPase, and therefore induces retention of extra water to maintain the tightly-controlled plasma sodium level. Thus, ENaC controls three essential functions of kidney filtration: first, the amount of sodium reabsorbed by the kidney; second, blood volume (and therefore pressure); third, the amount of potassium passed from the plasma into the urine.

A variety of ENaC mutations have dramatic effects on patients' blood pressure. One of the earliest described is Liddle syndrome<sup>4</sup>. Liddle syndrome (also called pseudoaldosteronism) results from an autosomal dominant gain-of-function mutation in ENaC. Severe hypertension, hypokalemia, and low renin and aldosterone are hallmarks of the disease. It is a rare disorder, with only 72 families described as of 2018<sup>5</sup>. However, after excluding patients with other clear causes (primary aldosteronism, kidney or heart diseases, and obstructive sleep apnea), approximately one in one hundred hypertensive patients had Liddle syndrome, indicating that the prevalence may be

Of note: Asher and colleagues report an aldosterone mediated increase in expression of the  $\beta$  and  $\gamma$ , but not  $\alpha$ , subunits. A trend of subunit-specific regulation is widely observed in the literature, but poorly understood.

## 1. ENaC in the Body

higher than is currently thought<sup>6,7</sup>. All but one of the described cases involve mutation of the  $\beta$  or  $\gamma$  subunits<sup>5,7</sup>.

A majority of the mutations in the  $\beta$  and  $\gamma$  subunits disrupt or remove entirely a proline-rich PY motif at the C-terminus of those channels<sup>8</sup>. This PY motif is the binding site for the E3 ubiquitin ligase NEDD4-2 (see section ). Study of ENaC surface dwell time is complicated by a reserve pool maintained by the cells to be cycled up to the membrane, but there is a consensus that ENaC is recycled quickly; surface half life estimates range from fifteen minutes to three hours, with the low end having more support<sup>9</sup>. ENaC lacking the PY motif cannot be pulled back in from the membrane, thus increasing sodium permeability by increasing  $N$  rather than the conductivity or  $P_O$  of the channels<sup>10-12</sup>. The remaining minority of described Liddle syndrome mutations (including the sole ENaC $_{\alpha}$  mutation) which do not affect PY-motif binding instead directly augment channel  $P_O$ <sup>5</sup>.

It is worth noting here that the  $\alpha$  subunit does have a PY motif, but no Liddle syndrome mutation of the  $\alpha$  PY has been described. This is discussed further in

## References

1. Guyton, A. C. Blood Pressure Control---Special Role of the Kidneys and Body Fluids. *Science* **252**, 1813--1816 (1991).
2. Asher, C., Wald, H., Rossier, B. C. & Garty, H. Aldosterone-induced increase in the abundance of Na<sup>+</sup> channel subunits. *American Journal of Physiology-Cell Physiology* **271**, C605--C611 (1996).
3. Frindt, G. & Palmer, L. G. Surface expression of sodium channels and transporters in rat kidney: Effects of dietary sodium. *Am. J. Physiol. Renal Physiol.* **297**, F1249--F1255 (2009).
4. Liddle, G. W., Bledsoe, T. & Coppage, W. S. A familial renal disorder simulating primary aldosteronism but with negligible aldosterone secretion. *Trans. Assoc. Am. Physicians* **76**, 199--213 (1963).
5. Tetti, M. *et al.* Liddle Syndrome: Review of the Literature and Description of a New Case. *International Journal of Molecular Sciences* **19**, (2018).
6. Wang, L.-P *et al.* Prevalence of Liddle Syndrome Among Young Hypertension Patients of Undetermined Cause in a Chinese Population. *The Journal of Clinical Hypertension* **17**, 902--907 (2015).
7. Liu, K. *et al.* Analysis of the genes involved in Mendelian forms of low-renin hypertension in Chinese early-onset hypertensive patients. *Journal of Hypertension* **36**, (2018).
8. Yang, K.-Q., Xiao, Y., Tian, T., Gao, L.-G. & Zhou, X.-L. Molecular genetics of Liddle's syndrome. *Clinica Chimica Acta* **436**, 202--206 (2014).
9. Butterworth, M. B., Edinger, R. S., Frizzell, R. A. & Johnson, J. P. Regulation of the epithelial sodium channel by membrane trafficking. *American Journal of Physiology-Renal Physiology* **296**, F10--F24 (2009).
10. Schild, L. *et al.* A mutation in the epithelial sodium channel causing Liddle disease increases channel activity in the *Xenopus laevis* oocyte expression system. *Proceedings of the National Academy of Sciences* **92**, 5699--5703 (1995).

## *References*

11. Abriel, H. *et al.* Defective regulation of the epithelial Na<sup>+</sup> channel by Nedd4 in Liddle's syndrome. *J Clin Invest* **103**, 667--673 (1999).
12. Lu, C., Pribanic, S., Debonneville, A., Jiang, C. & Rotin, D. The PY Motif of ENaC, Mutated in Liddle Syndrome, Regulates Channel Internalization, Sorting and Mobilization from Subapical Pool. *Traffic* **8**, 1246--1264 (2007).