

①

$$\begin{aligned}
 & \log_2\left(\frac{8\sqrt{2}}{16}\right) + \log_2(32) - 2\log_2(4) = \\
 & = \log_2(8\sqrt{2}) - \log_2 16 + \log_2 2^5 - 2\log_2 2^2 = \\
 & = \log_2 2^3 + \log_2 2^{\frac{1}{2}} - \log_2 2^4 + \log_2 2^5 - 2\log_2 2^2 = \\
 & = 3 + \frac{1}{2} - 4 + 5 - 4 = -8 + 8 + \frac{1}{2} = \frac{1}{2}
 \end{aligned}$$

②

$$\begin{aligned}
 & \log_3(x-1) + \log_3(x+1) = 2 \\
 & \log_3(x-1)(x+1) = 2 \\
 & \log_3(x^2-1^2) = 2 \\
 & x^2-1^2 = 3^2 \Rightarrow x^2 = 3^2 + 1 = 10 \\
 & x = \pm \sqrt{10} \approx 3, 16
 \end{aligned}$$

$$x = 3,16 \quad \checkmark$$

$$x = -3,16 \quad \times$$

$$x - 1 > 0$$

$$x - 1 < 0$$

$$x + 1 > 0$$

$$x + 1 < 0$$

$$x = 3,16$$

③

$$10000 \text{ \$; } 6\% \text{ per year} = 72000 \text{ \$}$$

$$10000 \left(1 + \frac{0,06}{4}\right) - 1st \text{ Quarter}$$

$$10000 \left(1 + \frac{0,06}{4}\right)^2 - 2nd \text{ Quarter}$$

$$10000 \left(1 + \frac{0,06}{4}\right)^3 - 3rd \text{ Quarter}$$

$$10000 \left(1 + \frac{0,06}{4}\right)^4 - 4 \text{ Quarters} = 1 \text{ year}$$

$t$ -years

$$10000 \left(1 + \frac{0,06}{4}\right)^{4t} = 20000$$

$$\left(1 + \frac{0,06}{4}\right)^{4t} = \frac{20000}{10000} = 2$$

$$1,015^{4t} = 2$$

$$\ln(1,015)^{4t} = \ln 2$$

$$4t \cdot \ln(1,015) = \ln 2$$

$$4t = \frac{\ln 2}{\ln(1,015)} \Rightarrow t = \frac{\ln 2}{4 \ln 1,015}$$

$$t = \frac{0,6931}{4 \cdot 0,0149} = \textcircled{11,64}$$

④  $N(t) = N_0 e^{-kt}$

$N_0$  - initial amount;  $k$  - const,  $t$  - years  
 half-life  $\lambda$  - 5 years, find  $k$ ?

$$\frac{N_0}{2} = N_0 e^{-k \cdot 5}; \frac{1}{2} = e^{-k \cdot 5}$$

→ 1

$$\ln\left(\frac{1}{2}\right) = -k \cdot 5 \quad (\ln(e))$$

$$-k = \frac{\ln\left(\frac{1}{2}\right)}{5},$$

$$-k = -0,138629; \Rightarrow k = 0,138629$$

$$K \approx 0,14$$

⑤

$$100g \xrightarrow{3 \text{ hours}} 70g$$

$$100g \xrightarrow{?} 20g$$

$$M(t) = 100 \cdot e^{-k \cdot 3}$$

$$\frac{4}{10} = e^{-k \cdot 3} \Rightarrow \ln\left(\frac{4}{10}\right) = -3k$$

$$k = \frac{\ln\left(\frac{4}{10}\right)}{-3} = 0,118891648$$

$$20 = 100 \cdot e^{-kt}$$

$$\ln\left(\frac{2}{10}\right) = -k \cdot t ; \Rightarrow t = \frac{\ln\left(\frac{2}{10}\right)}{-k}$$

?

$$t = \frac{\ln\left(\frac{2}{10}\right)}{-0,1189} = \boxed{13,54 \text{ hours}}$$

⑥ Find vector from  $A(1,2,3)$  to  
 $B(4,6,9)$

$$\vec{v}(AB) = (4-1, 6-2, 9-3) = (3, 4, 6)$$

$$|\vec{v}| = \sqrt{3^2 + 4^2 + 6^2} = \boxed{7,81}$$

⑦  $v = 4\hat{i} - 2\hat{j} + 4\hat{k}$  in matrix and  
 find magnitude

$$\vec{v} = \begin{bmatrix} 4 \\ -2 \\ 4 \end{bmatrix}$$

$$\vec{v} = \sqrt{4^2 - 2^2 + 4^2} = \boxed{\sqrt{30}}$$

⑧  $\vec{a} = (2, -1, 3)$ ;  $\vec{b} = (-1, 4, 2)$

Compute  $3\vec{a} - 2\vec{b}$

$$3\vec{a} = (6, -3, 9); \quad 2\vec{b} = (-2, 8, 4)$$

$$3\vec{a} - 2\vec{b} = (6\hat{i} - 3\hat{j} + 9\hat{k}) - (-2\hat{i} + 8\hat{j} + 4\hat{k})$$

$$= 8\hat{i} - 11\hat{j} + 5\hat{k} = \boxed{(8, -11, 5)}$$

⑨ Find the angle between  $\vec{P}$  and  $\vec{Q}$ ;  $\vec{P} = (1, 2, 3)$   $\vec{Q} = (4, -5, 6)$

$$\vec{P} \cdot \vec{Q} = \|\vec{P}\| \cdot \|\vec{Q}\| \cdot \cos \theta$$

$$\vec{P} \cdot \vec{Q} = 1 \cdot 4 + 2 \cdot (-5) + 3 \cdot 6 = 12$$

$$\sqrt{1^2 + 2^2 + 3^2} \cdot \sqrt{4^2 - 5^2 + 6^2} \cdot \cos \theta = 12$$

$$\frac{3\sqrt{14}}{\sqrt{32}} \cdot \frac{8\sqrt{74}}{\sqrt{32}} \cdot \cos \theta = 12$$

✓  
32,8329

$$\cos \theta = \frac{12}{32,8329} = 0,365487$$

$$\cos \theta = 0,365487 \Rightarrow \theta \approx 69^\circ$$

⑩  $\vec{u} = (2, -1, 4)$  and  $\vec{v} = (-8, 4, -16)$   
Orthogonal - ?

$$\vec{u} \cdot \vec{v} = \|\vec{u}\| \cdot \|\vec{v}\| \cdot \cos \theta = 0 \text{ if } \theta = 90^\circ$$

$$\vec{u} \cdot \vec{v} = 2 \cdot (-8) - 1 \cdot 4 + 4 \cdot (-16) = \\ = -16 - 4 - 64 = -84 \neq 0$$

Orthogonal