

Statistics 153 - Homework 4

Due on Wednesday, October 27, 3 pm

1 Theory: Bartlett's Formula

- (a) Assuming its conditions are met, show that for an ARMA(p, q) process X_t with $p = q = 0$ (ie. X_t is white noise) Bartlett's formula gives the following result: (10 points)

$$\sqrt{n} \begin{pmatrix} \hat{r}_1 \\ \vdots \\ \hat{r}_k \end{pmatrix} \xrightarrow{d} N_k(0, I_k)$$

***This is the asymptotic result for the sample correlations of white noise covered earlier in class*

- (b) For the AR(1) process $X_t = \phi X_{t-1} + W_t$, calculate the asymptotic variance of \hat{r}_k for $k \geq 1$. Note that, technically, the autocorrelation for AR(1) is $\rho(h) = \phi^{|h|}$, where the absolute value of h is in the exponent. (15 points)

2 Computation: Bartlett's Formula

Here you will use the results for the AR(1) model that you derived above. Assume $\phi = 0.7$.

- (a) Simulate $n = 300$ observations of the process using `arma.sim()`. Plot the sample ACF of this simulated stationary process. Note the blue dashed lines at $\pm 1.96/\sqrt{n}$ bars: these are the default for the `acf()` function and represent the distribution of r_h for white noise, aka ARMA(0,0). (5 point)
- (b) Now let's visualize the distribution of ARMA(1,0), usually simply called AR(1). Add to this plot the expectations and 95% intervals for r_1, \dots, r_{20} in red. Comment on the difference between the blue lines and the red lines. (10 points)

3 Prediction of MA(1)

Consider an invertible MA(1) model $X_t = W_t + \theta W_{t-1}$ for some i.i.d. white noise process $\{W_t\}$ with variance σ^2 .

- (a) Derive the explicit form of the minimum mean-square error one-step prediction

$$\tilde{X}_{n+1} = E(X_{n+1} | X_n, X_{n-1}, X_{n-2}, \dots)$$

for X_{n+1} based on the complete infinite past $X_n, X_{n-1}, X_{n-2}, \dots$. Hint: invertible processes can be put in $AR(\infty)$ form, right?

(15 points)

- (b) Derive the mean squared error $E[(\tilde{X}_{n+1} - X_{n+1})^2]$. (10 point)

- (c) Now consider the truncated estimate \tilde{X}_{n+1}^n , which equals \tilde{X}_{n+1} but with unobserved data being set to zero, that is, $0 = X_0 = X_{-1} = \dots$. Show that (15 points)

$$E[(X_{n+1} - \tilde{X}_{n+1}^n)^2] = \sigma^2(1 + \theta^{2+2n}).$$

- (d) Comment on how well the truncated estimate \tilde{X}_{n+1}^n works compared to \tilde{X}_{n+1} (i.e. compare the MSE's of the two). (5 point)

4 Prediction of MA(q)

Consider an invertible MA(q) model $X_t = \theta(B)W_t$ for some white noise $\{W_t\}$ with variance σ^2 .

- (a) Show that for any $m > q$ the best linear predictor of X_{n+m} based on X_1, \dots, X_n is always zero. (8 point)
- (b) Now assume that the white noise $\{W_t\}$ is also i.i.d.. Show that for any $m > q$ the best predictor (minimum mean-square error forecast) of X_{n+m} based on the full history $X_n, X_{n-1}, X_{n-2}, \dots$ is also zero. (7 point)