

Randomness and determinism in Diophantine approximation: small linear forms, lattice flows and some applications

Victor Beresnevich

**Department of Mathematics
University of York**

Imperial College London

18 January 2017

PLAN:

PLAN:

My main goal: To have a discussion of Diophantine approximation (DA):

PLAN:

My main goal: To have a discussion of Diophantine approximation (DA):

- Small linear forms, badly approximable systems

PLAN:

My main goal: To have a discussion of Diophantine approximation (DA):

- Small linear forms, badly approximable systems
- One dimensional DA and continued fractions

PLAN:

My main goal: To have a discussion of Diophantine approximation (DA):

- Small linear forms, badly approximable systems
- One dimensional DA and continued fractions
- Multiplicative DA

PLAN:

My main goal: To have a discussion of Diophantine approximation (DA):

- Small linear forms, badly approximable systems
- One dimensional DA and continued fractions
- Multiplicative DA
- Metric DA, the Khintchine-Groshev theorem,

PLAN:

My main goal: To have a discussion of Diophantine approximation (DA):

- Small linear forms, badly approximable systems
- One dimensional DA and continued fractions
- Multiplicative DA
- Metric DA, the Khintchine-Groshev theorem,
- DA on manifolds, and

My main goal: To have a discussion of Diophantine approximation (DA):

- Small linear forms, badly approximable systems
- One dimensional DA and continued fractions
- Multiplicative DA
- Metric DA, the Khintchine-Groshev theorem,
- DA on manifolds, and
- Effective metric theorems

PLAN:

My main goal: To have a discussion of Diophantine approximation (DA):

- Small linear forms, badly approximable systems
- One dimensional DA and continued fractions
- Multiplicative DA
- Metric DA, the Khintchine-Groshev theorem,
- DA on manifolds, and
- Effective metric theorems

Relevance to coding and lattices:

My main goal: To have a discussion of Diophantine approximation (DA):

- Small linear forms, badly approximable systems
- One dimensional DA and continued fractions
- Multiplicative DA
- Metric DA, the Khintchine-Groshev theorem,
- DA on manifolds, and
- Effective metric theorems

Relevance to coding and lattices:

- DA is of growing interest in electronics: lattice coding and interference alignment (the latter to be briefly described);

My main goal: To have a discussion of Diophantine approximation (DA):

- Small linear forms, badly approximable systems
- One dimensional DA and continued fractions
- Multiplicative DA
- Metric DA, the Khintchine-Groshev theorem,
- DA on manifolds, and
- Effective metric theorems

Relevance to coding and lattices:

- DA is of growing interest in electronics: lattice coding and interference alignment (the latter to be briefly described);
- Minkowski's theorems is a basis for much of DA, and a useful tool at the same time;

My main goal: To have a discussion of Diophantine approximation (DA):

- Small linear forms, badly approximable systems
- One dimensional DA and continued fractions
- Multiplicative DA
- Metric DA, the Khintchine-Groshev theorem,
- DA on manifolds, and
- Effective metric theorems

Relevance to coding and lattices:

- DA is of growing interest in electronics: lattice coding and interference alignment (the latter to be briefly described);
- Minkowski's theorems is a basis for much of DA, and a useful tool at the same time;
- Dictionary between Homogeneous dynamics (lattice orbits) and Diophantine Approximation (will be discussed where possible)

The idea of interference alignment

The idea of interference alignment

Consider the system of linear equations:

$$\begin{cases} y_1 = h_{11}x_1 + \cdots + h_{1K}x_K \\ \vdots \\ y_B = h_{B1}x_1 + \cdots + h_{BK}x_K \end{cases}$$

The idea of interference alignment

Consider the system of linear equations:

$$\begin{cases} y_1 = h_{11}x_1 + \cdots + h_{1K}x_K \\ \vdots \\ y_B = h_{B1}x_1 + \cdots + h_{BK}x_K \end{cases}$$

Interpretation: h_{ij} are channel coefficients

x_1, \dots, x_K are signals transmitted from K different transmitters

y_1, \dots, y_B are signals received at a receiver

B is the bandwidth (number of signaling dimensions)

The idea of interference alignment

Consider the system of linear equations:

$$\begin{cases} y_1 = h_{11}x_1 + \cdots + h_{1K}x_K \\ \vdots \\ y_B = h_{B1}x_1 + \cdots + h_{BK}x_K \end{cases}$$

Interpretation: h_{ij} are channel coefficients

x_1, \dots, x_K are signals transmitted from K different transmitters

y_1, \dots, y_B are signals received at a receiver

B is the bandwidth (number of signaling dimensions)

$$\text{If } \mathbf{rank}(h_{ij}) \geq K$$

the receiver can recover all the symbols x_1, \dots, x_K . This requires that

$$B \geq K.$$

The idea of interference alignment

Consider the system of linear equations:

$$\begin{cases} y_1 = h_{11}x_1 + \cdots + h_{1K}x_K \\ \vdots \\ y_B = h_{B1}x_1 + \cdots + h_{BK}x_K \end{cases}$$

Interpretation: h_{ij} are channel coefficients

x_1, \dots, x_K are signals transmitted from K different transmitters

y_1, \dots, y_B are signals received at a receiver

B is the bandwidth (number of signaling dimensions)

$$\text{If } \mathbf{rank}(h_{ij}) \geq K$$

the receiver can recover all the symbols x_1, \dots, x_K . This requires that

$$B \geq K.$$

In IC the receiver may be interested only in some of the transmitted symbols, maybe only x_1 , while the other symbols are intended for other receivers. Even then we still require B signaling dimensions to recover x_1 .

The idea of interference alignment

Consider the system of linear equations:

$$\begin{cases} y_1 = h_{11}x_1 + \cdots + h_{1K}x_K \\ \vdots \\ y_B = h_{B1}x_1 + \cdots + h_{BK}x_K \end{cases}$$

Interpretation: h_{ij} are channel coefficients

x_1, \dots, x_K are signals transmitted from K different transmitters

y_1, \dots, y_B are signals received at a receiver

B is the bandwidth (number of signaling dimensions)

$$\text{If } \mathbf{rank}(h_{ij}) \geq K$$

the receiver can recover all the symbols x_1, \dots, x_K . This requires that

$$B \geq K.$$

In IC the receiver may be interested only in some of the transmitted symbols, maybe only x_1 , while the other symbols are intended for other receivers. Even then we still require B signaling dimensions to recover x_1 .

\Rightarrow The bandwidth is simply divided between the users.

The idea of interference alignment

Let

$$\mathbf{H}_i = \begin{pmatrix} h_{1i} \\ \vdots \\ h_{Bi} \end{pmatrix} \quad \text{and} \quad \mathbf{Y} = \begin{pmatrix} y_1 \\ \vdots \\ y_B \end{pmatrix}$$

The idea of interference alignment

Let

$$\mathbf{H}_i = \begin{pmatrix} h_{1i} \\ \vdots \\ h_{Bi} \end{pmatrix} \quad \text{and} \quad \mathbf{Y} = \begin{pmatrix} y_1 \\ \vdots \\ y_B \end{pmatrix}$$

Then the system is

$$\mathbf{Y} = x_1 \mathbf{H}_1 + \underbrace{(x_2 \mathbf{H}_2 + \cdots + x_K \mathbf{H}_K)}_{\in \text{Span}\{\mathbf{H}_2, \dots, \mathbf{H}_K\}}$$

If

$$\mathbf{H}_1 \notin \text{Span}\{\mathbf{H}_2, \dots, \mathbf{H}_K\}$$

then x_1 can be uniquely determined from the above equations.

The idea of interference alignment

Let

$$\mathbf{H}_i = \begin{pmatrix} h_{1i} \\ \vdots \\ h_{Bi} \end{pmatrix} \quad \text{and} \quad \mathbf{Y} = \begin{pmatrix} y_1 \\ \vdots \\ y_B \end{pmatrix}$$

Then the system is

$$\mathbf{Y} = x_1 \mathbf{H}_1 + \underbrace{(x_2 \mathbf{H}_2 + \cdots + x_K \mathbf{H}_K)}_{\in \text{Span}\{\mathbf{H}_2, \dots, \mathbf{H}_K\}}$$

If

$$\mathbf{H}_1 \notin \text{Span}\{\mathbf{H}_2, \dots, \mathbf{H}_K\}$$

then x_1 can be uniquely determined from the above equations.

Cadambe & Jafar (2008) showed that it is theoretically possible that each of the K users enjoy half of the available signalling dimensions.

The idea of interference alignment

Let

$$\mathbf{H}_i = \begin{pmatrix} h_{1i} \\ \vdots \\ h_{Bi} \end{pmatrix} \quad \text{and} \quad \mathbf{Y} = \begin{pmatrix} y_1 \\ \vdots \\ y_B \end{pmatrix}$$

Then the system is

$$\mathbf{Y} = x_1 \mathbf{H}_1 + \underbrace{(x_2 \mathbf{H}_2 + \cdots + x_K \mathbf{H}_K)}_{\in \text{Span}\{\mathbf{H}_2, \dots, \mathbf{H}_K\}}$$

If

$$\mathbf{H}_1 \notin \text{Span}\{\mathbf{H}_2, \dots, \mathbf{H}_K\}$$

then x_1 can be uniquely determined from the above equations.

Cadambe & Jafar (2008) showed that it is theoretically possible that each of the K users enjoy half of the available signalling dimensions. More formally,

$$DOF = \frac{K}{2}.$$

The idea of interference alignment

Let

$$\mathbf{H}_i = \begin{pmatrix} h_{1i} \\ \vdots \\ h_{Bi} \end{pmatrix} \quad \text{and} \quad \mathbf{Y} = \begin{pmatrix} y_1 \\ \vdots \\ y_B \end{pmatrix}$$

Then the system is

$$\mathbf{Y} = x_1 \mathbf{H}_1 + \underbrace{(x_2 \mathbf{H}_2 + \cdots + x_K \mathbf{H}_K)}_{\in \text{Span}\{\mathbf{H}_2, \dots, \mathbf{H}_K\}}$$

If

$$\mathbf{H}_1 \notin \text{Span}\{\mathbf{H}_2, \dots, \mathbf{H}_K\}$$

then x_1 can be uniquely determined from the above equations.

Cadambe & Jafar (2008) showed that it is theoretically possible that each of the K users enjoy half of the available signalling dimensions. More formally,

$$DOF = \frac{K}{2}.$$

Various assumptions:

- varying channel coefficients,
- multiple antennae at receivers
- diagonal form of the channel coefficient matrices
- ...

Rational dimensions framework

Assume the transmit signals are taken from the constellation $\{0, \dots, Q\}$.

Rational dimensions framework

Assume the transmit signals are taken from the constellation $\{0, \dots, Q\}$.

By observing only

$$y = h_1 x_1 + \dots + h_K x_K$$

at the receiver we still can decode our signals, if h_1, \dots, h_K are linearly independent over \mathbb{Q} (or equivalently \mathbb{Z}).

Rational dimensions framework

Assume the transmit signals are taken from the constellation $\{0, \dots, Q\}$.

By observing only

$$y = h_1 x_1 + \dots + h_K x_K$$

at the receiver we still can decode our signals, if h_1, \dots, h_K are linearly independent over \mathbb{Q} (or equivalently \mathbb{Z}). Independence over \mathbb{Q} means that any linear combination $h_1 x_1 + \dots + h_K x_K$ with integer x_1, \dots, x_K will only vanish when all the x_i are zeros.

Rational dimensions framework

Assume the transmit signals are taken from the constellation $\{0, \dots, Q\}$.

By observing only

$$y = h_1 x_1 + \dots + h_K x_K$$

at the receiver we still can decode our signals, if h_1, \dots, h_K are linearly independent over \mathbb{Q} (or equivalently \mathbb{Z}). Independence over \mathbb{Q} means that any linear combination $h_1 x_1 + \dots + h_K x_K$ with integer x_1, \dots, x_K will only vanish when all the x_i are zeros.

If there is noise z , the receiver observes

$$y = h_1 x_1 + \dots + h_K x_K + z.$$

Rational dimensions framework

Assume the transmit signals are taken from the constellation $\{0, \dots, Q\}$.

By observing only

$$y = h_1 x_1 + \dots + h_K x_K$$

at the receiver we still can decode our signals, if h_1, \dots, h_K are linearly independent over \mathbb{Q} (or equivalently \mathbb{Z}). Independence over \mathbb{Q} means that any linear combination $h_1 x_1 + \dots + h_K x_K$ with integer x_1, \dots, x_K will only vanish when all the x_i are zeros.

If there is noise z , the receiver observes

$$y = h_1 x_1 + \dots + h_K x_K + z.$$

This observation can be distinguished from another one, say

$$y' = h_1 x'_1 + \dots + h_K x'_K + z'$$

only if y and y' are sufficiently separated.

Rational dimensions framework

Assume the transmit signals are taken from the constellation $\{0, \dots, Q\}$.

By observing only

$$y = h_1 x_1 + \dots + h_K x_K$$

at the receiver we still can decode our signals, if h_1, \dots, h_K are linearly independent over \mathbb{Q} (or equivalently \mathbb{Z}). Independence over \mathbb{Q} means that any linear combination $h_1 x_1 + \dots + h_K x_K$ with integer x_1, \dots, x_K will only vanish when all the x_i are zeros.

If there is noise z , the receiver observes

$$y = h_1 x_1 + \dots + h_K x_K + z.$$

This observation can be distinguished from another one, say

$$y' = h_1 x'_1 + \dots + h_K x'_K + z'$$

only if y and y' are sufficiently separated.

Fact: for any $\varepsilon > 0$ for almost every collection (h_1, \dots, h_K) of real numbers there exists a constant $\gamma = \gamma(\varepsilon, h_1, \dots, h_K) > 0$ such that

$$|h_1(x_1 - x'_1) + \dots + h_K(x_K - x'_K)| \geq \frac{\gamma}{Q^{K-1+\varepsilon}}$$

Using that **Fact** a single antenna can be turned into a multiple antennae to simultaneously transmit several data streams, say x_1, \dots, x_M , $x_i \in \{0, \dots, Q\}$.

Using that **Fact** a single antenna can be turned into a multiple antennae to simultaneously transmit several data streams, say x_1, \dots, x_M , $x_i \in \{0, \dots, Q\}$. The transmit signal may be taken to be

$$x = \lambda(c_1x_1 + \dots + c_Mx_M)$$

where $|c_i| = O(1)$ are precoding coefficients and λ is a scalar reflecting power constraints.

Using that **Fact** a single antenna can be turned into a multiple antennae to simultaneously transmit several data streams, say x_1, \dots, x_M , $x_i \in \{0, \dots, Q\}$. The transmit signal may be taken to be

$$x = \lambda(c_1x_1 + \dots + c_Mx_M)$$

where $|c_i| = O(1)$ are precoding coefficients and λ is a scalar reflecting power constraints. The separation between different x 's is at least $\lambda O(Q^{-M+1-\varepsilon})$.

Using that **Fact** a single antenna can be turned into a multiple antennae to simultaneously transmit several data streams, say x_1, \dots, x_M , $x_i \in \{0, \dots, Q\}$. The transmit signal may be taken to be

$$x = \lambda(c_1 x_1 + \dots + c_M x_M)$$

where $|c_i| = O(1)$ are precoding coefficients and λ is a scalar reflecting power constraints. The separation between different x 's is at least $\lambda O(Q^{-M+1-\varepsilon})$. Choosing

$$Q = O(P^{\frac{1-\varepsilon}{2(M+\varepsilon)}}) \quad \text{and} \quad \lambda = P^{1/2} Q^{-1}$$

ensures that the separation is bigger than the standard deviation of the noise, while meeting power constraints.

Using that **Fact** a single antenna can be turned into a multiple antennae to simultaneously transmit several data streams, say x_1, \dots, x_M , $x_i \in \{0, \dots, Q\}$. The transmit signal may be taken to be

$$x = \lambda(c_1x_1 + \dots + c_Mx_M)$$

where $|c_i| = O(1)$ are precoding coefficients and λ is a scalar reflecting power constraints. The separation between different x 's is at least $\lambda O(Q^{-M+1-\varepsilon})$. Choosing

$$Q = O(P^{\frac{1-\varepsilon}{2(M+\varepsilon)}}) \quad \text{and} \quad \lambda = P^{1/2}Q^{-1}$$

ensures that the separation is bigger than the standard deviation of the noise, while meeting power constraints.

More sophisticated examples require separability of linear forms when the coefficients the forms are functions of several other variables (in DA this is known as Diophantine approximation on manifolds).

Rational dimensions: single antenna, multiple data streams

Using that **Fact** a single antenna can be turned into a multiple antennae to simultaneously transmit several data streams, say x_1, \dots, x_M , $x_i \in \{0, \dots, Q\}$. The transmit signal may be taken to be

$$x = \lambda(c_1x_1 + \dots + c_Mx_M)$$

where $|c_i| = O(1)$ are precoding coefficients and λ is a scalar reflecting power constraints. The separation between different x 's is at least $\lambda O(Q^{-M+1-\varepsilon})$. Choosing

$$Q = O(P^{\frac{1-\varepsilon}{2(M+\varepsilon)}}) \quad \text{and} \quad \lambda = P^{1/2}Q^{-1}$$


ensures that the separation is bigger than the standard deviation of the noise, while meeting power constraints.

More sophisticated examples require separability of linear forms when the coefficients the forms are functions of several other variables (in DA this is known as Diophantine approximation on manifolds).


Just as with vector alignment, it is possible to align alone real numbers.

2-user X-channel (Motahari, *et al*)

2-user X-channel (Motahari, *et al*)




2-user X-channel (Motahari, *et al*)



T_1 simultaneously transmits two data streams u_1 (intended for R_1) and v_1 (intended for R_2). Similarly, T_2 transmits independent two data streams u_2 (intended for R_1) and v_2 (intended for R_2). h_{ij} are the the channel coefficients. Let x_i is the signal transmitted by T_i .

2-user X-channel (Motahari, *et al*)



T_1 simultaneously transmits two data streams u_1 (intended for R_1) and v_1 (intended for R_2). Similarly, T_2 transmits independent two data streams u_2 (intended for R_1) and v_2 (intended for R_2). h_{ij} are the the channel coefficients. Let x_i is the signal transmitted by T_i . The received signals by R_1 and R_2 are

$$y_1 = h_{11}x_1 + h_{12}x_2 + z_1,$$

$$y_2 = h_{21}x_1 + h_{22}x_2 + z_2.$$

$$y_1 = h_{11}x_1 + h_{12}x_2 + z_1 ,$$

$$y_2 = h_{21}x_1 + h_{22}x_2 + z_2 .$$

2-user X-channel (Motahari, *et al*)

$$y_1 = h_{11}x_1 + h_{12}x_2 + z_1 ,$$

$$y_2 = h_{21}x_1 + h_{22}x_2 + z_2 .$$

If $x_1 = \lambda(h_{22}u_1 + h_{12}v_1)$ and $x_2 = \lambda(h_{21}u_2 + h_{11}v_2)$ then

$$y_1 = \lambda \left(h_{11}h_{22}u_1 + h_{12}h_{21}u_2 + h_{11}h_{12}(v_1 + v_2) \right) + z_1 ,$$

$$y_2 = \lambda \left(h_{21}h_{22}(u_1 + u_2) + h_{12}h_{21}v_1 + h_{11}h_{22}v_2 \right) + z_1$$

At R_1 : v_1 and v_2 are aligned along the same real number, $h_{11}h_{12}$

At R_2 : u_1 and u_2 are aligned along the same real number, $h_{21}h_{22}$

2-user X-channel (Motahari, *et al*)

$$y_1 = h_{11}x_1 + h_{12}x_2 + z_1,$$

$$y_2 = h_{21}x_1 + h_{22}x_2 + z_2.$$

If $x_1 = \lambda(h_{22}u_1 + h_{12}v_1)$ and $x_2 = \lambda(h_{21}u_2 + h_{11}v_2)$ then

$$y_1 = \lambda(h_{11}h_{22}u_1 + h_{12}h_{21}u_2 + h_{11}h_{12}(v_1 + v_2)) + z_1,$$

$$y_2 = \lambda(h_{21}h_{22}(u_1 + u_2) + h_{12}h_{21}v_1 + h_{11}h_{22}v_2) + z_1$$

At R_1 : v_1 and v_2 are aligned along the same real number, $h_{11}h_{12}$

At R_2 : u_1 and u_2 are aligned along the same real number, $h_{21}h_{22}$

Again, assuming $u_i, v_i \in \{0, \dots, Q\}$ we achieve the required separation (and normalising power) in each of the equation by taking

$$Q = O(P^{\frac{1-\varepsilon}{2(3+\varepsilon)}}), \quad \lambda = O(P^{\frac{1+\varepsilon}{3+\varepsilon}}).$$

DOF = $\frac{4}{3}$ almost surely.

Diophantine approximation: the basics

In applications we often deal with small linear forms or systems of small linear forms at integer variables -

Diophantine approximation: the basics

In applications we often deal with small linear forms or systems of small linear forms at integer variables - a subject of Diophantine approximation.

Diophantine approximation: the basics

In applications we often deal with small linear forms or systems of small linear forms at integer variables - a subject of Diophantine approximation.

Dirichlet's theorem: Let $\alpha_{i,j} \in \mathbb{R}$, where $1 \leq j \leq n$, $1 \leq i \leq m$, and $Q > 1$. Then there exist $q_1, \dots, q_n, p_1, \dots, p_m \in \mathbb{Z}$ such that

$$|q_1\alpha_{i,1} + \dots + q_n\alpha_{i,n} - p_i| < Q^{-\frac{n}{m}} \quad (1 \leq i \leq m)$$

$$1 \leq \max_{1 \leq j \leq n} |q_j| \leq Q.$$

Diophantine approximation: the basics

In applications we often deal with small linear forms or systems of small linear forms at integer variables - a subject of Diophantine approximation.

Dirichlet's theorem: Let $\alpha_{i,j} \in \mathbb{R}$, where $1 \leq j \leq n$, $1 \leq i \leq m$, and $Q > 1$. Then there exist $q_1, \dots, q_n, p_1, \dots, p_m \in \mathbb{Z}$ such that

$$|q_1 \alpha_{i,1} + \dots + q_n \alpha_{i,n} - p_i| < Q^{-\frac{n}{m}} \quad (1 \leq i \leq m)$$
$$1 \leq \max_{1 \leq j \leq n} |q_j| \leq Q.$$

Minkowski's theorem for systems of linear forms: Let $\beta_{i,j} \in \mathbb{R}$, where $1 \leq i, j \leq k$, and let $C_1, \dots, C_k > 0$. If

$$|\det(\beta_{i,j})_{1 \leq i,j \leq k}| \leq \prod_{i=1}^k C_i, \quad (1)$$

then there exist a non-zero integer point $\mathbf{x} = (x_1, \dots, x_k)$ such that

$$\begin{cases} |x_1 \beta_{i,1} + \dots + x_k \beta_{i,k}| < C_i, & (1 \leq i \leq k-1) \\ |x_1 \beta_{k,1} + \dots + x_k \beta_{k,k}| \leq C_k. \end{cases} \quad (2)$$

Diophantine approximation: the basics

In applications we often deal with small linear forms or systems of small linear forms at integer variables - a subject of Diophantine approximation.

Dirichlet's theorem: Let $\alpha_{i,j} \in \mathbb{R}$, where $1 \leq j \leq n$, $1 \leq i \leq m$, and $Q > 1$. Then there exist $q_1, \dots, q_n, p_1, \dots, p_m \in \mathbb{Z}$ such that

$$|q_1\alpha_{i,1} + \dots + q_n\alpha_{i,n} - p_i| < Q^{-\frac{n}{m}} \quad (1 \leq i \leq m)$$
$$1 \leq \max_{1 \leq j \leq n} |q_j| \leq Q.$$

Minkowski's theorem for systems of linear forms: Let $\beta_{i,j} \in \mathbb{R}$, where $1 \leq i, j \leq k$, and let $C_1, \dots, C_k > 0$. If

$$|\det(\beta_{i,j})_{1 \leq i,j \leq k}| \leq \prod_{i=1}^k C_i, \quad (1)$$

then there exist a non-zero integer point $\mathbf{x} = (x_1, \dots, x_k)$ such that

$$\begin{cases} |x_1\beta_{i,1} + \dots + x_k\beta_{i,k}| < C_i, & (1 \leq i \leq k-1) \\ |x_1\beta_{k,1} + \dots + x_k\beta_{k,k}| \leq C_k. \end{cases} \quad (2)$$

Proof: uses Minkowski's theorem for convex bodies.

Minkowski's theorem: If B is a convex body in \mathbb{R}^n symmetric about $\mathbf{0}$ and $\text{vol } B > 2^n$ then B contains a non-zero integer point.

Minkowski's theorem: If B is a convex body in \mathbb{R}^n symmetric about $\mathbf{0}$ and $\text{vol } B > 2^n$ then B contains a non-zero integer point.

Minkowski can be used beyond the real case. Example:

Dirichlet's theorem for \mathbb{C} (one dimensional): For any $z \in \mathbb{C}$ and any $Q > 1$ there exist $p, q \in \mathbb{Z}[i] = \{a + bi : a, b \in \mathbb{Z}\}$ ($i = \sqrt{-1}$) such that

$$|qz - p| < \frac{4}{\pi Q}, \quad 1 \leq |q| \leq Q.$$

Diophantine approximation: the basics

Minkowski's theorem: If B is a convex body in \mathbb{R}^n symmetric about $\mathbf{0}$ and $\text{vol } B > 2^n$ then B contains a non-zero integer point.

Minkowski can be used beyond the real case. Example:

Dirichlet's theorem for \mathbb{C} (one dimensional): For any $z \in \mathbb{C}$ and any $Q > 1$ there exist $p, q \in \mathbb{Z}[i] = \{a + bi : a, b \in \mathbb{Z}\}$ ($i = \sqrt{-1}$) such that

$$|qz - p| < \frac{4}{\pi Q}, \quad 1 \leq |q| \leq Q.$$

Badly approximable systems/matrices: Let $\alpha_{i,j} \in \mathbb{R}$, where $1 \leq j \leq n$, $1 \leq i \leq m$, and $Q > 1$. Then $A = (\alpha_{i,j})_{i,j}$ is badly approximable if there exist $c > 0$ such that for all $Q > 1$ the only integer solution $(q_1, \dots, q_n, p_1, \dots, p_m)$ to the system

$$\begin{aligned} |q_1\alpha_{i,1} + \dots + q_n\alpha_{i,n} - p_i| &< cQ^{-\frac{n}{m}} \quad (1 \leq i \leq m) \\ |q_i| &\leq Q \quad (1 \leq j \leq n) \end{aligned}$$

is zero.

Diophantine approximation: the basics

Minkowski's theorem: If B is a convex body in \mathbb{R}^n symmetric about $\mathbf{0}$ and $\text{vol } B > 2^n$ then B contains a non-zero integer point.

Minkowski can be used beyond the real case. Example:

Dirichlet's theorem for \mathbb{C} (one dimensional): For any $z \in \mathbb{C}$ and any $Q > 1$ there exist $p, q \in \mathbb{Z}[i] = \{a + bi : a, b \in \mathbb{Z}\}$ ($i = \sqrt{-1}$) such that

$$|qz - p| < \frac{4}{\pi Q}, \quad 1 \leq |q| \leq Q.$$

Badly approximable systems/matrices: Let $\alpha_{i,j} \in \mathbb{R}$, where $1 \leq j \leq n$, $1 \leq i \leq m$, and $Q > 1$. Then $A = (\alpha_{i,j})_{i,j}$ is badly approximable if there exist $c > 0$ such that for all $Q > 1$ the only integer solution $(q_1, \dots, q_n, p_1, \dots, p_m)$ to the system

$$\begin{aligned} |q_1\alpha_{i,1} + \dots + q_n\alpha_{i,n} - p_i| &< cQ^{-\frac{n}{m}} \quad (1 \leq i \leq m) \\ |q_i| &\leq Q \quad (1 \leq j \leq n) \end{aligned}$$

is zero. **Bad** (n, m) is the set of badly approximable $m \times n$ matrices.

Dirichlet's theorem with weights: Let $\alpha_{i,j} \in \mathbb{R}$, where $1 \leq j \leq n$, $1 \leq i \leq m$. Let $c = 1$ and let $\mathbf{r} = (r_1, \dots, r_n)$, $\mathbf{s} = (s_1, \dots, s_m)$ be such that

$$s_j \geq 0, \quad r_i \geq 0, \quad s_1 + \dots + s_m = 1, \quad r_1 + \dots + r_n = 1.$$

Then for any $Q > 1$ there exist $q_1, \dots, q_n, p_1, \dots, p_m \in \mathbb{Z}$ such that

$$|q_1 \alpha_{i,1} + \dots + q_n \alpha_{i,n} - p_i| < c Q^{-s_i} \quad (1 \leq i \leq m)$$

$$|q_j| \leq Q^{r_j} \quad (1 \leq j \leq n)$$

$$(q_1, \dots, q_n) \neq \mathbf{0}.$$

Dirichlet's theorem with weights: Let $\alpha_{i,j} \in \mathbb{R}$, where $1 \leq j \leq n$, $1 \leq i \leq m$. Let $c = 1$ and let $\mathbf{r} = (r_1, \dots, r_n)$, $\mathbf{s} = (s_1, \dots, s_m)$ be such that

$$s_j \geq 0, \quad r_i \geq 0, \quad s_1 + \dots + s_m = 1, \quad r_1 + \dots + r_n = 1.$$

Then for any $Q > 1$ there exist $q_1, \dots, q_n, p_1, \dots, p_m \in \mathbb{Z}$ such that

$$|q_1 \alpha_{i,1} + \dots + q_n \alpha_{i,n} - p_i| < c Q^{-s_i} \quad (1 \leq i \leq m)$$

$$|q_j| \leq Q^{r_j} \quad (1 \leq j \leq n)$$

$$(q_1, \dots, q_n) \neq \mathbf{0}.$$

Weighted Badly approximable systems/matrices: Let $\alpha_{i,j} \in \mathbb{R}$, where $1 \leq j \leq n$, $1 \leq i \leq m$, and $Q > 1$. Then $A = (\alpha_{i,j})_{i,j}$ is (\mathbf{r}, \mathbf{s}) -badly approximable if there exist $c > 0$ such that for all $Q > 1$ the only integer solution $(q_1, \dots, q_n, p_1, \dots, p_m)$ to the above system is zero.

Dirichlet's theorem with weights: Let $\alpha_{i,j} \in \mathbb{R}$, where $1 \leq j \leq n$, $1 \leq i \leq m$. Let $c = 1$ and let $\mathbf{r} = (r_1, \dots, r_n)$, $\mathbf{s} = (s_1, \dots, s_m)$ be such that

$$s_j \geq 0, \quad r_i \geq 0, \quad s_1 + \dots + s_m = 1, \quad r_1 + \dots + r_n = 1.$$

Then for any $Q > 1$ there exist $q_1, \dots, q_n, p_1, \dots, p_m \in \mathbb{Z}$ such that

$$|q_1 \alpha_{i,1} + \dots + q_n \alpha_{i,n} - p_i| < c Q^{-s_i} \quad (1 \leq i \leq m)$$

$$|q_j| \leq Q^{r_j} \quad (1 \leq j \leq n)$$

$$(q_1, \dots, q_n) \neq \mathbf{0}.$$

Weighted Badly approximable systems/matrices: Let $\alpha_{i,j} \in \mathbb{R}$, where $1 \leq j \leq n$, $1 \leq i \leq m$, and $Q > 1$. Then $A = (\alpha_{i,j})_{i,j}$ is (\mathbf{r}, \mathbf{s}) -badly approximable if there exist $c > 0$ such that for all $Q > 1$ the only integer solution $(q_1, \dots, q_n, p_1, \dots, p_m)$ to the above system is zero.

Let $\mathbf{Bad}(\mathbf{r}, \mathbf{s})$ be the set of (\mathbf{r}, \mathbf{s}) -badly approximable $m \times n$ matrices.

Dirichlet's theorem with weights: Let $\alpha_{i,j} \in \mathbb{R}$, where $1 \leq j \leq n$, $1 \leq i \leq m$. Let $c = 1$ and let $\mathbf{r} = (r_1, \dots, r_n)$, $\mathbf{s} = (s_1, \dots, s_m)$ be such that

$$s_j \geq 0, \quad r_i \geq 0, \quad s_1 + \dots + s_m = 1, \quad r_1 + \dots + r_n = 1.$$

Then for any $Q > 1$ there exist $q_1, \dots, q_n, p_1, \dots, p_m \in \mathbb{Z}$ such that

$$|q_1 \alpha_{i,1} + \dots + q_n \alpha_{i,n} - p_i| < c Q^{-s_i} \quad (1 \leq i \leq m)$$

$$|q_j| \leq Q^{r_j} \quad (1 \leq j \leq n)$$

$$(q_1, \dots, q_n) \neq \mathbf{0}.$$

Weighted Badly approximable systems/matrices: Let $\alpha_{i,j} \in \mathbb{R}$, where $1 \leq j \leq n$, $1 \leq i \leq m$, and $Q > 1$. Then $A = (\alpha_{i,j})_{i,j}$ is (\mathbf{r}, \mathbf{s}) -badly approximable if there exist $c > 0$ such that for all $Q > 1$ the only integer solution $(q_1, \dots, q_n, p_1, \dots, p_m)$ to the above system is zero.

Let $\mathbf{Bad}(\mathbf{r}, \mathbf{s})$ be the set of (\mathbf{r}, \mathbf{s}) -badly approximable $m \times n$ matrices.

$$\mathbf{Bad}(n, m) = \mathbf{Bad}\left(\left(\frac{1}{n}, \dots, \frac{1}{n}\right), \left(\frac{1}{m}, \dots, \frac{1}{m}\right)\right).$$

Dirichlet's theorem with weights: Let $\alpha_{i,j} \in \mathbb{R}$, where $1 \leq j \leq n$, $1 \leq i \leq m$. Let $c = 1$ and let $\mathbf{r} = (r_1, \dots, r_n)$, $\mathbf{s} = (s_1, \dots, s_m)$ be such that

$$s_j \geq 0, \quad r_i \geq 0, \quad s_1 + \dots + s_m = 1, \quad r_1 + \dots + r_n = 1.$$

Then for any $Q > 1$ there exist $q_1, \dots, q_n, p_1, \dots, p_m \in \mathbb{Z}$ such that

$$|q_1 \alpha_{i,1} + \dots + q_n \alpha_{i,n} - p_i| < c Q^{-s_i} \quad (1 \leq i \leq m)$$

$$|q_j| \leq Q^{r_j} \quad (1 \leq j \leq n)$$

$$(q_1, \dots, q_n) \neq \mathbf{0}.$$

Weighted Badly approximable systems/matrices: Let $\alpha_{i,j} \in \mathbb{R}$, where $1 \leq j \leq n$, $1 \leq i \leq m$, and $Q > 1$. Then $A = (\alpha_{i,j})_{ij}$ is (\mathbf{r}, \mathbf{s}) -badly approximable if there exist $c > 0$ such that for all $Q > 1$ the only integer solution $(q_1, \dots, q_n, p_1, \dots, p_m)$ to the above system is zero.

Let $\mathbf{Bad}(\mathbf{r}, \mathbf{s})$ be the set of (\mathbf{r}, \mathbf{s}) -badly approximable $m \times n$ matrices.

$$\mathbf{Bad}(n, m) = \mathbf{Bad}\left(\left(\frac{1}{n}, \dots, \frac{1}{n}\right), \left(\frac{1}{m}, \dots, \frac{1}{m}\right)\right).$$

The transpose of $\mathbf{Bad}(\mathbf{r}, \mathbf{s})$ is $\mathbf{Bad}(\mathbf{s}, \mathbf{r})$.

Weighted Diophantine approximation

Dirichlet's theorem with weights: Let $\alpha_{i,j} \in \mathbb{R}$, where $1 \leq j \leq n$, $1 \leq i \leq m$. Let $c = 1$ and let $\mathbf{r} = (r_1, \dots, r_n)$, $\mathbf{s} = (s_1, \dots, s_m)$ be such that

$$s_j \geq 0, \quad r_i \geq 0, \quad s_1 + \dots + s_m = 1, \quad r_1 + \dots + r_n = 1.$$

Then for any $Q > 1$ there exist $q_1, \dots, q_n, p_1, \dots, p_m \in \mathbb{Z}$ such that

$$|q_1 \alpha_{i,1} + \dots + q_n \alpha_{i,n} - p_i| < c Q^{-s_i} \quad (1 \leq i \leq m)$$

$$|q_j| \leq Q^{r_j} \quad (1 \leq j \leq n)$$

$$(q_1, \dots, q_n) \neq \mathbf{0}.$$

Weighted Badly approximable systems/matrices: Let $\alpha_{i,j} \in \mathbb{R}$, where $1 \leq j \leq n$, $1 \leq i \leq m$, and $Q > 1$. Then $A = (\alpha_{i,j})_{i,j}$ is (\mathbf{r}, \mathbf{s}) -badly approximable if there exist $c > 0$ such that for all $Q > 1$ the only integer solution $(q_1, \dots, q_n, p_1, \dots, p_m)$ to the above system is zero.

Let $\mathbf{Bad}(\mathbf{r}, \mathbf{s})$ be the set of (\mathbf{r}, \mathbf{s}) -badly approximable $m \times n$ matrices.

$$\mathbf{Bad}(n, m) = \mathbf{Bad}\left(\left(\frac{1}{n}, \dots, \frac{1}{n}\right), \left(\frac{1}{m}, \dots, \frac{1}{m}\right)\right).$$

The transpose of $\mathbf{Bad}(\mathbf{r}, \mathbf{s})$ is $\mathbf{Bad}(\mathbf{s}, \mathbf{r})$.

The transpose of $\mathbf{Bad}(n, m)$ equals $\mathbf{Bad}(m, n)$.

Multiplicative Diophantine approximation

Littlewood's conjecture (LC), (1930): For any $\alpha_1, \alpha_2 \in \mathbb{R}$ and any $\varepsilon > 0$ there exists $q \in \mathbb{Z}_{\neq 0}$, $p_1, p_2 \in \mathbb{Z}$ such that

$$|q| \cdot |q\alpha_1 - p_1| \cdot |q\alpha_2 - p_2| < \varepsilon.$$

Multiplicative Diophantine approximation

Littlewood's conjecture (LC), (1930): For any $\alpha_1, \alpha_2 \in \mathbb{R}$ and any $\varepsilon > 0$ there exists $q \in \mathbb{Z}_{\neq 0}$, $p_1, p_2 \in \mathbb{Z}$ such that

$$|q| \cdot |q\alpha_1 - p_1| \cdot |q\alpha_2 - p_2| < \varepsilon.$$

What's known:



Multiplicative Diophantine approximation

Littlewood's conjecture (LC), (1930): For any $\alpha_1, \alpha_2 \in \mathbb{R}$ and any $\varepsilon > 0$ there exists $q \in \mathbb{Z}_{\neq 0}$, $p_1, p_2 \in \mathbb{Z}$ such that

$$|q| \cdot |q\alpha_1 - p_1| \cdot |q\alpha_2 - p_2| < \varepsilon.$$

What's known:

- LC holds if either α_1 or α_2 is not badly approximable.
-

Littlewood's conjecture (LC), (1930): For any $\alpha_1, \alpha_2 \in \mathbb{R}$ and any $\varepsilon > 0$ there exists $q \in \mathbb{Z}_{\neq 0}$, $p_1, p_2 \in \mathbb{Z}$ such that

$$|q| \cdot |q\alpha_1 - p_1| \cdot |q\alpha_2 - p_2| < \varepsilon.$$

What's known:

- LC holds if either α_1 or α_2 is not badly approximable.
- LC holds if α_1 and α_2 lie in the same cubic field (Cassels & Swinnerton-Dyer, 1955)
-

Littlewood's conjecture (LC), (1930): For any $\alpha_1, \alpha_2 \in \mathbb{R}$ and any $\varepsilon > 0$ there exists $q \in \mathbb{Z}_{\neq 0}$, $p_1, p_2 \in \mathbb{Z}$ such that

$$|q| \cdot |q\alpha_1 - p_1| \cdot |q\alpha_2 - p_2| < \varepsilon.$$

What's known:

- LC holds if either α_1 or α_2 is not badly approximable.
- LC holds if α_1 and α_2 lie in the same cubic field (Cassels & Swinnerton-Dyer, 1955)
- The set of counterexample to LC has zero dimension (Einsiedler, Katok and Lindenstrauss, 2006), led Lindenstrauss to win a Fields medal in 2010
-

Littlewood's conjecture (LC), (1930): For any $\alpha_1, \alpha_2 \in \mathbb{R}$ and any $\varepsilon > 0$ there exists $q \in \mathbb{Z}_{\neq 0}$, $p_1, p_2 \in \mathbb{Z}$ such that

$$|q| \cdot |q\alpha_1 - p_1| \cdot |q\alpha_2 - p_2| < \varepsilon.$$

What's known:

- LC holds if either α_1 or α_2 is not badly approximable.
- LC holds if α_1 and α_2 lie in the same cubic field (Cassels & Swinnerton-Dyer, 1955)
- The set of counterexample to LC has zero dimension (Einsiedler, Katok and Lindenstrauss, 2006), led Lindenstrauss to win a Fields medal in 2010
- It is not known if LC holds for $(\sqrt{2}, \sqrt{3})$ or any other explicit examples of badly approximable pairs.

Littlewood's conjecture (LC), (1930): For any $\alpha_1, \alpha_2 \in \mathbb{R}$ and any $\varepsilon > 0$ there exists $q \in \mathbb{Z}_{\neq 0}$, $p_1, p_2 \in \mathbb{Z}$ such that

$$|q| \cdot |q\alpha_1 - p_1| \cdot |q\alpha_2 - p_2| < \varepsilon.$$

Multiplicative Diophantine approximation

Littlewood's conjecture (LC), (1930): For any $\alpha_1, \alpha_2 \in \mathbb{R}$ and any $\varepsilon > 0$ there exists $q \in \mathbb{Z}_{\neq 0}$, $p_1, p_2 \in \mathbb{Z}$ such that

$$|q| \cdot |q\alpha_1 - p_1| \cdot |q\alpha_2 - p_2| < \varepsilon.$$

Let

$$\Lambda = \begin{pmatrix} 1 & 0 & \alpha_1 \\ 0 & 1 & \alpha_2 \\ 0 & 0 & 1 \end{pmatrix} \mathbb{Z}^3 = \left\{ \begin{pmatrix} q\alpha_1 - p \\ q\alpha_2 - p \\ q \end{pmatrix} : q, p_1, p_2 \in \mathbb{Z} \right\}.$$

Littlewood's conjecture (LC), (1930): For any $\alpha_1, \alpha_2 \in \mathbb{R}$ and any $\varepsilon > 0$ there exists $q \in \mathbb{Z}_{\neq 0}$, $p_1, p_2 \in \mathbb{Z}$ such that

$$|q| \cdot |q\alpha_1 - p_1| \cdot |q\alpha_2 - p_2| < \varepsilon.$$

Let

$$\Lambda = \begin{pmatrix} 1 & 0 & \alpha_1 \\ 0 & 1 & \alpha_2 \\ 0 & 0 & 1 \end{pmatrix} \mathbb{Z}^3 = \left\{ \begin{pmatrix} q\alpha_1 - p \\ q\alpha_2 - p \\ q \end{pmatrix} : q, p_1, p_2 \in \mathbb{Z} \right\}.$$

LC:

$$\inf_{(\lambda_1, \lambda_2, \lambda_3) \in \Lambda, \lambda_3 \neq 0} |\lambda_1 \lambda_2 \lambda_3| = 0.$$

Multiplicative Diophantine approximation

Littlewood's conjecture (LC), (1930): For any $\alpha_1, \alpha_2 \in \mathbb{R}$ and any $\varepsilon > 0$ there exists $q \in \mathbb{Z}_{\neq 0}$, $p_1, p_2 \in \mathbb{Z}$ such that

$$|q| \cdot |q\alpha_1 - p_1| \cdot |q\alpha_2 - p_2| < \varepsilon.$$

Let

$$\Lambda = \begin{pmatrix} 1 & 0 & \alpha_1 \\ 0 & 1 & \alpha_2 \\ 0 & 0 & 1 \end{pmatrix} \mathbb{Z}^3 = \left\{ \begin{pmatrix} q\alpha_1 - p \\ q\alpha_2 - p \\ q \end{pmatrix} : q, p_1, p_2 \in \mathbb{Z} \right\}.$$

LC:

$$\inf_{(\lambda_1, \lambda_2, \lambda_3) \in \Lambda, \lambda_3 \neq 0} |\lambda_1 \lambda_2 \lambda_3| = 0.$$

Admissible lattices: $\Lambda \in \mathcal{L}_n := \mathrm{GL}(n, \mathbb{R}) / \mathrm{SL}(n, \mathbb{Z})$ is admissible if

$$\mathrm{Nm}(\Lambda) := \inf\{|\lambda_1 \cdots \lambda_n| : (\lambda_1, \dots, \lambda_n) \in \Lambda_{\neq 0}\} > 0.$$

Littlewood's conjecture (LC), (1930): For any $\alpha_1, \alpha_2 \in \mathbb{R}$ and any $\varepsilon > 0$ there exists $q \in \mathbb{Z}_{\neq 0}$, $p_1, p_2 \in \mathbb{Z}$ such that

$$|q| \cdot |q\alpha_1 - p_1| \cdot |q\alpha_2 - p_2| < \varepsilon.$$

Let

$$\Lambda = \begin{pmatrix} 1 & 0 & \alpha_1 \\ 0 & 1 & \alpha_2 \\ 0 & 0 & 1 \end{pmatrix} \mathbb{Z}^3 = \left\{ \begin{pmatrix} q\alpha_1 - p \\ q\alpha_2 - p \\ q \end{pmatrix} : q, p_1, p_2 \in \mathbb{Z} \right\}.$$

LC:

$$\inf_{(\lambda_1, \lambda_2, \lambda_3) \in \Lambda, \lambda_3 \neq 0} |\lambda_1 \lambda_2 \lambda_3| = 0.$$

Admissible lattices: $\Lambda \in \mathcal{L}_n := \mathrm{GL}(n, \mathbb{R}) / \mathrm{SL}(n, \mathbb{Z})$ is admissible if

$$\mathrm{Nm}(\Lambda) := \inf\{|\lambda_1 \cdots \lambda_n| : (\lambda_1, \dots, \lambda_n) \in \Lambda_{\neq \mathbf{0}}\} > 0.$$

Fact: The set of admissible lattices is dense in \mathcal{L}_n .

Multiplicative Diophantine approximation

Littlewood's conjecture (LC), (1930): For any $\alpha_1, \alpha_2 \in \mathbb{R}$ and any $\varepsilon > 0$ there exists $q \in \mathbb{Z}_{\neq 0}$, $p_1, p_2 \in \mathbb{Z}$ such that

$$|q| \cdot |q\alpha_1 - p_1| \cdot |q\alpha_2 - p_2| < \varepsilon.$$

Let

$$\Lambda = \begin{pmatrix} 1 & 0 & \alpha_1 \\ 0 & 1 & \alpha_2 \\ 0 & 0 & 1 \end{pmatrix} \mathbb{Z}^3 = \left\{ \begin{pmatrix} q\alpha_1 - p \\ q\alpha_2 - p \\ q \end{pmatrix} : q, p_1, p_2 \in \mathbb{Z} \right\}.$$

LC:

$$\inf_{(\lambda_1, \lambda_2, \lambda_3) \in \Lambda, \lambda_3 \neq 0} |\lambda_1 \lambda_2 \lambda_3| = 0.$$

Admissible lattices: $\Lambda \in \mathcal{L}_n := \mathrm{GL}(n, \mathbb{R}) / \mathrm{SL}(n, \mathbb{Z})$ is admissible if

$$\mathrm{Nm}(\Lambda) := \inf\{|\lambda_1 \cdots \lambda_n| : (\lambda_1, \dots, \lambda_n) \in \Lambda_{\neq \mathbf{0}}\} > 0.$$

Fact: The set of admissible lattices is dense in \mathcal{L}_n .

$n = 2$:

Multiplicative Diophantine approximation

Littlewood's conjecture (LC), (1930): For any $\alpha_1, \alpha_2 \in \mathbb{R}$ and any $\varepsilon > 0$ there exists $q \in \mathbb{Z}_{\neq 0}$, $p_1, p_2 \in \mathbb{Z}$ such that

$$|q| \cdot |q\alpha_1 - p_1| \cdot |q\alpha_2 - p_2| < \varepsilon.$$

Let

$$\Lambda = \begin{pmatrix} 1 & 0 & \alpha_1 \\ 0 & 1 & \alpha_2 \\ 0 & 0 & 1 \end{pmatrix} \mathbb{Z}^3 = \left\{ \begin{pmatrix} q\alpha_1 - p \\ q\alpha_2 - p \\ q \end{pmatrix} : q, p_1, p_2 \in \mathbb{Z} \right\}.$$

LC:

$$\inf_{(\lambda_1, \lambda_2, \lambda_3) \in \Lambda, \lambda_3 \neq 0} |\lambda_1 \lambda_2 \lambda_3| = 0.$$

Admissible lattices: $\Lambda \in \mathcal{L}_n := \mathrm{GL}(n, \mathbb{R}) / \mathrm{SL}(n, \mathbb{Z})$ is admissible if

$$\mathrm{Nm}(\Lambda) := \inf\{|\lambda_1 \cdots \lambda_n| : (\lambda_1, \dots, \lambda_n) \in \Lambda_{\neq \mathbf{0}}\} > 0.$$

Fact: The set of admissible lattices is dense in \mathcal{L}_n .

$n = 2$:

$$\Lambda = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mathbb{Z}^2 \text{ is admissible}$$

Littlewood's conjecture (LC), (1930): For any $\alpha_1, \alpha_2 \in \mathbb{R}$ and any $\varepsilon > 0$ there exists $q \in \mathbb{Z}_{\neq 0}$, $p_1, p_2 \in \mathbb{Z}$ such that

$$|q| \cdot |q\alpha_1 - p_1| \cdot |q\alpha_2 - p_2| < \varepsilon.$$

Let

$$\Lambda = \begin{pmatrix} 1 & 0 & \alpha_1 \\ 0 & 1 & \alpha_2 \\ 0 & 0 & 1 \end{pmatrix} \mathbb{Z}^3 = \left\{ \begin{pmatrix} q\alpha_1 - p \\ q\alpha_2 - p \\ q \end{pmatrix} : q, p_1, p_2 \in \mathbb{Z} \right\}.$$

LC:

$$\inf_{(\lambda_1, \lambda_2, \lambda_3) \in \Lambda, \lambda_3 \neq 0} |\lambda_1 \lambda_2 \lambda_3| = 0.$$

Admissible lattices: $\Lambda \in \mathcal{L}_n := \mathrm{GL}(n, \mathbb{R}) / \mathrm{SL}(n, \mathbb{Z})$ is admissible if

$$\mathrm{Nm}(\Lambda) := \inf\{|\lambda_1 \cdots \lambda_n| : (\lambda_1, \dots, \lambda_n) \in \Lambda_{\neq \mathbf{0}}\} > 0.$$

Fact: The set of admissible lattices is dense in \mathcal{L}_n .

$n = 2$:

$$\Lambda = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mathbb{Z}^2 \text{ is admissible} \iff$$

Multiplicative Diophantine approximation

Littlewood's conjecture (LC), (1930): For any $\alpha_1, \alpha_2 \in \mathbb{R}$ and any $\varepsilon > 0$ there exists $q \in \mathbb{Z}_{\neq 0}$, $p_1, p_2 \in \mathbb{Z}$ such that

$$|q| \cdot |q\alpha_1 - p_1| \cdot |q\alpha_2 - p_2| < \varepsilon.$$

Let

$$\Lambda = \begin{pmatrix} 1 & 0 & \alpha_1 \\ 0 & 1 & \alpha_2 \\ 0 & 0 & 1 \end{pmatrix} \mathbb{Z}^3 = \left\{ \begin{pmatrix} q\alpha_1 - p \\ q\alpha_2 - p \\ q \end{pmatrix} : q, p_1, p_2 \in \mathbb{Z} \right\}.$$

LC:

$$\inf_{(\lambda_1, \lambda_2, \lambda_3) \in \Lambda, \lambda_3 \neq 0} |\lambda_1 \lambda_2 \lambda_3| = 0.$$

Admissible lattices: $\Lambda \in \mathcal{L}_n := \mathrm{GL}(n, \mathbb{R}) / \mathrm{SL}(n, \mathbb{Z})$ is admissible if

$$\mathrm{Nm}(\Lambda) := \inf\{|\lambda_1 \cdots \lambda_n| : (\lambda_1, \dots, \lambda_n) \in \Lambda_{\neq \mathbf{0}}\} > 0.$$

Fact: The set of admissible lattices is dense in \mathcal{L}_n .

$n = 2$:

$$\Lambda = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mathbb{Z}^2 \text{ is admissible} \iff \frac{a}{b}, \frac{c}{d} \in \mathbf{Bad} = \mathbf{Bad}(1, 1) \subset \mathbb{R}.$$

Admissible lattices: examples

Admissible lattices: $\Lambda \in \mathcal{L}_n := \mathrm{GL}(n, \mathbb{R})/\mathrm{SL}(n, \mathbb{Z})$ is admissible if

$$\mathrm{Nm}(\Lambda) := \inf\{|\lambda_1 \cdots \lambda_n| : (\lambda_1, \dots, \lambda_n) \in \Lambda_{\neq \mathbf{0}}\} > 0.$$

$n = 2$:

$$\Lambda = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mathbb{Z}^2 \text{ is admissible} \iff \frac{a}{b}, \frac{c}{d} \in \mathbf{Bad} = \mathbf{Bad}(1, 1) \subset \mathbb{R}.$$

Example (admissible lattices in \mathcal{L}_n for $n \geq 3$):

Admissible lattices: examples

Admissible lattices: $\Lambda \in \mathcal{L}_n := \mathrm{GL}(n, \mathbb{R})/\mathrm{SL}(n, \mathbb{Z})$ is admissible if

$$\mathrm{Nm}(\Lambda) := \inf\{|\lambda_1 \cdots \lambda_n| : (\lambda_1, \dots, \lambda_n) \in \Lambda_{\neq \mathbf{0}}\} > 0.$$

$n = 2$:

$$\Lambda = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mathbb{Z}^2 \text{ is admissible} \iff \frac{a}{b}, \frac{c}{d} \in \mathbf{Bad} = \mathbf{Bad}(1, 1) \subset \mathbb{R}.$$

Example (admissible lattices in \mathcal{L}_n for $n \geq 3$): Let $f \in \mathbb{Z}[x]$, monic, irreducible over \mathbb{Q} , $\deg f = n$ with all real roots. Thus

$$f(x) = (x - \alpha_1) \cdots (x - \alpha_n) \text{ with } \alpha_i \in \mathbb{R}.$$

Admissible lattices: $\Lambda \in \mathcal{L}_n := \mathrm{GL}(n, \mathbb{R})/\mathrm{SL}(n, \mathbb{Z})$ is admissible if

$$\mathrm{Nm}(\Lambda) := \inf\{|\lambda_1 \cdots \lambda_n| : (\lambda_1, \dots, \lambda_n) \in \Lambda_{\neq \mathbf{0}}\} > 0.$$

$n = 2$:

$$\Lambda = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mathbb{Z}^2 \text{ is admissible} \iff \frac{a}{b}, \frac{c}{d} \in \mathbf{Bad} = \mathbf{Bad}(1, 1) \subset \mathbb{R}.$$

Example (admissible lattices in \mathcal{L}_n for $n \geq 3$): Let $f \in \mathbb{Z}[x]$, monic, irreducible over \mathbb{Q} , $\deg f = n$ with all real roots. Thus

$f(x) = (x - \alpha_1) \cdots (x - \alpha_n)$ with $\alpha_i \in \mathbb{R}$. Define

$$\Lambda = \begin{pmatrix} 1 & \alpha_1 & \cdots & \alpha_1^{n-1} \\ \vdots & & \ddots & \vdots \\ 1 & \alpha_n & \cdots & \alpha_n^{n-1} \end{pmatrix} \mathbb{Z}^n.$$

Admissible lattices: $\Lambda \in \mathcal{L}_n := \mathrm{GL}(n, \mathbb{R})/\mathrm{SL}(n, \mathbb{Z})$ is admissible if

$$\mathrm{Nm}(\Lambda) := \inf\{|\lambda_1 \cdots \lambda_n| : (\lambda_1, \dots, \lambda_n) \in \Lambda_{\neq 0}\} > 0.$$

$n = 2$:

$$\Lambda = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mathbb{Z}^2 \text{ is admissible} \iff \frac{a}{b}, \frac{c}{d} \in \mathbf{Bad} = \mathbf{Bad}(1, 1) \subset \mathbb{R}.$$

Example (admissible lattices in \mathcal{L}_n for $n \geq 3$): Let $f \in \mathbb{Z}[x]$, monic, irreducible over \mathbb{Q} , $\deg f = n$ with all real roots. Thus

$f(x) = (x - \alpha_1) \cdots (x - \alpha_n)$ with $\alpha_i \in \mathbb{R}$. Define

$$\Lambda = \begin{pmatrix} 1 & \alpha_1 & \cdots & \alpha_1^{n-1} \\ \vdots & & \ddots & \vdots \\ 1 & \alpha_n & \cdots & \alpha_n^{n-1} \end{pmatrix} \mathbb{Z}^n.$$

$$\begin{aligned} \mathrm{Nm}(\Lambda) &= \inf \left\{ \prod_{1 \leq i \leq n} |g(\alpha_i)| : g \in \mathbb{Z}[x]_{\neq 0}, \deg g \leq n-1 \right\} \\ &= \inf \left\{ |\mathrm{Resultant}(f, g)| : g \in \mathbb{Z}[x]_{\neq 0}, \deg g \leq n-1 \right\} \geq 1. \end{aligned}$$

Admissible lattices and **Bad**

Admissible lattices and **Bad**

Let g be an $(n+1) \times (n+1)$ matrix such that $\Lambda = g\mathbb{Z}^{n+1}$ is admissible.
Let (x_0, x_1, \dots, x_n) be any row (column) of g .

Admissible lattices and **Bad**

Let g be an $(n+1) \times (n+1)$ matrix such that $\Lambda = g\mathbb{Z}^{n+1}$ is admissible. Let (x_0, x_1, \dots, x_n) be any row (column) of g . Then $x_0 \neq 0$ and

$$\left(\frac{x_1}{x_0}, \dots, \frac{x_n}{x_0}\right) \in \mathbf{Bad}(n, 1).$$

Proof.

Admissible lattices and **Bad**

Let g be an $(n+1) \times (n+1)$ matrix such that $\Lambda = g\mathbb{Z}^{n+1}$ is admissible. Let (x_0, x_1, \dots, x_n) be any row (column) of g . Then $x_0 \neq 0$ and

$$\left(\frac{x_1}{x_0}, \dots, \frac{x_n}{x_0}\right) \in \mathbf{Bad}(n, 1).$$

Proof. WLOG $x_0 = 1$ and g is the first column.

Admissible lattices and **Bad**

Let g be an $(n+1) \times (n+1)$ matrix such that $\Lambda = g\mathbb{Z}^{n+1}$ is admissible. Let (x_0, x_1, \dots, x_n) be any row (column) of g . Then $x_0 \neq 0$ and

$$\left(\frac{x_1}{x_0}, \dots, \frac{x_n}{x_0}\right) \in \mathbf{Bad}(n, 1).$$

Proof. WLOG $x_0 = 1$ and g is the first column. Let $Q > 1$ and $(q_1, \dots, q_n, p) \in \mathbb{Z}_{\neq 0}^{n+1}$ with $|q_i| \leq Q$.

Admissible lattices and **Bad**

Let g be an $(n+1) \times (n+1)$ matrix such that $\Lambda = g\mathbb{Z}^{n+1}$ is admissible. Let (x_0, x_1, \dots, x_n) be any row (column) of g . Then $x_0 \neq 0$ and

$$\left(\frac{x_1}{x_0}, \dots, \frac{x_n}{x_0}\right) \in \mathbf{Bad}(n, 1).$$

Proof. WLOG $x_0 = 1$ and g is the first column. Let $Q > 1$ and $(q_1, \dots, q_n, p) \in \mathbb{Z}_{\neq 0}^{n+1}$ with $|q_i| \leq Q$. Recall,

$$0 < \mathrm{Nm}(\Lambda) \leq |\lambda_0 \cdots \lambda_n|, \quad (\lambda_1, \dots, \lambda_n) \in \Lambda_{\neq 0}.$$

Admissible lattices and **Bad**

Let g be an $(n+1) \times (n+1)$ matrix such that $\Lambda = g\mathbb{Z}^{n+1}$ is admissible. Let (x_0, x_1, \dots, x_n) be any row (column) of g . Then $x_0 \neq 0$ and

$$\left(\frac{x_1}{x_0}, \dots, \frac{x_n}{x_0}\right) \in \mathbf{Bad}(n, 1).$$

Proof. WLOG $x_0 = 1$ and g is the first column. Let $Q > 1$ and $(q_1, \dots, q_n, p) \in \mathbb{Z}_{\neq 0}^{n+1}$ with $|q_i| \leq Q$. Recall,

$$0 < \mathrm{Nm}(\Lambda) \leq |\lambda_0 \cdots \lambda_n|, \quad (\lambda_1, \dots, \lambda_n) \in \Lambda_{\neq 0}.$$

Let $(\lambda_0, \dots, \lambda_n)^T = g(-p, q_1, \dots, q_n)^T \neq \mathbf{0}$.

Admissible lattices and **Bad**

Let g be an $(n+1) \times (n+1)$ matrix such that $\Lambda = g\mathbb{Z}^{n+1}$ is admissible. Let (x_0, x_1, \dots, x_n) be any row (column) of g . Then $x_0 \neq 0$ and

$$\left(\frac{x_1}{x_0}, \dots, \frac{x_n}{x_0}\right) \in \mathbf{Bad}(n, 1).$$

Proof. WLOG $x_0 = 1$ and g is the first column. Let $Q > 1$ and $(q_1, \dots, q_n, p) \in \mathbb{Z}_{\neq 0}^{n+1}$ with $|q_i| \leq Q$. Recall,

$$0 < \mathrm{Nm}(\Lambda) \leq |\lambda_0 \cdots \lambda_n|, \quad (\lambda_1, \dots, \lambda_n) \in \Lambda_{\neq 0}.$$

Let $(\lambda_0, \dots, \lambda_n)^T = g(-p, q_1, \dots, q_n)^T \neq \mathbf{0}$. Note $|\lambda_i| \leq C_g Q$.

Admissible lattices and **Bad**

Let g be an $(n+1) \times (n+1)$ matrix such that $\Lambda = g\mathbb{Z}^{n+1}$ is admissible. Let (x_0, x_1, \dots, x_n) be any row (column) of g . Then $x_0 \neq 0$ and

$$\left(\frac{x_1}{x_0}, \dots, \frac{x_n}{x_0}\right) \in \mathbf{Bad}(n, 1).$$

Proof. WLOG $x_0 = 1$ and g is the first column. Let $Q > 1$ and $(q_1, \dots, q_n, p) \in \mathbb{Z}_{\neq 0}^{n+1}$ with $|q_i| \leq Q$. Recall,

$$0 < \mathrm{Nm}(\Lambda) \leq |\lambda_0 \cdots \lambda_n|, \quad (\lambda_1, \dots, \lambda_n) \in \Lambda_{\neq 0}.$$

Let $(\lambda_0, \dots, \lambda_n)^T = g(-p, q_1, \dots, q_n)^T \neq \mathbf{0}$. Note $|\lambda_i| \leq C_g Q$. Then

$$|\lambda_0| =$$

Admissible lattices and **Bad**

Let g be an $(n+1) \times (n+1)$ matrix such that $\Lambda = g\mathbb{Z}^{n+1}$ is admissible. Let (x_0, x_1, \dots, x_n) be any row (column) of g . Then $x_0 \neq 0$ and

$$\left(\frac{x_1}{x_0}, \dots, \frac{x_n}{x_0}\right) \in \mathbf{Bad}(n, 1).$$

Proof. WLOG $x_0 = 1$ and g is the first column. Let $Q > 1$ and $(q_1, \dots, q_n, p) \in \mathbb{Z}_{\neq 0}^{n+1}$ with $|q_i| \leq Q$. Recall,

$$0 < \mathrm{Nm}(\Lambda) \leq |\lambda_0 \cdots \lambda_n|, \quad (\lambda_1, \dots, \lambda_n) \in \Lambda_{\neq 0}.$$

Let $(\lambda_0, \dots, \lambda_n)^T = g(-p, q_1, \dots, q_n)^T \neq \mathbf{0}$. Note $|\lambda_i| \leq C_g Q$. Then

$$|\lambda_0| = |q_1 x_1 + \cdots + q_n x_n - p| \geq$$

Admissible lattices and **Bad**

Let g be an $(n+1) \times (n+1)$ matrix such that $\Lambda = g\mathbb{Z}^{n+1}$ is admissible. Let (x_0, x_1, \dots, x_n) be any row (column) of g . Then $x_0 \neq 0$ and

$$\left(\frac{x_1}{x_0}, \dots, \frac{x_n}{x_0}\right) \in \mathbf{Bad}(n, 1).$$

Proof. WLOG $x_0 = 1$ and g is the first column. Let $Q > 1$ and $(q_1, \dots, q_n, p) \in \mathbb{Z}_{\neq 0}^{n+1}$ with $|q_i| \leq Q$. Recall,

$$0 < \mathrm{Nm}(\Lambda) \leq |\lambda_0 \cdots \lambda_n|, \quad (\lambda_1, \dots, \lambda_n) \in \Lambda_{\neq 0}.$$

Let $(\lambda_0, \dots, \lambda_n)^T = g(-p, q_1, \dots, q_n)^T \neq \mathbf{0}$. Note $|\lambda_i| \leq C_g Q$. Then

$$|\lambda_0| = |q_1 x_1 + \cdots + q_n x_n - p| \geq \frac{|\mathrm{Nm}(\Lambda)|}{|\lambda_1 \cdots \lambda_n|} \geq$$

Admissible lattices and **Bad**

Let g be an $(n+1) \times (n+1)$ matrix such that $\Lambda = g\mathbb{Z}^{n+1}$ is admissible. Let (x_0, x_1, \dots, x_n) be any row (column) of g . Then $x_0 \neq 0$ and

$$\left(\frac{x_1}{x_0}, \dots, \frac{x_n}{x_0}\right) \in \mathbf{Bad}(n, 1).$$

Proof. WLOG $x_0 = 1$ and g is the first column. Let $Q > 1$ and $(q_1, \dots, q_n, p) \in \mathbb{Z}_{\neq 0}^{n+1}$ with $|q_i| \leq Q$. Recall,

$$0 < \mathrm{Nm}(\Lambda) \leq |\lambda_0 \cdots \lambda_n|, \quad (\lambda_1, \dots, \lambda_n) \in \Lambda_{\neq 0}.$$

Let $(\lambda_0, \dots, \lambda_n)^T = g(-p, q_1, \dots, q_n)^T \neq \mathbf{0}$. Note $|\lambda_i| \leq C_g Q$. Then

$$|\lambda_0| = |q_1 x_1 + \cdots + q_n x_n - p| \geq \frac{|\mathrm{Nm}(\Lambda)|}{|\lambda_1 \cdots \lambda_n|} \geq \frac{|\mathrm{Nm}(\Lambda)|}{C_g^n Q^n}.$$

Admissible lattices and **Bad**

Let g be an $(n+1) \times (n+1)$ matrix such that $\Lambda = g\mathbb{Z}^{n+1}$ is admissible. Let (x_0, x_1, \dots, x_n) be any row (column) of g . Then $x_0 \neq 0$ and

$$\left(\frac{x_1}{x_0}, \dots, \frac{x_n}{x_0}\right) \in \mathbf{Bad}(n, 1).$$

Proof. WLOG $x_0 = 1$ and g is the first column. Let $Q > 1$ and $(q_1, \dots, q_n, p) \in \mathbb{Z}_{\neq 0}^{n+1}$ with $|q_i| \leq Q$. Recall,

$$0 < \mathrm{Nm}(\Lambda) \leq |\lambda_0 \cdots \lambda_n|, \quad (\lambda_1, \dots, \lambda_n) \in \Lambda_{\neq 0}.$$

Let $(\lambda_0, \dots, \lambda_n)^T = g(-p, q_1, \dots, q_n)^T \neq \mathbf{0}$. Note $|\lambda_i| \leq C_g Q$. Then

$$|\lambda_0| = |q_1 x_1 + \cdots + q_n x_n - p| \geq \frac{|\mathrm{Nm}(\Lambda)|}{|\lambda_1 \cdots \lambda_n|} \geq \frac{|\mathrm{Nm}(\Lambda)|}{C_g^n Q^n}.$$

Similarly one can give examples of matrices in $\mathbf{Bad}(n, m)$.

Admissible lattices and **Bad**

Let g be an $(n+1) \times (n+1)$ matrix such that $\Lambda = g\mathbb{Z}^{n+1}$ is admissible. Let (x_0, x_1, \dots, x_n) be any row (column) of g . Then $x_0 \neq 0$ and

$$\left(\frac{x_1}{x_0}, \dots, \frac{x_n}{x_0}\right) \in \mathbf{Bad}(n, 1).$$

Proof. WLOG $x_0 = 1$ and g is the first column. Let $Q > 1$ and $(q_1, \dots, q_n, p) \in \mathbb{Z}_{\neq 0}^{n+1}$ with $|q_i| \leq Q$. Recall,

$$0 < \mathrm{Nm}(\Lambda) \leq |\lambda_0 \cdots \lambda_n|, \quad (\lambda_1, \dots, \lambda_n) \in \Lambda_{\neq 0}.$$

Let $(\lambda_0, \dots, \lambda_n)^T = g(-p, q_1, \dots, q_n)^T \neq \mathbf{0}$. Note $|\lambda_i| \leq C_g Q$. Then

$$|\lambda_0| = |q_1 x_1 + \cdots + q_n x_n - p| \geq \frac{|\mathrm{Nm}(\Lambda)|}{|\lambda_1 \cdots \lambda_n|} \geq \frac{|\mathrm{Nm}(\Lambda)|}{C_g^n Q^n}.$$

Similarly one can give examples of matrices in **Bad**(n, m).

The proof does not extend to weighted badly approximable points.

Q: How likely is that a random matrix lies in **Bad**(n, m)?

Answer:

Q: How likely is that a random matrix lies in **Bad**(n, m)?

Answer: Unlikely! The set **Bad**(n, m) has Lebesgue measure 0.

Q: How likely is that a random matrix lies in **Bad**(n, m)?

Answer: Unlikely! The set **Bad**(n, m) has Lebesgue measure 0.

However: **Bad**(n, m) has Hausdorff dimension nm , the same as the dimension of the ambient space. (Schmidt, 1960s)

More on **Bad**

Q: How likely is that a random matrix lies in **Bad**(n, m)?

Answer: Unlikely! The set **Bad**(n, m) has Lebesgue measure 0.

However: **Bad**(n, m) has Hausdorff dimension nm , the same as the dimension of the ambient space. (Schmidt, 1960s)

Bad := **Bad**(1, 1) can be characterise using continued fractions:

Q: How likely is that a random matrix lies in **Bad**(n, m)?

Answer: Unlikely! The set **Bad**(n, m) has Lebesgue measure 0.

However: **Bad**(n, m) has Hausdorff dimension nm , the same as the dimension of the ambient space. (Schmidt, 1960s)

Bad := **Bad**(1, 1) can be characterise using continued fractions:

$$x = a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{\ddots}}}} = [a_0; a_1, a_2, \dots],$$

where all $a_n \in \mathbb{Z}$, $a_n \geq 1$ for $n = 1, 2, \dots$.

Q: How likely is that a random matrix lies in **Bad**(n, m)?

Answer: Unlikely! The set **Bad**(n, m) has Lebesgue measure 0.

However: **Bad**(n, m) has Hausdorff dimension nm , the same as the dimension of the ambient space. (Schmidt, 1960s)

Bad := **Bad**(1, 1) can be characterise using continued fractions:

$$x = a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{\ddots}}}$$

where all $a_n \in \mathbb{Z}$, $a_n \geq 1$ for $n = 1, 2, \dots$. The convergents

$$\frac{p_n}{q_n} := [a_1, a_2, a_3, \dots, a_n]$$

are best approximations and satisfy

$$\frac{1}{(a_{n+1} + 2)q_n^2} < \left| x - \frac{p_n}{q_n} \right| < \frac{1}{a_{n+1}q_n^2}.$$

Q: How likely is that a random matrix lies in **Bad**(n, m)?

Answer: Unlikely! The set **Bad**(n, m) has Lebesgue measure 0.

However: **Bad**(n, m) has Hausdorff dimension nm , the same as the dimension of the ambient space. (Schmidt, 1960s)

Bad := **Bad**(1, 1) can be characterise using continued fractions:

$$x = a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{\ddots}}}$$

where all $a_n \in \mathbb{Z}$, $a_n \geq 1$ for $n = 1, 2, \dots$. The convergents

$$\frac{p_n}{q_n} := [a_1, a_2, a_3, \dots, a_n]$$

are best approximations and satisfy

$$\frac{1}{(a_{n+1} + 2)q_n^2} < \left| x - \frac{p_n}{q_n} \right| < \frac{1}{a_{n+1}q_n^2}.$$

Fact: An irrational $x = [a_0; a_1, a_2, \dots]$ is in **Bad** $\Leftrightarrow \exists M \forall i \in \mathbb{N} a_i \leq M$

More on dimension 1

Hurwitz's Theorem (1891): $\forall x \in \mathbb{R} \setminus \mathbb{Q}$ there are infinitely many coprime integers p and $q > 0$ such that

$$\left| x - \frac{p}{q} \right| < \frac{1}{\sqrt{5}q^2}.$$

Hurwitz's Theorem (1891): $\forall x \in \mathbb{R} \setminus \mathbb{Q}$ there are infinitely many coprime integers p and $q > 0$ such that

$$\left| x - \frac{p}{q} \right| < \frac{1}{\sqrt{5}q^2}.$$

The constant $1/(\sqrt{5})$ is best possible.

Recall:

$$\mathbf{Bad} := \{x \in \mathbb{R} : \exists c(x) > 0 \text{ s.t. } \left| x - \frac{p}{q} \right| \geq \frac{c(x)}{q^2} \quad \forall q \in \mathbb{N}, p \in \mathbb{Z}\}.$$

- quadratic irrationals are in **Bad**

Hurwitz's Theorem (1891): $\forall x \in \mathbb{R} \setminus \mathbb{Q}$ there are infinitely many coprime integers p and $q > 0$ such that

$$\left| x - \frac{p}{q} \right| < \frac{1}{\sqrt{5}q^2}.$$

The constant $1/(\sqrt{5})$ is best possible.

Recall:

$$\mathbf{Bad} := \{x \in \mathbb{R} : \exists c(x) > 0 \text{ s.t. } \left| x - \frac{p}{q} \right| \geq \frac{c(x)}{q^2} \quad \forall q \in \mathbb{N}, p \in \mathbb{Z}\}.$$

- quadratic irrationals are in **Bad**
- Folklore Conjecture:** Cubic irrationals are not in **Bad**

Hurwitz's Theorem (1891): $\forall x \in \mathbb{R} \setminus \mathbb{Q}$ there are infinitely many coprime integers p and $q > 0$ such that

$$\left| x - \frac{p}{q} \right| < \frac{1}{\sqrt{5}q^2}.$$

The constant $1/(\sqrt{5})$ is best possible.

Recall:

$$\mathbf{Bad} := \{x \in \mathbb{R} : \exists c(x) > 0 \text{ s.t. } \left| x - \frac{p}{q} \right| \geq \frac{c(x)}{q^2} \quad \forall q \in \mathbb{N}, p \in \mathbb{Z}\}.$$

- quadratic irrationals are in **Bad**
- Folklore Conjecture:** Cubic irrationals are not in **Bad**
- However, for any irrational algebraic x and any $\varepsilon > 0$ there exists a constant $c(x, \varepsilon) > 0$ such that

$$\text{(Roth, 1955)} \quad \left| x - \frac{p}{q} \right| \geq \frac{c(x, \varepsilon)}{q^{2+\varepsilon}}.$$

More on dimension 1

Badly approximable numbers seem like worst approximable.

Badly approximable numbers seem like worst approximable. However, they are the only irrationals that Dirichlet's theorem can be improved for!!!

Dirichlet's Theorem (1848): *Let $x \in \mathbb{R} \setminus \mathbb{Q}$ and $c = 1$. Let $Q > 1$. Then there are integers p and q such that*

$$\left| x - \frac{p}{q} \right| < \frac{c}{qQ}, \quad 1 \leq q \leq Q. \quad (3)$$

Badly approximable numbers seem like worst approximable. However, they are the only irrationals that Dirichlet's theorem can be improved for!!!

Dirichlet's Theorem (1848): *Let $x \in \mathbb{R} \setminus \mathbb{Q}$ and $c = 1$. Let $Q > 1$. Then there are integers p and q such that*

$$\left| x - \frac{p}{q} \right| < \frac{c}{qQ}, \quad 1 \leq q \leq Q. \quad (3)$$

Definition: x is called Dirichlet Improvable if there exists a positive $c < 1$ such that for all sufficiently large Q there are integers p and q satisfying (3).

Badly approximable numbers seem like worst approximable. However, they are the only irrationals that Dirichlet's theorem can be improved for!!!

Dirichlet's Theorem (1848): *Let $x \in \mathbb{R} \setminus \mathbb{Q}$ and $c = 1$. Let $Q > 1$. Then there are integers p and q such that*

$$\left| x - \frac{p}{q} \right| < \frac{c}{qQ}, \quad 1 \leq q \leq Q. \quad (3)$$

Definition: x is called Dirichlet Improvable if there exists a positive $c < 1$ such that for all sufficiently large Q there are integers p and q satisfying (3).

Theorem (Davenport and Schmidt (1970)): An irrational x is Dirichlet improvable if and only if $x \in \mathbf{Bad}$.

Badly approximable numbers seem like worst approximable. However, they are the only irrationals that Dirichlet's theorem can be improved for!!!

Dirichlet's Theorem (1848): *Let $x \in \mathbb{R} \setminus \mathbb{Q}$ and $c = 1$. Let $Q > 1$. Then there are integers p and q such that*

$$\left| x - \frac{p}{q} \right| < \frac{c}{qQ}, \quad 1 \leq q \leq Q. \quad (3)$$

Definition: x is called Dirichlet Improvable if there exists a positive $c < 1$ such that for all sufficiently large Q there are integers p and q satisfying (3).

Theorem (Davenport and Schmidt (1970)): An irrational x is Dirichlet improvable if and only if $x \in \mathbf{Bad}$.

Remark. Dirichlet Improvable points/matrices can be introduced in higher dimensions, but their characterisation is not that simple. It is known that almost every matrix is not Dirichlet Improvable.

More on dimension 1: the Three Distance Theorem

To understand the gaps between $q\alpha - p$, where $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ is fixed and $p, q \in \mathbb{Z}$, $0 \leq q \leq Q$ it is enough to describe the distribution of

$$\{\alpha\}, \{2\alpha\}, \dots, \{Q\alpha\} \quad (4)$$

where $\{\cdot\}$ denotes the fractional part.

More on dimension 1: the Three Distance Theorem

To understand the gaps between $q\alpha - p$, where $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ is fixed and $p, q \in \mathbb{Z}$, $0 \leq q \leq Q$ it is enough to describe the distribution of

$$\{\alpha\}, \{2\alpha\}, \dots, \{Q\alpha\} \quad (4)$$

where $\{\cdot\}$ denotes the fractional part.

The Three Distance Theorem: For any $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and any integer $Q \geq 1$ the points (4) partition $[0, 1]$ into $Q + 1$ intervals which lengths take at most 3 different values δ_A , δ_B and δ_C with $\delta_C = \delta_A + \delta_B$.

More on dimension 1: the Three Distance Theorem

To understand the gaps between $q\alpha - p$, where $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ is fixed and $p, q \in \mathbb{Z}$, $0 \leq q \leq Q$ it is enough to describe the distribution of

$$\{\alpha\}, \{2\alpha\}, \dots, \{Q\alpha\} \quad (4)$$

where $\{\cdot\}$ denotes the fractional part.

The Three Distance Theorem: For any $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and any integer $Q \geq 1$ the points (4) partition $[0, 1]$ into $Q + 1$ intervals which lengths take at most 3 different values δ_A , δ_B and δ_C with $\delta_C = \delta_A + \delta_B$.

The length of the intervals, the number of intervals of each type and even the order in which the intervals of various type emerge can be determined using continued fractions!!!

More on dimension 1: the Three Distance Theorem

Theorem: Let $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and $[a_0; a_1, a_2, \dots]$ be the continued fraction expansion of α and $p_k/q_k = [a_0; a_1, \dots, a_k]$ and $D_k = q_k\alpha - p_k$ ($k \geq 0$). Then for any $Q \in \mathbb{N}$ there exists a unique integer $k \geq 0$ such that

$$q_k + q_{k-1} \leq Q < q_{k+1} + q_k \quad (5)$$

and unique integers r and s satisfying

$$Q = rq_k + q_{k-1} + s, \quad 1 \leq r \leq a_{k+1} \quad \text{and} \quad 0 \leq s \leq q_k - 1 \quad (6)$$

such that the points

$$\{\alpha\}, \{2\alpha\}, \dots, \{Q\alpha\}$$

partition $[0, 1]$ into $Q + 1$ intervals, of which

$$\begin{aligned} N_A &= Q + 1 - q_k && \text{are of length } \delta_A = |D_k|, \\ N_B &= s + 1 && \text{are of length } \delta_B = |D_{k+1}| + (a_{k+1} - r)|D_k|, \\ N_C &= q_k - s - 1 && \text{are of length } \delta_C = \delta_A + \delta_B. \end{aligned}$$

More on dimension 1: Metric viewpoint

More on dimension 1: Metric viewpoint

Let $\psi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ and let $W(\psi)$ be the set of $x \in [0, 1]$ such that

$$|qx - p| < \psi(q) \tag{7}$$

for infinitely many coprime $(p, q) \in \mathbb{Z} \times \mathbb{N}$.

More on dimension 1: Metric viewpoint

Let $\psi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ and let $W(\psi)$ be the set of $x \in [0, 1]$ such that

$$|qx - p| < \psi(q) \quad (7)$$

for infinitely many coprime $(p, q) \in \mathbb{Z} \times \mathbb{N}$.

Khinchine's Theorem (1924) *If ψ is monotonic, then*

$$m(W(\psi)) = \begin{cases} 0 & \text{if } \sum_{q=1}^{\infty} \psi(q) < \infty, \\ 1 & \text{if } \sum_{q=1}^{\infty} \psi(q) = \infty. \end{cases}$$

More on dimension 1: Metric viewpoint

Let $\psi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ and let $W(\psi)$ be the set of $x \in [0, 1]$ such that

$$|qx - p| < \psi(q) \quad (7)$$

for infinitely many coprime $(p, q) \in \mathbb{Z} \times \mathbb{N}$.

Khintchine's Theorem (1924) *If ψ is monotonic, then*

$$m(W(\psi)) = \begin{cases} 0 & \text{if } \sum_{q=1}^{\infty} \psi(q) < \infty, \\ 1 & \text{if } \sum_{q=1}^{\infty} \psi(q) = \infty. \end{cases}$$

Example. Take $\psi(q) = q^{-1}(\log q)^{-1-\varepsilon}$ with $\varepsilon > 0$.

More on dimension 1: Metric viewpoint

Let $\psi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ and let $W(\psi)$ be the set of $x \in [0, 1]$ such that

$$|qx - p| < \psi(q) \quad (7)$$

for infinitely many coprime $(p, q) \in \mathbb{Z} \times \mathbb{N}$.

Khintchine's Theorem (1924) *If ψ is monotonic, then*

$$m(W(\psi)) = \begin{cases} 0 & \text{if } \sum_{q=1}^{\infty} \psi(q) < \infty, \\ 1 & \text{if } \sum_{q=1}^{\infty} \psi(q) = \infty. \end{cases}$$

Example. Take $\psi(q) = q^{-1}(\log q)^{-1-\varepsilon}$ with $\varepsilon > 0$. The Khintchine sum converges. Hence for x picked at random (7) has only a finite number of solution. Also means that for almost every x there exists a constant $c(x, \psi) > 0$ such that

$$|qx - p| \geq \frac{c(x, \psi)}{(q \log q)^{1+\varepsilon}}$$

for all $(p, q) \in \mathbb{Z} \times \mathbb{N}$.

The Khintchine-Groshev theorem

The Khintchine-Groshev theorem

Let $\psi : \mathbb{R}_{>0} \rightarrow \mathbb{R}_{>0}$ and $W(n, m; \psi)$ denote the set of real matrices $X = (x_{i,j})_{\substack{1 \leq j \leq n \\ 1 \leq i \leq m}}$ with $0 \leq x_{i,j} \leq 1$ such that

$$|q_1 \alpha_{i,1} + \cdots + q_n \alpha_{i,n} - p_i| < \psi(Q) \quad (1 \leq i \leq m)$$
$$1 \leq \max_{1 \leq j \leq n} |q_j| \leq Q$$

holds for infinitely many $(q_1, \dots, q_n, p_1, \dots, p_m) \in \mathbb{Z}^{n+m}$.

The Khintchine-Groshev theorem

Let $\psi : \mathbb{R}_{>0} \rightarrow \mathbb{R}_{>0}$ and $W(n, m; \psi)$ denote the set of real matrices $X = (x_{i,j})_{\substack{1 \leq j \leq n \\ 1 \leq i \leq m}}$ with $0 \leq x_{i,j} \leq 1$ such that

$$|q_1 \alpha_{i,1} + \cdots + q_n \alpha_{i,n} - p_i| < \psi(Q) \quad (1 \leq i \leq m)$$
$$1 \leq \max_{1 \leq j \leq n} |q_j| \leq Q$$

holds for infinitely many $(q_1, \dots, q_n, p_1, \dots, p_m) \in \mathbb{Z}^{n+m}$.

Recall that Dirichlet's theorem $\Rightarrow W(n, m)(\psi)$ is everything if $\psi(Q) = Q^{-n/m}$.

The Khintchine-Groshev theorem

Let $\psi : \mathbb{R}_{>0} \rightarrow \mathbb{R}_{>0}$ and $W(n, m; \psi)$ denote the set of real matrices $X = (x_{i,j})_{\substack{1 \leq j \leq n \\ 1 \leq i \leq m}}$ with $0 \leq x_{i,j} \leq 1$ such that

$$|q_1 \alpha_{i,1} + \cdots + q_n \alpha_{i,n} - p_i| < \psi(Q) \quad (1 \leq i \leq m)$$
$$1 \leq \max_{1 \leq j \leq n} |q_j| \leq Q$$

holds for infinitely many $(q_1, \dots, q_n, p_1, \dots, p_m) \in \mathbb{Z}^{n+m}$.

Recall that Dirichlet's theorem $\Rightarrow W(n, m)(\psi)$ is everything if $\psi(Q) = Q^{-n/m}$.

Theorem (Khintchine-Groshev /1924-1938/): Suppose that ψ is monotonic. Then

$$\mathbf{Prob}(X \in W(n, m; \psi)) = \begin{cases} 0, & \text{if } \sum_{q=1}^{\infty} q^{n-1} \psi(q)^m < \infty, \\ 1, & \text{if } \sum_{q=1}^{\infty} q^{n-1} \psi(q)^m = \infty. \end{cases}$$

The Khintchine-Groshev theorem for manifolds

The Khintchine-Groshev theorem for manifolds

Q: What can be said about the probability of $W(n, m; \psi)$ under the condition that the entries of the matrix X are functionally related?

The Khintchine-Groshev theorem for manifolds

Q: What can be said about the probability of $W(n, m; \psi)$ under the condition that the entries of the matrix X are functionally related?

Functional relations \Rightarrow that X lies on a submanifold (curve or surface).

The Khintchine-Groshev theorem for manifolds

Q: What can be said about the probability of $W(n, m; \psi)$ under the condition that the entries of the matrix X are functionally related?

Functional relations \Rightarrow that X lies on a submanifold (curve or surface).

We'll restrict to the case of one linear form: $m = 1$.

The Khintchine-Groshev theorem for manifolds

Q: What can be said about the probability of $W(n, m; \psi)$ under the condition that the entries of the matrix X are functionally related?

Functional relations \Rightarrow that X lies on a submanifold (curve or surface).

We'll restrict to the case of one linear form: $m = 1$.

Also restrict ourselves to the case when the manifold is analytic, e.g. is given by polynomial relations. The coordinates of $X = (x_1, \dots, x_n)$ are analytic functions of several (possibly just one) other variables. Example: $X = (x, x^2, x^3, \dots, x^n)$.

The Khintchine-Groshev theorem for manifolds

Q: What can be said about the probability of $W(n, m; \psi)$ under the condition that the entries of the matrix X are functionally related?

Functional relations \Rightarrow that X lies on a submanifold (curve or surface).

We'll restrict to the case of one linear form: $m = 1$.

Also restrict ourselves to the case when the manifold is analytic, e.g. is given by polynomial relations. The coordinates of $X = (x_1, \dots, x_n)$ are analytic functions of several (possibly just one) other variables. Example: $X = (x, x^2, x^3, \dots, x^n)$.

Non-degeneracy: A connected analytic submanifold \mathcal{M} of \mathbb{R}^n is non-degenerate if it is not entirely contained in a hyperplane of \mathbb{R}^n .

The Khintchine-Groshev theorem for manifolds

Q: What can be said about the probability of $W(n, m; \psi)$ under the condition that the entries of the matrix X are functionally related?

Functional relations \Rightarrow that X lies on a submanifold (curve or surface).

We'll restrict to the case of one linear form: $m = 1$.

Also restrict ourselves to the case when the manifold is analytic, e.g. is given by polynomial relations. The coordinates of $X = (x_1, \dots, x_n)$ are analytic functions of several (possibly just one) other variables. Example: $X = (x, x^2, x^3, \dots, x^n)$.

Non-degeneracy: A connected analytic submanifold \mathcal{M} of \mathbb{R}^n is non-degenerate if it is not entirely contained in a hyperplane of \mathbb{R}^n .

The Khintchine-Groshev theorem for non-degenerate manifolds

(BBKM, 1998-2002): Suppose that ψ is monotonic and $\mathcal{M} \subset \mathbb{R}^n$ is non-degenerate. Then

$$\mathbf{Prob}(X \in W(n, 1; \psi) | X \in \mathcal{M}) = \begin{cases} 0, & \text{if } \sum_{q=1}^{\infty} q^{n-1} \psi(q) < \infty, \\ 1, & \text{if } \sum_{q=1}^{\infty} q^{n-1} \psi(q) = \infty. \end{cases}$$

Extremal manifolds and homogeneous flows

Extremal manifolds and homogeneous flows

Consider $\psi_\tau(q) = q^{-\tau}$

Extremal manifolds and homogeneous flows

Consider $\psi_\tau(q) = q^{-\tau}$

Extremal matrices: X is extremal if $X \notin W(n, m; \psi_\tau)$ for any $\tau > n/m$.
If X is not extremal it is called very well approximable (VWA).

Extremal manifolds and homogeneous flows

Consider $\psi_\tau(q) = q^{-\tau}$

Extremal matrices: X is extremal if $X \notin W(n, m; \psi_\tau)$ for any $\tau > n/m$.
If X is not extremal it is called very well approximable (VWA).

Theorem of Kleinbock and Margulis, 1998): Almost every point on any non-degenerate manifold \mathcal{M} in \mathbb{R}^n is extremal.

Extremal manifolds and homogeneous flows

Consider $\psi_\tau(q) = q^{-\tau}$

Extremal matrices: X is extremal if $X \notin W(n, m; \psi_\tau)$ for any $\tau > n/m$.
If X is not extremal it is called very well approximable (VWA).

Theorem of Kleinbock and Margulis, 1998): Almost every point on any non-degenerate manifold \mathcal{M} in \mathbb{R}^n is extremal.

Relation to homogeneous flows: Let $g_t = \text{diag}\{e^t, e^{-t/n}, \dots, e^{-t/n}\}$ and

$$\Lambda_X = \begin{pmatrix} 1 & x_1 & \dots & x_n \\ 0 & 1 & \dots & 0 \\ 0 & & \ddots & \\ 0 & \dots & 0 & 1 \end{pmatrix} \mathbb{Z}^{n+1}$$

Extremal manifolds and homogeneous flows

Consider $\psi_\tau(q) = q^{-\tau}$

Extremal matrices: X is extremal if $X \notin W(n, m; \psi_\tau)$ for any $\tau > n/m$.
If X is not extremal it is called very well approximable (VWA).

Theorem of Kleinbock and Margulis, 1998): Almost every point on any non-degenerate manifold \mathcal{M} in \mathbb{R}^n is extremal.

Relation to homogeneous flows: Let $g_t = \text{diag}\{e^t, e^{-t/n}, \dots, e^{-t/n}\}$ and

$$\Lambda_X = \begin{pmatrix} 1 & x_1 & \dots & x_n \\ 0 & 1 & \dots & 0 \\ 0 & & \ddots & \\ 0 & \dots & 0 & 1 \end{pmatrix} \mathbb{Z}^{n+1}$$

Let

$$\delta(\Lambda) = \inf\{\|\lambda\| : \lambda \in \Lambda_{\neq 0}\}.$$

Extremal manifolds and homogeneous flows

Consider $\psi_\tau(q) = q^{-\tau}$

Extremal matrices: X is extremal if $X \notin W(n, m; \psi_\tau)$ for any $\tau > n/m$.
If X is not extremal it is called very well approximable (VWA).

Theorem of Kleinbock and Margulis, 1998): Almost every point on any non-degenerate manifold \mathcal{M} in \mathbb{R}^n is extremal.

Relation to homogeneous flows: Let $g_t = \text{diag}\{e^t, e^{-t/n}, \dots, e^{-t/n}\}$ and

$$\Lambda_X = \begin{pmatrix} 1 & x_1 & \dots & x_n \\ 0 & 1 & \dots & 0 \\ 0 & & \ddots & \\ 0 & \dots & 0 & 1 \end{pmatrix} \mathbb{Z}^{n+1}$$

Let

$$\delta(\Lambda) = \inf\{\|\lambda\| : \lambda \in \Lambda_{\neq 0}\}.$$

Fact: X is VWA if and only if there exists $\varepsilon > 0$ such that

$$\delta(g_t \Lambda_X) \leq e^{-\varepsilon t} \quad \text{for arbitrarily large } t.$$