

```

CROSSTABS
/TABLES=diagPca BY g894NOS3
/FORMAT=AVALUE TABLES
/STATISTICS=CHISQ RISK CMH(1)
/CELLS=COUNT
/COUNT ROUND CELL.

```

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:47:19
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=diagPca BY g894NOS3 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.016
	Elapsed Time	0:00:00.016
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for diagPca * g894NOS3, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
diagPca * g894NOS3	300	75,0%	100	25,0%	400	100,0%

diagPca * g894NOS3 Crosstabulation

Count

		g894NOS3			Total
		GG	GT	TT	
diagPca	no	78	59	13	150
	yes	76	65	9	150
	Total	154	124	22	300

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1,044 ^a	2	,593
Likelihood Ratio	1,048	2	,592
Linear-by-Linear Association	,397	1	,528
N of Valid Cases	300		

a. 0 cells (,0%) have expected count less than 5. The minimum expected count is 11,00.

Risk Estimate

	Value
Odds Ratio for diagPca (no / yes)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=cmDiagPca0Kont BY g894NOS3

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Output Created	22-lip-2012 11:47:19
Comments	
Input Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmv\rez\SPSS\Stat.sav
Active Dataset	DataSet1
Filter	<none>
Weight	<none>
Split File	<none>
N of Rows in Working Data File	400

Notes

Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=cmDiagPca0Kont BY g894NOS3 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
	Resources	
	Processor Time	0:00:00.031
	Elapsed Time	0:00:00.150
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmb
dmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmDiagPca0Kont * g894NOS3, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmDiagPca0Kont * g894NOS3	250	62,5%	150	37,5%	400	100,0%

cmDiagPca0Kont * g894NOS3 Crosstabulation

Count

		g894NOS3			Total
		GG	GT	TT	
cmDiagPca0Kont	no	78	59	13	150
	control	54	40	6	100
	Total	132	99	19	250

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	,614 ^a	2	,736
Likelihood Ratio	,630	2	,730
Linear-by-Linear Association	,566	1	,452
N of Valid Cases	250		

a. 0 cells (,0%) have expected count less than 5. The minimum expected count is 7,60.

Risk Estimate

	Value
Odds Ratio for cmDiagPca0Kont (no / control)	a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

```
CROSSTABS
/TABLES=cmDiagPca1Kont BY g894NOS3
/FORMAT=AVALUE TABLES
/STATISTICS=CHISQ RISK CMH(1)
/CELLS=COUNT
/COUNT ROUND CELL.
```

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:47:19
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=cmDiagPca1Kont BY g894NOS3 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.016
	Elapsed Time	0:00:00.017
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmDiagPca1Kont * g894NOS3, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmDiagPca1Kont * g894NOS3	250	62,5%	150	37,5%	400	100,0%

cmDiagPca1Kont * g894NOS3 Crosstabulation

Count

		g894NOS3			Total
		GG	GT	TT	
cmDiagPca1Kont	yes	76	65	9	150
	control	54	40	6	100
	Total	130	105	15	250

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	,287 ^a	2	,866
Likelihood Ratio	,287	2	,866
Linear-by-Linear Association	,047	1	,828
N of Valid Cases	250		

a. 0 cells (,0%) have expected count less than 5. The minimum expected count is 6,00.

Risk Estimate

	Value
Odds Ratio for cmDiagPca1Kont (yes / control)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=kontrol BY g894NOS3

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:47:20
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=kontrol BY g894NOS3 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.015
	Elapsed Time	0:00:00.380
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for kontrol * g894NOS3, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
kontrol * g894NOS3	400	100,0%	0	,0%	400	100,0%

kontrol * g894NOS3 Crosstabulation

Count

		g894NOS3			Total
		GG	GT	TT	
kontrol	no kontrol	154	124	22	300
	kontrol	54	40	6	100
	Total	208	164	28	400

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	,326 ^a	2	,850
Likelihood Ratio	,331	2	,847
Linear-by-Linear Association	,303	1	,582
N of Valid Cases	400		

a. 0 cells (,0%) have expected count less than 5. The minimum expected count is 7,00.

Risk Estimate

	Value
Odds Ratio for kontrol (no kontrol / kontrol)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=cmTStadOnly12 BY g894NOS3

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:47:20
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvfrez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=cmTStadOnly12 BY g894NOS3 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.

Notes

Resources	Processor Time	0:00:00.000
	Elapsed Time	0:00:00.014
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmb
dmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmTstadOnly12 * g894NOS3, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmTstadOnly12 * g894NOS3	100	25,0%	300	75,0%	400	100,0%

cmTstadOnly12 * g894NOS3 Crosstabulation

Count

		g894NOS3			Total
		GG	GT	TT	
cmTstadOnly12	T1	13	13	2	28
	T2	33	32	7	72
	Total	46	45	9	100

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	,168 ^a	2	,919
Likelihood Ratio	,175	2	,916
Linear-by-Linear Association	,126	1	,723
N of Valid Cases	100		

a. 1 cells (16,7%) have expected count less than 5. The minimum expected count is 2,52.

Risk Estimate

	Value
Odds Ratio for cmTstadOnly12 (T1 / T2)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.


```

/TABLES=cmTStadOnly13 BY g894NOS3
/FORMAT=AVALUE TABLES
/STATISTICS=CHISQ RISK CMH(1)
/CELLS=COUNT
/COUNT ROUND CELL.

```

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:47:20
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
Resources	Syntax	CROSSTABS /TABLES=cmTStadOnly13 BY g894NOS3 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
	Processor Time	0:00:00.031
	Elapsed Time	0:00:00.135
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmTStadOnly13 * g894NOS3, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

cmTStadOnly13 * g894NOS3 Crosstabulation

Count		g894NOS3			Total
		GG	GT	TT	
cmTStadOnly13	T1	13	13	2	28

cmTStadOnly13 * g894NOS3 Crosstabulation

Count

		g894NOS3			Total
		GG	GT	TT	
cmTStadOnly13	T3,T4	30	20	0	50
	Total	43	33	2	78

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	4,346 ^a	2	,114
Likelihood Ratio	4,887	2	,087
Linear-by-Linear Association	4,140	1	,042
N of Valid Cases	78		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,72.

Risk Estimate

	Value
Odds Ratio for cmTStadOnly13 (T1 / T3, T4)	a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=cmTStadOnly23 BY g894NOS3

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Output Created	22-lip-2012 11:47:21
Comments	
Input Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmv\rez\SPSS\Stat.sav
Active Dataset	DataSet1
Filter	<none>
Weight	<none>
Split File	<none>
N of Rows in Working Data File	400

Notes

Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=cmTstadOnly23 BY g894NOS3 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
	Resources	
	Processor Time	0:00:00.031
	Elapsed Time	0:00:00.021
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmb
dmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmTstadOnly23 * g894NOS3, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmTstadOnly23 * g894NOS3	122	30,5%	278	69,5%	400	100,0%

cmTstadOnly23 * g894NOS3 Crosstabulation

Count

		g894NOS3			Total
		GG	GT	TT	
cmTstadOnly23	T2	33	32	7	72
	T3,T4	30	20	0	50
	Total	63	52	7	122

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	6,145 ^a	2	,046
Likelihood Ratio	8,652	2	,013
Linear-by-Linear Association	6,093	1	,014
N of Valid Cases	122		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is 2,87.

Risk Estimate

	Value
Odds Ratio for cmTStadOnly23 (T2 / T3, T4)	a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=cmPsaLT10vs10to20FonPCA1 BY g894NOS3

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:47:21
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmvdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=cmPsaLT10vs10to20FonPCA1 BY g894NOS3 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.015
	Elapsed Time	0:00:00.015
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmvdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmPsaLT10vs10to20FonPCA1 * g894NOS3, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmPsaLT10vs10to20FonPCA1 * g894NOS3	88	22,0%	312	78,0%	400	100,0%

cmPsaLT10vs10to20FonPCA1 * g894NOS3 Crosstabulation

Count

		g894NOS3			Total
		GG	GT	TT	
cmPsaLT10vs10to20FonPCA1	<10	19	25	5	49
	10-20	19	18	2	39
	Total	38	43	7	88

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1,306 ^a	2	,521
Likelihood Ratio	1,334	2	,513
Linear-by-Linear Association	1,161	1	,281
N of Valid Cases	88		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is 3,10.

Risk Estimate

	Value
Odds Ratio for cmPsaLT10vs10to20FonPCA1 (<10 / 10-20)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=cmPsaLT10vsGT20FonPCA1 BY g894NOS3

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:47:22
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=cmPsaLT10vsGT20FonPCA1 BY g894NOS3 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.016
	Elapsed Time	0:00:00.077
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmPsaLT10vsGT20FonPCA1 * g894NOS3, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmPsaLT10vsGT20FonPCA1 * g894NOS3	111	27,8%	289	72,3%	400	100,0%

cmPsaLT10vsGT20FonPCA1 * g894NOS3 Crosstabulation

Count

		g894NOS3			Total
		GG	GT	TT	
cmPsaLT10vsGT20FonPCA1	<10	19	25	5	49
	>20	38	22	2	62
	Total	57	47	7	111

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	6,375 ^a	2	,041
Likelihood Ratio	6,450	2	,040
Linear-by-Linear Association	4,743	1	,029
N of Valid Cases	111		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is 3,09.

Risk Estimate

	Value
Odds Ratio for cmPsaLT10vsGT20Fon PCA1 (<10 / >20)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=cmPsa10to20vsGT20FonPCA1 BY g894NOS3

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:47:22
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvfrez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
Missing Value Handling	Split File	<none>
	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.

Notes

Syntax		CROSSTABS /TABLES=cmPsa10to20vsGT20Fon PCA1 BY g894NOS3 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.	
Resources	Processor Time	0:00:00.015	
	Elapsed Time	0:00:00.014	
	Dimensions Requested		2
	Cells Available		174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmb
dmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmPsa10to20vsGT20FonPCA1 * g894NOS3, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmPsa10to20vsGT20Fon PCA1 * g894NOS3	101	25,3%	299	74,8%	400	100,0%

cmPsa10to20vsGT20FonPCA1 * g894NOS3 Crosstabulation

Count

		g894NOS3			Total
		GG	GT	TT	
cmPsa10to20vsGT20Fon PCA1	10-20	19	18	2	39
	>20	38	22	2	62
	Total	57	40	4	101

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1,578 ^a	2	,454
Likelihood Ratio	1,573	2	,455
Linear-by-Linear Association	,934	1	,334
N of Valid Cases	101		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is 1,54.

Risk Estimate

	Value
Odds Ratio for cmPsa10to20vsGT20Fon PCA1 (10-20 / >20)	a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=cmPsaLT20vsGT20onPCA1 BY g894NOS3

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:47:22
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmvdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=cmPsaLT20vsGT20onPCA1 BY g894NOS3 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.046
	Elapsed Time	0:00:00.267
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmvdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmPsaLT20vsGT20onPCA1 * g894NOS3, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmPsaLT20vsGT20on PCA1 * g894NOS3	150	37,5%	250	62,5%	400	100,0%

cmPsaLT20vsGT20onPCA1 * g894NOS3 Crosstabulation

Count

		g894NOS3			Total
		GG	GT	TT	
cmPsaLT20vsGT20on PCA1	,00	38	43	7	88
	<10	38	22	2	62
	Total	76	65	9	150

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	5,212 ^a	2	,074
Likelihood Ratio	5,320	2	,070
Linear-by-Linear Association	3,549	1	,060
N of Valid Cases	150		

a. 1 cells (16,7%) have expected count less than 5. The minimum expected count is 3,72.

Risk Estimate

	Value
Odds Ratio for cmPsaLT20vsGT20on PCA1 (,00 / <10)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=cmGgLtvsGt7F BY g894NOS3

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:47:22
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=cmGgLtvsGt7F BY g894NOS3 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.015
	Elapsed Time	0:00:00.240
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmGgLtvsGt7F * g894NOS3, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmGgLtvsGt7F * g894NOS3	93	23,3%	307	76,8%	400	100,0%

cmGgLtvsGt7F * g894NOS3 Crosstabulation

Count		g894NOS3			Total
		GG	GT	TT	
cmGgLtvsGt7F	<7	33	34	4	71
	>7	13	9	0	22
	Total	46	43	4	93

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1,956 ^a	2	,376
Likelihood Ratio	2,860	2	,239
Linear-by-Linear Association	1,903	1	,168
N of Valid Cases	93		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,95.

Risk Estimate

	Value
Odds Ratio for cmGgLtvsGt7F (<7 / >7)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=cmGgLt7vsEq7F BY g894NOS3

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:47:23
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvfrez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=cmGgLt7vsEq7F BY g894NOS3 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.

Notes

Resources	Processor Time	0:00:00.032
	Elapsed Time	0:00:00.024
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmb
dmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmGgLt7vsEq7F * g894NOS3, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmGgLt7vsEq7F * g894NOS3	128	32,0%	272	68,0%	400	100,0%

cmGgLt7vsEq7F * g894NOS3 Crosstabulation

Count

		g894NOS3			Total
		GG	GT	TT	
cmGgLt7vsEq7F	<7	33	34	4	71
	=7	30	22	5	57
	Total	63	56	9	128

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1,310 ^a	2	,519
Likelihood Ratio	1,311	2	,519
Linear-by-Linear Association	,082	1	,775
N of Valid Cases	128		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is 4,01.

Risk Estimate

	Value
Odds Ratio for cmGgLt7vsEq7F (<7 / =7)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

```

/TABLES=cmGgEq7vsGt7F BY g894NOS3
/FORMAT=AVALUE TABLES
/STATISTICS=CHISQ RISK CMH(1)
/CELLS=COUNT
/COUNT ROUND CELL.

```

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:47:23
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmvdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
Resources	Syntax	CROSSTABS /TABLES=cmGgEq7vsGt7F BY g894NOS3 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
	Processor Time	0:00:00.031
	Elapsed Time	0:00:00.104
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmvdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmGgEq7vsGt7F * g894NOS3, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

cmGgEq7vsGt7F * g894NOS3 Crosstabulation

Count		g894NOS3			Total
		GG	GT	TT	
cmGgEq7vsGt7F	=7	30	22	5	57

cmGgEq7vsGt7F * g894NOS3 Crosstabulation

Count

		g894NOS3			Total
		GG	GT	TT	
cmGgEq7vsGt7F	>7	13	9	0	22
	Total	43	31	5	79

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	2,073 ^a	2	,355
Likelihood Ratio	3,405	2	,182
Linear-by-Linear Association	1,838	1	,175
N of Valid Cases	79		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is 1,39.

Risk Estimate

	Value
Odds Ratio for cmGgEq7vsGt7F (=7 / >7)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=mMeta BY g894NOS3

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Output Created	22-lip-2012 11:47:23
Comments	
Input Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
Active Dataset	DataSet1
Filter	<none>
Weight	<none>
Split File	<none>
N of Rows in Working Data File	400

Notes

Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=mMeta BY g894NOS3 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.016
	Elapsed Time	0:00:00.016
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmb
dmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for mMeta * g894NOS3, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
mMeta * g894NOS3	150	37,5%	250	62,5%	400	100,0%

mMeta * g894NOS3 Crosstabulation

Count

		g894NOS3			Total
		GG	GT	TT	
mMeta	no	45	42	8	95
	yes	31	23	1	55
	Total	76	65	9	150

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	3,133 ^a	2	,209
Likelihood Ratio	3,630	2	,163
Linear-by-Linear Association	3,108	1	,078
N of Valid Cases	150		

a. 1 cells (16,7%) have expected count less than 5. The minimum expected count is 3,30.

Risk Estimate

	Value
Odds Ratio for mMeta (no / yes)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=mRiskEAU BY g894NOS3

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:47:24
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=mRiskEAU BY g894NOS3 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.015
	Elapsed Time	0:00:00.138
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for mRiskEAU * g894NOS3, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
mRiskEAU * g894NOS3	150	37,5%	250	62,5%	400	100,0%

mRiskEAU * g894NOS3 Crosstabulation

Count

		g894NOS3			Total
		GG	GT	TT	
mRiskEAU	low	5	8	1	14
	medium	25	25	5	55
	high	46	32	3	81
	Total	76	65	9	150

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	4,077 ^a	4	,396
Likelihood Ratio	4,079	4	,395
Linear-by-Linear Association	2,505	1	,113
N of Valid Cases	150		

a. 3 cells (33,3%) have expected count less than 5. The minimum expected count is ,84.

Risk Estimate

	Value
Odds Ratio for mRiskEAU (low / medium)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=mRiskEAULowMedium BY g894NOS3

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:47:24
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=mRiskEAULowMedium BY g894NOS3 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.016
	Elapsed Time	0:00:00.015
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for mRiskEAULowMedium * g894NOS3, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
mRiskEAULowMedium * g894NOS3	69	17,3%	331	82,8%	400	100,0%

mRiskEAULowMedium * g894NOS3 Crosstabulation

Count		g894NOS3			Total
		GG	GT	TT	
mRiskEAULowMedium	low	5	8	1	14
	medium	25	25	5	55
	Total	30	33	6	69

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	,611 ^a	2	,737
Likelihood Ratio	,611	2	,737
Linear-by-Linear Association	,002	1	,962
N of Valid Cases	69		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is 1,22.

Risk Estimate

	Value
Odds Ratio for mRiskEAULowMedium (low / medium)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=mRiskEAULowHigh BY g894NOS3

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:47:24
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvfrez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.

Notes

Syntax	CROSSTABS /TABLES=mRiskEAULowHigh BY g894NOS3 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.		
Resources	Processor Time	0:00:00.016	
	Elapsed Time	0:00:00.017	
	Dimensions Requested		2
	Cells Available		174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmb
dmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for mRiskEAULowHigh * g894NOS3, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
mRiskEAULowHigh * g894NOS3	95	23,8%	305	76,3%	400	100,0%

mRiskEAULowHigh * g894NOS3 Crosstabulation

Count

		g894NOS3			Total
		GG	GT	TT	
mRiskEAULowHigh	low	5	8	1	14
	high	46	32	3	81
	Total	51	40	4	95

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	2,205 ^a	2	,332
Likelihood Ratio	2,195	2	,334
Linear-by-Linear Association	1,333	1	,248
N of Valid Cases	95		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,59.

Risk Estimate

	Value
Odds Ratio for mRiskEAULowHigh (low / high)	a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=mRiskEAUMediumHigh BY g894NOS3

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:47:25
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=mRiskEAUMediumHigh BY g894NOS3 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.016
	Elapsed Time	0:00:00.012
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for mRiskEAUMediumHigh * g894NOS3, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
mRiskEAUMediumHigh * g894NOS3	136	34,0%	264	66,0%	400	100,0%

mRiskEAUMediumHigh * g894NOS3 Crosstabulation

Count

		g894NOS3			Total
		GG	GT	TT	
mRiskEAUMediumHigh	medium	25	25	5	55
	high	46	32	3	81
	Total	71	57	8	136

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	2,699 ^a	2	,259
Likelihood Ratio	2,671	2	,263
Linear-by-Linear Association	2,546	1	,111
N of Valid Cases	136		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is 3,24.

Risk Estimate

	Value
Odds Ratio for mRiskEAUMediumHigh (medium / high)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=mRiskMed BY g894NOS3

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:47:25
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=mRiskMed BY g894NOS3 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.016
	Elapsed Time	0:00:00.012
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for mRiskMed * g894NOS3, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
mRiskMed * g894NOS3	150	37,5%	250	62,5%	400	100,0%

mRiskMed * g894NOS3 Crosstabulation

Count

		g894NOS3			Total
		GG	GT	TT	
mRiskMed	low	24	27	4	55
	high	52	38	5	95
	Total	76	65	9	150

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1,746 ^a	2	,418
Likelihood Ratio	1,748	2	,417
Linear-by-Linear Association	,908	1	,341
N of Valid Cases	150		

a. 1 cells (16,7%) have expected count less than 5. The minimum expected count is 3,30.

Risk Estimate

	Value
Odds Ratio for mRiskMed (low / high)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=mRiskMedLowMedium BY g894NOS3

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:47:25
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvfrez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=mRiskMedLowMedium BY g894NOS3 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.

Notes

Resources	Processor Time	0:00:00.016
	Elapsed Time	0:00:00.012
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmb
dmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for mRiskMedLowMedium * g894NOS3, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

No measures of association are computed for the crosstabulation of mRiskMedLowMedium * g894NOS3. At least one variable in each 2-way table upon which measures of association are computed is a constant.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
mRiskMedLowMedium * g894NOS3	55	13,8%	345	86,3%	400	100,0%

mRiskMedLowMedium * g894NOS3 Crosstabulation

Count

		g894NOS3			Total
		GG	GT	TT	
mRiskMedLowMedium	low	24	27	4	55
	Total	24	27	4	55

Chi-Square Tests

	Value
Pearson Chi-Square	a.
N of Valid Cases	55

a. No statistics are computed because mRiskMedLowMedium is a constant.

Risk Estimate

	Value
Odds Ratio for mRiskMedLowMedium (low / .)	a.

a. No statistics are computed because mRiskMedLowMedium is a constant.

CROSSTABS

/TABLES=mRiskMedLowHigh BY g894NOS3

```

/FORMAT=AVALUE TABLES
/STATISTICS=CHISQ RISK CMH(1)
/CELLS=COUNT
/COUNT ROUND CELL.

```

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:47:26
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=mRiskMedLowHigh BY g894NOS3 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.031
	Elapsed Time	0:00:00.016
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for mRiskMedLowHigh * g894NOS3, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
mRiskMedLowHigh * g894NOS3	150	37,5%	250	62,5%	400	100,0%

mRiskMedLowHigh * g894NOS3 Crosstabulation

Count

		g894NOS3			Total
		GG	GT	TT	
mRiskMedLowHigh	low	24	27	4	55
	high	52	38	5	95
	Total	76	65	9	150

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1,746 ^a	2	,418
Likelihood Ratio	1,748	2	,417
Linear-by-Linear Association	,908	1	,341
N of Valid Cases	150		

a. 1 cells (16,7%) have expected count less than 5. The minimum expected count is 3,30.

Risk Estimate

	Value
Odds Ratio for mRiskMedLowHigh (low / high)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=mRiskMedMediumHigh BY g894NOS3

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Output Created	22-lip-2012 11:47:26
Comments	
Input Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
Active Dataset	DataSet1
Filter	<none>
Weight	<none>
Split File	<none>
N of Rows in Working Data File	400

Notes

Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=mRiskMedMediumHigh BY g894NOS3 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.000
	Elapsed Time	0:00:00.014
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmb
dmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for mRiskMedMediumHigh * g894NOS3, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

No measures of association are computed for the crosstabulation of mRiskMedMediumHigh * g894NOS3. At least one variable in each 2-way table upon which measures of association are computed is a constant.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
mRiskMedMediumHigh * g894NOS3	95	23,8%	305	76,3%	400	100,0%

mRiskMedMediumHigh * g894NOS3 Crosstabulation

Count

		g894NOS3			Total
		GG	GT	TT	
mRiskMedMediumHigh	high	52	38	5	95
	Total	52	38	5	95

Chi-Square Tests

	Value
Pearson Chi-Square	a.
N of Valid Cases	95

a. No statistics are computed because mRiskMedMediumHigh is a constant.

Risk Estimate

	Value
Odds Ratio for mRiskMedMediumHigh (high / .)	a.

a. No statistics are computed
because mRiskMedMediumHigh
is a constant.