

```

CROSSTABS
/TABLES=diagPca BY g4242
/FORMAT=AVALUE TABLES
/STATISTICS=CHISQ RISK CMH(1)
/CELLS=COUNT
/COUNT ROUND CELL.

```

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:57:47
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=diagPca BY g4242 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.000
	Elapsed Time	0:00:00.017
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for diagPca * g4242, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
diagPca * g4242	300	75,0%	100	25,0%	400	100,0%

diagPca * g4242 Crosstabulation

Count

		g4242			Total
		AA	GA	GG	
diagPca	no	1	25	124	150
	yes	2	42	106	150
	Total	3	67	230	300

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	6,055 ^a	2	,048
Likelihood Ratio	6,111	2	,047
Linear-by-Linear Association	4,034	1	,045
N of Valid Cases	300		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is 1,50.

Risk Estimate

	Value
Odds Ratio for diagPca (no / yes)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=cmDiagPca0Kont BY g4242

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Output Created	22-lip-2012 11:57:47
Comments	
Input Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmv\rez\SPSS\Stat.sav
Active Dataset	DataSet1
Filter	<none>
Weight	<none>
Split File	<none>
N of Rows in Working Data File	400

Notes

Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=cmDiagPca0Kont BY g4242 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
	Resources	
	Processor Time	0:00:00.016
	Elapsed Time	0:00:00.017
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmb
dmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmDiagPca0Kont * g4242, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmDiagPca0Kont * g4242	250	62,5%	150	37,5%	400	100,0%

cmDiagPca0Kont * g4242 Crosstabulation

Count

		g4242			Total
		AA	GA	GG	
cmDiagPca0Kont	no	1	25	124	150
	control	0	20	80	100
	Total	1	45	204	250

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1,089 ^a	2	,580
Likelihood Ratio	1,440	2	,487
Linear-by-Linear Association	,000	1	1,000
N of Valid Cases	250		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,40.

Risk Estimate

	Value
Odds Ratio for cmDiagPca0Kont (no / control)	a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=cmDiagPca1Kont BY g4242

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:57:48
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=cmDiagPca1Kont BY g4242 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.000
	Elapsed Time	0:00:00.027
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmDiagPca1Kont * g4242, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmDiagPca1Kont * g4242	250	62,5%	150	37,5%	400	100,0%

cmDiagPca1Kont * g4242 Crosstabulation

Count

		g4242			Total
		AA	GA	GG	
cmDiagPca1Kont	yes	2	42	106	150
	control	0	20	80	100
	Total	2	62	186	250

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	3,584 ^a	2	,167
Likelihood Ratio	4,330	2	,115
Linear-by-Linear Association	3,521	1	,061
N of Valid Cases	250		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,80.

Risk Estimate

	Value
Odds Ratio for cmDiagPca1Kont (yes / control)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=kontrol BY g4242

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:57:48
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=kontrol BY g4242 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.031
	Elapsed Time	0:00:00.027
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for kontrol * g4242, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
kontrol * g4242	400	100,0%	0	,0%	400	100,0%

kontrol * g4242 Crosstabulation

Count

		g4242			Total
		AA	GA	GG	
kontrol	no control	3	67	230	300
	control	0	20	80	100
	Total	3	87	310	400

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1,295 ^a	2	,523
Likelihood Ratio	2,024	2	,363
Linear-by-Linear Association	1,183	1	,277
N of Valid Cases	400		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,75.

Risk Estimate

	Value
Odds Ratio for kontrol (no kontrol / kontrol)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=cmTStadOnly12 BY g4242

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:57:48
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvfrez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=cmTStadOnly12 BY g4242 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.

Notes

Resources	Processor Time	0:00:00.016
	Elapsed Time	0:00:00.122
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmb
dmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmTStadOnly12 * g4242, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmTStadOnly12 * g4242	100	25,0%	300	75,0%	400	100,0%

cmTStadOnly12 * g4242 Crosstabulation

Count

		g4242			Total
		AA	GA	GG	
cmTStadOnly12	T1	0	5	23	28
	T2	1	23	48	72
	Total	1	28	71	100

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	2,498 ^a	2	,287
Likelihood Ratio	2,882	2	,237
Linear-by-Linear Association	2,096	1	,148
N of Valid Cases	100		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,28.

Risk Estimate

	Value
Odds Ratio for cmTStadOnly12 (T1 / T2)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.


```

/TABLES=cmTStadOnly13 BY g4242
/FORMAT=AVALUE TABLES
/STATISTICS=CHISQ RISK CMH(1)
/CELLS=COUNT
/COUNT ROUND CELL.

```

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:57:48
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=cmTStadOnly13 BY g4242 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.000
	Elapsed Time	0:00:00.016
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmTStadOnly13 * g4242, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmTStadOnly13 * g4242	78	19,5%	322	80,5%	400	100,0%

cmTStadOnly13 * g4242 Crosstabulation

Count

		g4242			Total
		AA	GA	GG	
cmTStadOnly13	T1	0	5	23	28
	T3,T4	1	14	35	50
	Total	1	19	58	78

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1,674 ^a	2	,433
Likelihood Ratio	2,036	2	,361
Linear-by-Linear Association	1,531	1	,216
N of Valid Cases	78		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,36.

Risk Estimate

	Value
Odds Ratio for cmTStadOnly13 (T1 / T3, T4)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=cmTStadOnly23 BY g4242

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Output Created	22-lip-2012 11:57:49
Comments	
Input Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
Active Dataset	DataSet1
Filter	<none>
Weight	<none>
Split File	<none>
N of Rows in Working Data File	400

Notes

Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=cmTStadOnly23 BY g4242 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.015
	Elapsed Time	0:00:00.014
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmb
dmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmTStadOnly23 * g4242, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmTStadOnly23 * g4242	122	30,5%	278	69,5%	400	100,0%

cmTStadOnly23 * g4242 Crosstabulation

Count

		g4242			Total
		AA	GA	GG	
cmTStadOnly23	T2	1	23	48	72
	T3,T4	1	14	35	50
	Total	2	37	83	122

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	,267 ^a	2	,875
Likelihood Ratio	,267	2	,875
Linear-by-Linear Association	,004	1	,949
N of Valid Cases	122		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,82.

Risk Estimate

	Value
Odds Ratio for cmTStadOnly23 (T2 / T3, T4)	a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=cmPsaLT10vs10to20FonPCA1 BY g4242

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:57:49
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmvdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=cmPsaLT10vs10to20FonPCA1 BY g4242 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.016
	Elapsed Time	0:00:00.013
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmvdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmPsaLT10vs10to20FonPCA1 * g4242, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmPsaLT10vs10to20FonPCA1 * g4242	88	22,0%	312	78,0%	400	100,0%

cmPsaLT10vs10to20FonPCA1 * g4242 Crosstabulation

Count

		g4242			Total
		AA	GA	GG	
cmPsaLT10vs10to20FonPCA1	<10	0	15	34	49
	10-20	1	7	31	39
	Total	1	22	65	88

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	2,949 ^a	2	,229
Likelihood Ratio	3,363	2	,186
Linear-by-Linear Association	,000	1	,991
N of Valid Cases	88		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,44.

Risk Estimate

	Value
Odds Ratio for cmPsaLT10vs10to20FonPCA1 (<10 / 10-20)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=cmPsaLT10vsGT20FonPCA1 BY g4242

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:57:49
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=cmPsaLT10vsGT20FonPCA1 BY g4242 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.016
	Elapsed Time	0:00:00.019
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmPsaLT10vsGT20FonPCA1 * g4242, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmPsaLT10vsGT20FonPCA1 * g4242	111	27,8%	289	72,3%	400	100,0%

cmPsaLT10vsGT20FonPCA1 * g4242 Crosstabulation

Count

		g4242			Total
		AA	GA	GG	
cmPsaLT10vsGT20FonPCA1	<10	0	15	34	49
	>20	1	20	41	62
	Total	1	35	75	111

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	,857 ^a	2	,652
Likelihood Ratio	1,231	2	,540
Linear-by-Linear Association	,623	1	,430
N of Valid Cases	111		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,44.

Risk Estimate

	Value
Odds Ratio for cmPsaLT10vsGT20Fon PCA1 (<10 / >20)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=cmPsa10to20vsGT20FonPCA1 BY g4242

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:57:49
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvfrez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.

Notes

Syntax	CROSSTABS /TABLES=cmPsa10to20vsGT20Fon PCA1 BY g4242 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.		
Resources	Processor Time	0:00:00.016	
	Elapsed Time	0:00:00.226	
	Dimensions Requested		2
	Cells Available		174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmb
dmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmPsa10to20vsGT20FonPCA1 * g4242, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmPsa10to20vsGT20Fon PCA1 * g4242	101	25,3%	299	74,8%	400	100,0%

cmPsa10to20vsGT20FonPCA1 * g4242 Crosstabulation

Count

		g4242			Total
		AA	GA	GG	
cmPsa10to20vsGT20Fon PCA1	10-20	1	7	31	39
	>20	1	20	41	62
	Total	2	27	72	101

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	2,542 ^a	2	,280
Likelihood Ratio	2,636	2	,268
Linear-by-Linear Association	,344	1	,557
N of Valid Cases	101		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,77.

Risk Estimate

	Value
Odds Ratio for cmPsa10to20vsGT20Fon PCA1 (10-20 / >20)	a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=cmPsaLT20vsGT20onPCA1 BY g4242

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:57:50
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmvdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=cmPsaLT20vsGT20onPCA1 BY g4242 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.015
	Elapsed Time	0:00:00.017
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmvdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmPsaLT20vsGT20onPCA1 * g4242, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmPsaLT20vsGT20onPCA1 * g4242	150	37,5%	250	62,5%	400	100,0%

cmPsaLT20vsGT20onPCA1 * g4242 Crosstabulation

Count

		g4242			Total
		AA	GA	GG	
cmPsaLT20vsGT20onPCA1	,00	1	22	65	88
	<10	1	20	41	62
	Total	2	42	106	150

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1,054 ^a	2	,590
Likelihood Ratio	1,047	2	,592
Linear-by-Linear Association	,681	1	,409
N of Valid Cases	150		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,83.

Risk Estimate

	Value
Odds Ratio for cmPsaLT20vsGT20onPCA1 (,00 / <10)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=cmGgLtvsGt7F BY g4242

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:57:50
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=cmGgLtvsGt7F BY g4242 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.031
	Elapsed Time	0:00:00.177
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmGgLtvsGt7F * g4242, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmGgLtvsGt7F * g4242	93	23,3%	307	76,8%	400	100,0%

cmGgLtvsGt7F * g4242 Crosstabulation

Count		g4242			Total
		AA	GA	GG	
cmGgLtvsGt7F	<7	1	19	51	71
	>7	0	5	17	22
	Total	1	24	68	93

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	,484 ^a	2	,785
Likelihood Ratio	,716	2	,699
Linear-by-Linear Association	,474	1	,491
N of Valid Cases	93		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,24.

Risk Estimate

	Value
Odds Ratio for cmGgLtvsGt7F (<7 / >7)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=cmGgLt7vsEq7F BY g4242

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:57:50
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvfrez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=cmGgLt7vsEq7F BY g4242 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.

Notes

Resources	Processor Time	0:00:00.016
	Elapsed Time	0:00:00.014
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmb
dmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmGgL7vsEq7F * g4242, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmGgL7vsEq7F * g4242	128	32,0%	272	68,0%	400	100,0%

cmGgL7vsEq7F * g4242 Crosstabulation

Count

		g4242			Total
		AA	GA	GG	
cmGgL7vsEq7F	<7	1	19	51	71
	=7	1	18	38	57
	Total	2	37	89	128

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	,399 ^a	2	,819
Likelihood Ratio	,398	2	,819
Linear-by-Linear Association	,247	1	,619
N of Valid Cases	128		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,89.

Risk Estimate

	Value
Odds Ratio for cmGgL7vsEq7F (<7 / =7)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

```

/TABLES=cmGgEq7vsGt7F BY g4242
/FORMAT=AVALUE TABLES
/STATISTICS=CHISQ RISK CMH(1)
/CELLS=COUNT
/COUNT ROUND CELL.

```

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:57:51
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmvdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
Resources	Syntax	CROSSTABS /TABLES=cmGgEq7vsGt7F BY g4242 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
	Processor Time	0:00:00.015
	Elapsed Time	0:00:00.191
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmvdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmGgEq7vsGt7F * g4242, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmGgEq7vsGt7F * g4242	79	19,8%	321	80,3%	400	100,0%

cmGgEq7vsGt7F * g4242 Crosstabulation

Count

	g4242			Total
	AA	GA	GG	
cmGgEq7vsGt7F =7	1	18	38	57
>7	0	5	17	22
Total	1	23	55	79

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1,070 ^a	2	,586
Likelihood Ratio	1,353	2	,508
Linear-by-Linear Association	1,010	1	,315
N of Valid Cases	79		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,28.

Risk Estimate

	Value
Odds Ratio for cmGgEq7vsGt7F (=7 / >7)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=mMeta BY g4242

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Output Created	22-lip-2012 11:57:51
Comments	
Input Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmdbmv\rez\SPSS\Stat.sav
Active Dataset	DataSet1
Filter	<none>
Weight	<none>
Split File	<none>
N of Rows in Working Data File	400

Notes

Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=mMeta BY g4242 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
	Resources	
	Processor Time	0:00:00.015
	Elapsed Time	0:00:00.015
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmb
dmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for mMeta * g4242, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
mMeta * g4242	150	37,5%	250	62,5%	400	100,0%

mMeta * g4242 Crosstabulation

Count

		g4242			Total
		AA	GA	GG	
mMeta	no	1	25	69	95
	yes	1	17	37	55
	Total	2	42	106	150

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	,557 ^a	2	,757
Likelihood Ratio	,549	2	,760
Linear-by-Linear Association	,498	1	,481
N of Valid Cases	150		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,73.

Risk Estimate

	Value
Odds Ratio for mMeta (no / yes)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=mRiskEAU BY g4242

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:57:51
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=mRiskEAU BY g4242 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.016
	Elapsed Time	0:00:00.110
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for mRiskEAU * g4242, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
mRiskEAU * g4242	150	37,5%	250	62,5%	400	100,0%

mRiskEAU * g4242 Crosstabulation

Count

		g4242			Total
		AA	GA	GG	
mRiskEAU	low	0	3	11	14
	medium	1	15	39	55
	high	1	24	56	81
	Total	2	42	106	150

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	,744 ^a	4	,946
Likelihood Ratio	,934	4	,920
Linear-by-Linear Association	,238	1	,626
N of Valid Cases	150		

a. 4 cells (44,4%) have expected count less than 5. The minimum expected count is ,19.

Risk Estimate

	Value
Odds Ratio for mRiskEAU (low / medium)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=mRiskEAULowMedium BY g4242

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:57:52
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=mRiskEAULowMedium BY g4242 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.016
	Elapsed Time	0:00:00.014
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for mRiskEAULowMedium * g4242, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
mRiskEAULowMedium * g4242	69	17,3%	331	82,8%	400	100,0%

mRiskEAULowMedium * g4242 Crosstabulation

Count

		g4242			Total
		AA	GA	GG	
mRiskEAULowMedium	low	0	3	11	14
	medium	1	15	39	55
	Total	1	18	50	69

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	,491 ^a	2	,782
Likelihood Ratio	,695	2	,706
Linear-by-Linear Association	,479	1	,489
N of Valid Cases	69		

a. 3 cells (50,0%) have expected count less than 5. The minimum expected count is ,20.

Risk Estimate

	Value
Odds Ratio for mRiskEAULowMedium (low / medium)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=mRiskEAULowHigh BY g4242

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:57:52
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvfrez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.

Notes

Syntax	CROSSTABS /TABLES=mRiskEAULowHigh BY g4242 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.		
Resources	Processor Time	0:00:00.016	
	Elapsed Time	0:00:00.014	
	Dimensions Requested		2
	Cells Available		174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmb
dmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for mRiskEAULowHigh * g4242, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
mRiskEAULowHigh * g4242	95	23,8%	305	76,3%	400	100,0%

mRiskEAULowHigh * g4242 Crosstabulation

Count

		g4242			Total
		AA	GA	GG	
mRiskEAULowHigh	low	0	3	11	14
	high	1	24	56	81
	Total	1	27	67	95

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	,606 ^a	2	,739
Likelihood Ratio	,770	2	,681
Linear-by-Linear Association	,562	1	,453
N of Valid Cases	95		

a. 3 cells (50,0%) have expected count less than 5. The minimum expected count is ,15.

Risk Estimate

	Value
Odds Ratio for mRiskEAULowHigh (low / high)	a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=mRiskEAUMediumHigh BY g4242

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:57:52
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=mRiskEAUMediumHigh BY g4242 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.015
	Elapsed Time	0:00:00.128
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for mRiskEAUMediumHigh * g4242, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
mRiskEAUMediumHigh * g4242	136	34,0%	264	66,0%	400	100,0%

mRiskEAUMediumHigh * g4242 Crosstabulation

Count

		g4242			Total
		AA	GA	GG	
mRiskEAUMediumHigh	medium	1	15	39	55
	high	1	24	56	81
	Total	2	39	95	136

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	,154 ^a	2	,926
Likelihood Ratio	,153	2	,926
Linear-by-Linear Association	,002	1	,965
N of Valid Cases	136		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,81.

Risk Estimate

	Value
Odds Ratio for mRiskEAUMediumHigh (medium / high)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=mRiskMed BY g4242

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:57:53
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=mRiskMed BY g4242 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.016
	Elapsed Time	0:00:00.014
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for mRiskMed * g4242, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
mRiskMed * g4242	150	37,5%	250	62,5%	400	100,0%

mRiskMed * g4242 Crosstabulation

Count

		g4242			Total
		AA	GA	GG	
mRiskMed	low	1	13	41	55
	high	1	29	65	95
	Total	2	42	106	150

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	,929 ^a	2	,629
Likelihood Ratio	,936	2	,626
Linear-by-Linear Association	,066	1	,798
N of Valid Cases	150		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,73.

Risk Estimate

	Value
Odds Ratio for mRiskMed (low / high)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=mRiskMedLowMedium BY g4242

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:57:53
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvfrez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=mRiskMedLowMedium BY g4242 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.

Notes

Resources	Processor Time	0:00:00.015
	Elapsed Time	0:00:00.256
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmb
dmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for mRiskMedLowMedium * g4242, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

No measures of association are computed for the crosstabulation of mRiskMedLowMedium * g4242. At least one variable in each 2-way table upon which measures of association are computed is a constant.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
mRiskMedLowMedium * g4242	55	13,8%	345	86,3%	400	100,0%

mRiskMedLowMedium * g4242 Crosstabulation

Count

		g4242			Total
		AA	GA	GG	
mRiskMedLowMedium	low	1	13	41	55
	Total	1	13	41	55

Chi-Square Tests

	Value
Pearson Chi-Square	a
N of Valid Cases	55

a. No statistics are computed because mRiskMedLowMedium is a constant.

Risk Estimate

	Value
Odds Ratio for mRiskMedLowMedium (low / .)	a

a. No statistics are computed because mRiskMedLowMedium is a constant.

CROSSTABS

/TABLES=mRiskMedLowHigh BY g4242

```

/FORMAT=AVALUE TABLES
/STATISTICS=CHISQ RISK CMH(1)
/CELLS=COUNT
/COUNT ROUND CELL.

```

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:57:53
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=mRiskMedLowHigh BY g4242 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.016
	Elapsed Time	0:00:00.013
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for mRiskMedLowHigh * g4242, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
mRiskMedLowHigh * g4242	150	37,5%	250	62,5%	400	100,0%

mRiskMedLowHigh * g4242 Crosstabulation

Count

		g4242			Total
		AA	GA	GG	
mRiskMedLowHigh	low	1	13	41	55
	high	1	29	65	95
	Total	2	42	106	150

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	,929 ^a	2	,629
Likelihood Ratio	,936	2	,626
Linear-by-Linear Association	,066	1	,798
N of Valid Cases	150		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,73.

Risk Estimate

	Value
Odds Ratio for mRiskMedLowHigh (low / high)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=mRiskMedMediumHigh BY g4242

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 11:57:53
	Comments	
Data	Active Dataset	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Filter	DataSet1
Weight	Filter	<none>
	Weight	<none>
Split File	Split File	<none>
	N of Rows in Working Data File	400

Notes

Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=mRiskMedMediumHigh BY g4242 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.000
	Elapsed Time	0:00:00.162
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmb
dmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for mRiskMedMediumHigh * g4242, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

No measures of association are computed for the crosstabulation of mRiskMedMediumHigh * g4242. At least one variable in each 2-way table upon which measures of association are computed is a constant.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
mRiskMedMediumHigh * g4242	95	23,8%	305	76,3%	400	100,0%

mRiskMedMediumHigh * g4242 Crosstabulation

Count

		g4242			Total
		AA	GA	GG	
mRiskMedMediumHigh	high	1	29	65	95
	Total	1	29	65	95

Chi-Square Tests

	Value
Pearson Chi-Square	a.
N of Valid Cases	95

a. No statistics are computed because mRiskMedMediumHigh is a constant.

Risk Estimate

	Value
Odds Ratio for mRiskMedMediumHigh (high / .)	a

a. No statistics are computed
because mRiskMedMediumHigh
is a constant.