

```

CROSSTABS
/TABLES=diagPca BY g7837
/FORMAT=AVALUE TABLES
/STATISTICS=CHISQ RISK CMH(1)
/CELLS=COUNT
/COUNT ROUND CELL.

```

Crosstabs

Notes

Input	Output Created	22-lip-2012 12:07:06
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=diagPca BY g7837 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.031
	Elapsed Time	0:00:00.022
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for diagPca * g7837, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
diagPca * g7837	300	75,0%	100	25,0%	400	100,0%

diagPca * g7837 Crosstabulation

Count

		g7837			Total
		GG	GT	TT	
diagPca	no	122	28	0	150
	yes	101	47	2	150
	Total	223	75	2	300

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	8,791 ^a	2	,012
Likelihood Ratio	9,619	2	,008
Linear-by-Linear Association	8,306	1	,004
N of Valid Cases	300		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is 1,00.

Risk Estimate

	Value
Odds Ratio for diagPca (no / yes)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=cmDiagPca0Kont BY g7837

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Output Created	22-lip-2012 12:07:06
Comments	
Input Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
Active Dataset	DataSet1
Filter	<none>
Weight	<none>
Split File	<none>
N of Rows in Working Data File	400

Notes

Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing. Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table. CROSSTABS /TABLES=cmDiagPca0Kont BY g7837 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
	Cases Used	
	Syntax	
	Resources	
	Processor Time	0:00:00.016
	Elapsed Time	0:00:00.011
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmb
dmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmDiagPca0Kont * g7837, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmDiagPca0Kont * g7837	250	62,5%	150	37,5%	400	100,0%

cmDiagPca0Kont * g7837 Crosstabulation

Count

		g7837			Total
		GG	GT	TT	
cmDiagPca0Kont	no	122	28	0	150
	control	82	16	2	100
	Total	204	44	2	250

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	3,246 ^a	2	,197
Likelihood Ratio	3,914	2	,141
Linear-by-Linear Association	,976	1	,323
N of Valid Cases	250		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,80.

Risk Estimate

	Value
Odds Ratio for cmDiagPca0Kont (no / control)	a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=cmDiagPca1Kont BY g7837

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 12:07:06
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=cmDiagPca1Kont BY g7837 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.016
	Elapsed Time	0:00:00.011
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmDiagPca1Kont * g7837, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmDiagPca1Kont * g7837	250	62,5%	150	37,5%	400	100,0%

cmDiagPca1Kont * g7837 Crosstabulation

Count

		g7837			Total
		GG	GT	TT	
cmDiagPca1Kont	yes	101	47	2	150
	control	82	16	2	100
	Total	183	63	4	250

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	7,528 ^a	2	,023
Likelihood Ratio	7,847	2	,020
Linear-by-Linear Association	1,588	1	,208
N of Valid Cases	250		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is 1,60.

Risk Estimate

	Value
Odds Ratio for cmDiagPca1Kont (yes / control)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=kontrol BY g7837

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 12:07:06
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=kontrol BY g7837 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.015
	Elapsed Time	0:00:00.013
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for kontrol * g7837, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
kontrol * g7837	400	100,0%	0	,0%	400	100,0%

kontrol * g7837 Crosstabulation

Count		g7837			Total
		GG	GT	TT	
kontrol	no control	223	75	2	300
	control	82	16	2	100
	Total	305	91	4	400

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	4,582 ^a	2	,101
Likelihood Ratio	4,603	2	,100
Linear-by-Linear Association	,102	1	,750
N of Valid Cases	400		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is 1,00.

Risk Estimate

	Value
Odds Ratio for kontrol (no kontrol / kontrol)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=cmTStadOnly12 BY g7837

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 12:07:06
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvfrez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=cmTStadOnly12 BY g7837 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.

Notes

Resources	Processor Time	0:00:00.016
	Elapsed Time	0:00:00.028
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmb
dmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmTStadOnly12 * g7837, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmTStadOnly12 * g7837	100	25,0%	300	75,0%	400	100,0%

cmTStadOnly12 * g7837 Crosstabulation

Count

		g7837			Total
		GG	GT	TT	
cmTStadOnly12	T1	21	6	1	28
	T2	47	25	0	72
	Total	68	31	1	100

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	4,001 ^a	2	,135
Likelihood Ratio	4,059	2	,131
Linear-by-Linear Association	,097	1	,756
N of Valid Cases	100		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,28.

Risk Estimate

	Value
Odds Ratio for cmTStadOnly12 (T1 / T2)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.


```

/TABLES=cmTStadOnly13 BY g7837
/FORMAT=AVALUE TABLES
/STATISTICS=CHISQ RISK CMH(1)
/CELLS=COUNT
/COUNT ROUND CELL.

```

Crosstabs

Notes

Input	Output Created	22-lip-2012 12:07:06
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=cmTStadOnly13 BY g7837 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.016
	Elapsed Time	0:00:00.021
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmTStadOnly13 * g7837, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmTStadOnly13 * g7837	78	19,5%	322	80,5%	400	100,0%

cmTStadOnly13 * g7837 Crosstabulation

Count

		g7837			Total
		GG	GT	TT	
cmTStadOnly13	T1	21	6	1	28
	T3,T4	33	16	1	50
	Total	54	22	2	78

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1,094 ^a	2	,579
Likelihood Ratio	1,115	2	,573
Linear-by-Linear Association	,017	1	,895
N of Valid Cases	78		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,72.

Risk Estimate

	Value
Odds Ratio for cmTStadOnly13 (T1 / T3, T4)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=cmTStadOnly23 BY g7837

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Output Created	22-lip-2012 12:07:07
Comments	
Input Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
Active Dataset	DataSet1
Filter	<none>
Weight	<none>
Split File	<none>
N of Rows in Working Data File	400

Notes

Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=cmTstadOnly23 BY g7837 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.000
	Elapsed Time	0:00:00.040
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmb
dmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmTstadOnly23 * g7837, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmTstadOnly23 * g7837	122	30,5%	278	69,5%	400	100,0%

cmTstadOnly23 * g7837 Crosstabulation

Count

		g7837			Total
		GG	GT	TT	
cmTstadOnly23	T2	47	25	0	72
	T3,T4	33	16	1	50
	Total	80	41	1	122

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1,507 ^a	2	,471
Likelihood Ratio	1,852	2	,396
Linear-by-Linear Association	,389	1	,533
N of Valid Cases	122		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,41.

Risk Estimate

	Value
Odds Ratio for cmTStadOnly23 (T2 / T3, T4)	a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=cmPsaLT10vs10to20FonPCA1 BY g7837

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 12:07:07
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmvdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=cmPsaLT10vs10to20FonPCA1 BY g7837 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.000
	Elapsed Time	0:00:00.017
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmvdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmPsaLT10vs10to20FonPCA1 * g7837, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmPsaLT10vs10to20FonPCA1 * g7837	88	22,0%	312	78,0%	400	100,0%

cmPsaLT10vs10to20FonPCA1 * g7837 Crosstabulation

Count

		g7837			Total
		GG	GT	TT	
cmPsaLT10vs10to20FonPCA1	<10	34	14	1	49
	10-20	28	11	0	39
	Total	62	25	1	88

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	,815 ^a	2	,665
Likelihood Ratio	1,190	2	,552
Linear-by-Linear Association	,531	1	,466
N of Valid Cases	88		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,44.

Risk Estimate

	Value
Odds Ratio for cmPsaLT10vs10to20FonPCA1 (<10 / 10-20)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=cmPsaLT10vsGT20FonPCA1 BY g7837

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 12:07:07
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=cmPsaLT10vsGT20FonPCA1 BY g7837 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.031
	Elapsed Time	0:00:00.020
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmPsaLT10vsGT20FonPCA1 * g7837, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmPsaLT10vsGT20FonPCA1 * g7837	111	27,8%	289	72,3%	400	100,0%

cmPsaLT10vsGT20FonPCA1 * g7837 Crosstabulation

Count		g7837			Total
		GG	GT	TT	
cmPsaLT10vsGT20FonPCA1	<10	34	14	1	49
	>20	39	22	1	62
	Total	73	36	2	111

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	,606 ^a	2	,739
Likelihood Ratio	,609	2	,737
Linear-by-Linear Association	,102	1	,749
N of Valid Cases	111		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,88.

Risk Estimate

	Value
Odds Ratio for cmPsaLT10vsGT20Fon PCA1 (<10 / >20)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=cmPsa10to20vsGT20FonPCA1 BY g7837

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 12:07:07
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvfrez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
Missing Value Handling	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.

Notes

Syntax	CROSSTABS /TABLES=cmPsa10to20vsGT20Fon PCA1 BY g7837 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.		
Resources	Processor Time	0:00:00.016	
	Elapsed Time	0:00:00.011	
	Dimensions Requested		2
	Cells Available		174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmb
dmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmPsa10to20vsGT20FonPCA1 * g7837, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmPsa10to20vsGT20Fon PCA1 * g7837	101	25,3%	299	74,8%	400	100,0%

cmPsa10to20vsGT20FonPCA1 * g7837 Crosstabulation

Count

		g7837			Total
		GG	GT	TT	
cmPsa10to20vsGT20Fon PCA1	10-20	28	11	0	39
	>20	39	22	1	62
	Total	67	33	1	101

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1,303 ^a	2	,521
Likelihood Ratio	1,654	2	,437
Linear-by-Linear Association	1,290	1	,256
N of Valid Cases	101		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,39.

Risk Estimate

	Value
Odds Ratio for cmPsa10to20vsGT20Fon PCA1 (10-20 / >20)	a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=cmPsaLT20vsGT20onPCA1 BY g7837

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 12:07:07
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=cmPsaLT20vsGT20onPCA1 BY g7837 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.016
	Elapsed Time	0:00:00.020
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmPsaLT20vsGT20onPCA1 * g7837, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmPsaLT20vsGT20onPCA1 * g7837	150	37,5%	250	62,5%	400	100,0%

cmPsaLT20vsGT20onPCA1 * g7837 Crosstabulation

Count

		g7837			Total
		GG	GT	TT	
cmPsaLT20vsGT20onPCA1	,00	62	25	1	88
	<10	39	22	1	62
	Total	101	47	2	150

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	,951 ^a	2	,622
Likelihood Ratio	,946	2	,623
Linear-by-Linear Association	,643	1	,422
N of Valid Cases	150		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,83.

Risk Estimate

	Value
Odds Ratio for cmPsaLT20vsGT20onPCA1 (,00 / <10)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=cmGgLtvsGt7F BY g7837

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 12:07:07
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=cmGgLtvsGt7F BY g7837 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.015
	Elapsed Time	0:00:00.012
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmGgLtvsGt7F * g7837, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmGgLtvsGt7F * g7837	93	23,3%	307	76,8%	400	100,0%

cmGgLtvsGt7F * g7837 Crosstabulation

Count		g7837			Total
		GG	GT	TT	
cmGgLtvsGt7F	<7	50	21	0	71
	>7	15	6	1	22
	Total	65	27	1	93

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	3,270 ^a	2	,195
Likelihood Ratio	2,926	2	,231
Linear-by-Linear Association	1,572	1	,210
N of Valid Cases	93		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,24.

Risk Estimate

	Value
Odds Ratio for cmGgLtvsGt7F (<7 / >7)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=cmGgLt7vsEq7F BY g7837

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 12:07:07
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvfrez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=cmGgLt7vsEq7F BY g7837 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.

Notes

Resources	Processor Time	0:00:00.015
	Elapsed Time	0:00:00.015
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmb
dmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmGgLt7vsEq7F * g7837, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmGgLt7vsEq7F * g7837	128	32,0%	272	68,0%	400	100,0%

cmGgLt7vsEq7F * g7837 Crosstabulation

Count

		g7837			Total
		GG	GT	TT	
cmGgLt7vsEq7F	<7	50	21	0	71
	=7	36	20	1	57
	Total	86	41	1	128

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1,794 ^a	2	,408
Likelihood Ratio	2,166	2	,339
Linear-by-Linear Association	1,656	1	,198
N of Valid Cases	128		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,45.

Risk Estimate

	Value
Odds Ratio for cmGgLt7vsEq7F (<7 / =7)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

```

/TABLES=cmGgEq7vsGt7F BY g7837
/FORMAT=AVALUE TABLES
/STATISTICS=CHISQ RISK CMH(1)
/CELLS=COUNT
/COUNT ROUND CELL.

```

Crosstabs

Notes

Input	Output Created	22-lip-2012 12:07:07
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmvdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
Resources	Syntax	CROSSTABS /TABLES=cmGgEq7vsGt7F BY g7837 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
	Processor Time	0:00:00.015
	Elapsed Time	0:00:00.016
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmvdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for cmGgEq7vsGt7F * g7837, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
cmGgEq7vsGt7F * g7837	79	19,8%	321	80,3%	400	100,0%

cmGgEq7vsGt7F * g7837 Crosstabulation

Count

	g7837			Total
	GG	GT	TT	
cmGgEq7vsGt7F =7	36	20	1	57
>7	15	6	1	22
Total	51	26	2	79

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	,845 ^a	2	,655
Likelihood Ratio	,804	2	,669
Linear-by-Linear Association	,078	1	,780
N of Valid Cases	79		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,56.

Risk Estimate

	Value
Odds Ratio for cmGgEq7vsGt7F (=7 / >7)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=mMeta BY g7837

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Output Created	22-lip-2012 12:07:07
Comments	
Input Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmv\rez\SPSS\Stat.sav
Active Dataset	DataSet1
Filter	<none>
Weight	<none>
Split File	<none>
N of Rows in Working Data File	400

Notes

Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=mMeta BY g7837 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.016
	Elapsed Time	0:00:00.017
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmb
dmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for mMeta * g7837, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
mMeta * g7837	150	37,5%	250	62,5%	400	100,0%

mMeta * g7837 Crosstabulation

Count

		g7837			Total
		GG	GT	TT	
mMeta	no	66	28	1	95
	yes	35	19	1	55
	Total	101	47	2	150

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	,615 ^a	2	,735
Likelihood Ratio	,607	2	,738
Linear-by-Linear Association	,546	1	,460
N of Valid Cases	150		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,73.

Risk Estimate

	Value
Odds Ratio for mMeta (no / yes)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=mRiskEAU BY g7837

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 12:07:07
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=mRiskEAU BY g7837 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.015
	Elapsed Time	0:00:00.016
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for mRiskEAU * g7837, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
mRiskEAU * g7837	150	37,5%	250	62,5%	400	100,0%

mRiskEAU * g7837 Crosstabulation

Count

		g7837			Total
		GG	GT	TT	
mRiskEAU	low	11	3	0	14
	medium	38	17	0	55
	high	52	27	2	81
	Total	101	47	2	150

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	2,655 ^a	4	,617
Likelihood Ratio	3,457	4	,484
Linear-by-Linear Association	2,217	1	,137
N of Valid Cases	150		

a. 4 cells (44,4%) have expected count less than 5. The minimum expected count is ,19.

Risk Estimate

	Value
Odds Ratio for mRiskEAU (low / medium)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=mRiskEAULowMedium BY g7837

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 12:07:07
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=mRiskEAULowMedium BY g7837 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.031
	Elapsed Time	0:00:00.027
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
mRiskEAULowMedium * g7837	69	17,3%	331	82,8%	400	100,0%

mRiskEAULowMedium * g7837 Crosstabulation

Count		g7837		Total
		GG	GT	
mRiskEAULowMedium	low	11	3	14
	medium	38	17	55
	Total	49	20	69

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	,487 ^a	1	,485		
Continuity Correction ^b	,136	1	,713		
Likelihood Ratio	,510	1	,475		
Fisher's Exact Test				,742	,367
Linear-by-Linear Association	,480	1	,488		
N of Valid Cases	69				

a. 1 cells (25,0%) have expected count less than 5. The minimum expected count is 4,06.

b. Computed only for a 2x2 table

Risk Estimate

	95% Confidence Interval		
	Value	Lower	Upper
Odds Ratio for mRiskEAULowMedium (low / medium)	1,640	,405	6,644
For cohort g7837 = GG	1,137	,821	1,575
For cohort g7837 = GT	,693	,236	2,038
N of Valid Cases	69		

Tests of Homogeneity of the Odds Ratio

	Chi-Squared	df	Asymp. Sig. (2-sided)
Breslow-Day	,000	0	.
Tarone's	,000	0	.

Tests of Conditional Independence

	Chi-Squared	df	Asymp. Sig. (2-sided)
Cochran's	,487	1	,485
Mantel-Haenszel	,134	1	,715

Under the conditional independence assumption, Cochran's statistic is asymptotically distributed as a 1 df chi-squared distribution, only if the number of strata is fixed, while the Mantel-Haenszel statistic is always asymptotically distributed as a 1 df chi-squared distribution. Note that the continuity correction is removed from the Mantel-Haenszel statistic when the sum of the differences between the observed and the expected is 0.

Mantel-Haenszel Common Odds Ratio Estimate

Estimate	1,640
ln(Estimate)	,495
Std. Error of ln(Estimate)	,714
Asymp. Sig. (2-sided)	,488

The Mantel-Haenszel common odds ratio estimate is asymptotically normally distributed under the common odds ratio of 1,000 assumption. So is the natural log of the estimate.

Mantel-Haenszel Common Odds Ratio Estimate

Asymp. 95% Confidence Interval	Common Odds Ratio	Lower Bound	,405
		Upper Bound	6,644
	In(Common Odds Ratio)	Lower Bound	-,904
		Upper Bound	1,894

The Mantel-Haenszel common odds ratio estimate is asymptotically normally distributed under the common odds ratio of 1,000 assumption. So is the natural log of the estimate.

CROSSTABS

/TABLES=mRiskEAULowHigh BY g7837

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 12:07:07
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=mRiskEAULowHigh BY g7837 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.015
	Elapsed Time	0:00:00.014
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for mRiskEAULowHigh * g7837, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
mRiskEAULowHigh * g7837	95	23,8%	305	76,3%	400	100,0%

mRiskEAULowHigh * g7837 Crosstabulation

Count

		g7837			Total
		GG	GT	TT	
mRiskEAULowHigh	low	11	3	0	14
	high	52	27	2	81
	Total	63	30	2	95

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1,253 ^a	2	,534
Likelihood Ratio	1,585	2	,453
Linear-by-Linear Association	1,045	1	,307
N of Valid Cases	95		

a. 3 cells (50,0%) have expected count less than 5. The minimum expected count is ,29.

Risk Estimate

	Value
Odds Ratio for mRiskEAULowHigh (low / high)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=mRiskEAUMediumHigh BY g7837

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 12:07:07
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=mRiskEAUMediumHigh BY g7837 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.032
	Elapsed Time	0:00:00.038
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for mRiskEAUMediumHigh * g7837, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
mRiskEAUMediumHigh * g7837	136	34,0%	264	66,0%	400	100,0%

mRiskEAUMediumHigh * g7837 Crosstabulation

Count		g7837			Total
		GG	GT	TT	
mRiskEAUMediumHigh	medium	38	17	0	55
	high	52	27	2	81
	Total	90	44	2	136

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1,536 ^a	2	,464
Likelihood Ratio	2,251	2	,325
Linear-by-Linear Association	1,330	1	,249
N of Valid Cases	136		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,81.

Risk Estimate

	Value
Odds Ratio for mRiskEAUMediumHigh (medium / high)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=mRiskMed BY g7837

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 12:07:07
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvfrez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
Missing Value Handling	Split File	<none>
	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=mRiskMed BY g7837 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.

Notes

Resources	Processor Time	0:00:00.000
	Elapsed Time	0:00:00.014
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmb
dmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for mRiskMed * g7837, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
mRiskMed * g7837	150	37,5%	250	62,5%	400	100,0%

mRiskMed * g7837 Crosstabulation

Count

		g7837			Total
		GG	GT	TT	
mRiskMed	low	41	14	0	55
	high	60	33	2	95
	Total	101	47	2	150

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	2,787 ^a	2	,248
Likelihood Ratio	3,476	2	,176
Linear-by-Linear Association	2,703	1	,100
N of Valid Cases	150		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,73.

Risk Estimate

	Value
Odds Ratio for mRiskMed (low / high)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

```

/TABLES=mRiskMedLowMedium BY g7837
/FORMAT=AVALUE TABLES
/STATISTICS=CHISQ RISK CMH(1)
/CELLS=COUNT
/COUNT ROUND CELL.

```

Crosstabs

Notes

Input	Output Created	22-lip-2012 12:07:08
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
Resources	Syntax	CROSSTABS /TABLES=mRiskMedLowMedium BY g7837 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
	Processor Time	0:00:00.015
	Elapsed Time	0:00:00.022
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

No measures of association are computed for the crosstabulation of mRiskMedLowMedium * g7837. At least one variable in each 2-way table upon which measures of association are computed is a constant.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
mRiskMedLowMedium * g7837	55	13,8%	345	86,3%	400	100,0%

mRiskMedLowMedium * g7837 Crosstabulation

Count

		g7837		Total
		GG	GT	
mRiskMedLowMedium	low	41	14	55
	Total	41	14	55

Chi-Square Tests

	Value
Pearson Chi-Square	a.
N of Valid Cases	55

a. No statistics are computed because mRiskMedLowMedium is a constant.

Risk Estimate

	Value
Odds Ratio for mRiskMedLowMedium (low / .)	a.

a. No statistics are computed because mRiskMedLowMedium is a constant.

CROSSTABS

/TABLES=mRiskMedLowHigh BY g7837

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 12:07:08
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvfrez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.

Notes

Syntax	CROSSTABS /TABLES=mRiskMedLowHigh BY g7837 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.		
Resources	Processor Time	0:00:00.015	
	Elapsed Time	0:00:00.014	
	Dimensions Requested		2
	Cells Available		174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmb
dmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for mRiskMedLowHigh * g7837, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
mRiskMedLowHigh * g7837	150	37,5%	250	62,5%	400	100,0%

mRiskMedLowHigh * g7837 Crosstabulation

Count

		g7837			Total
		GG	GT	TT	
mRiskMedLowHigh	low	41	14	0	55
	high	60	33	2	95
	Total	101	47	2	150

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	2,787 ^a	2	,248
Likelihood Ratio	3,476	2	,176
Linear-by-Linear Association	2,703	1	,100
N of Valid Cases	150		

a. 2 cells (33,3%) have expected count less than 5. The minimum expected count is ,73.

Risk Estimate

	Value
Odds Ratio for mRiskMedLowHigh (low / high)	a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

CROSSTABS

/TABLES=mRiskMedMediumHigh BY g7837

/FORMAT=AVALUE TABLES

/STATISTICS=CHISQ RISK CMH(1)

/CELLS=COUNT

/COUNT ROUND CELL.

Crosstabs

Notes

Input	Output Created	22-lip-2012 12:07:08
	Comments	
	Data	U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
Missing Value Handling	N of Rows in Working Data File	400
	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
	Syntax	CROSSTABS /TABLES=mRiskMedMediumHigh BY g7837 /FORMAT=AVALUE TABLES /STATISTICS=CHISQ RISK CMH(1) /CELLS=COUNT /COUNT ROUND CELL.
Resources	Processor Time	0:00:00.016
	Elapsed Time	0:00:00.017
	Dimensions Requested	2
	Cells Available	174762

[DataSet1] U:\Personal Data\My Folders\Science\WorkCurrent_rad_b01_x_dsmbdmvf\rez\SPSS\Stat.sav

Warnings

The Tests for Homogeneity of the Odds Ratio table and the Mantel-Haenszel Common Odds Ratio Estimate table are not computed for mRiskMedMediumHigh * g7837, because either (1) the group variable does not have exactly two distinct non-missing values or/and (2) the response variable does not have exactly two distinct non-missing values.

No measures of association are computed for the crosstabulation of mRiskMedMediumHigh * g7837. At least one variable in each 2-way table upon which measures of association are computed is a constant.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
mRiskMedMediumHigh * g7837	95	23,8%	305	76,3%	400	100,0%

mRiskMedMediumHigh * g7837 Crosstabulation

Count

		g7837			Total
		GG	GT	TT	
mRiskMedMediumHigh	high	60	33	2	95
	Total	60	33	2	95

Chi-Square Tests

	Value
Pearson Chi-Square	a
N of Valid Cases	95

a. No statistics are computed because mRiskMedMediumHigh is a constant.

Risk Estimate

	Value
Odds Ratio for mRiskMedMediumHigh (high / .)	a

a. No statistics are computed because mRiskMedMediumHigh is a constant.