Identities used: binomial therom
$(x+a)^r = \sum_{b=0}^{r} \left[\binom{r}{b} x^b a^{r-b} \right]$
pascals identity
$\binom{a}{n} = \binom{a-1}{n} + \binom{a-1}{n-1}$

Reucrsions from Binomial therom

Non falling sums

$$(x+a)^r = \sum_{b=0}^{r} \left[\binom{r}{b} x^b a^{r-b} \right]$$

let $a = 1$

$$(x+1)^r = \sum_{b=0}^{r} \left[\binom{r}{b} x^b \right]$$

let $x = x' - 1$

$$((x'-1)+1)^r = x^r = \sum_{b=0}^{r} \left[\binom{r}{b} (x'-1)^b \right]$$

in general

$$((x-b)+a)^r = \sum_{b=0}^{r} \left[\binom{r}{b} (x-b)^b a^{r-b} \right]$$

if b=a

$$x^r = \sum_{b=0}^{r} \left[\binom{r}{b} (x-a)^b a^{r-b} \right]$$

we can subtitute the defintion of x^r into $(x-a)^b$ or a^{r-b}
$(x-a)^b$ is a good substitution as it the nested power term will shrink by a
after $\frac{x}{a}$ expansions the inner power term will be 0

$$x^r = \sum_{b=0}^{r} \left[\binom{r}{b} (x-a)^b a^{r-b} \right]$$

$$(x-a)^b = \sum_{c=0}^{b} \left[\binom{b}{c} ((x-a)-a)^c a^{b-c} \right]$$

$$x^r = \sum_{b=0}^{r} \left[\binom{r}{b} \sum_{c=0}^{b} \left[\binom{b}{c} ((x-a)-a)^c a^{b-c} \right] a^{r-b} \right]$$

every layer strips one a from inner power's base

$$x^r = \sum_{b=0}^{r} \left[\binom{r}{b} \sum_{c=0}^{b} \left[\binom{b}{c} ((x-a)-a)^c a^{b-c} \right] a^{r-b} \right]$$

$$x^r = \overbrace{\sum_{b=0}^{r} \left[\binom{r}{b} \sum_{c=0}^{b} \left[\binom{b}{c} \sum_{d=0}^{c} \left[\binom{c}{d} \dots \left[\sum_{n_{\frac{x}{a}}=0}^{n_{(\frac{x}{a})-1}} \left[\binom{n_{(\frac{x}{a})-1}}{n_{\frac{x}{a}}} 0^{n_{\frac{x}{a}}} a^{n_{(\frac{x}{a})-1}-n_{\frac{x}{a}}} \right] \right] a^{c-d} \right] a^{b-c} \right] a^{r-b} \right]}^{\frac{x}{a} \text{ sums}}$$

you can stop at any layer you want, you dont need to fully expand

$$x^r = \overbrace{\sum_{b=0}^{r} \left[\binom{r}{b} \sum_{c=0}^{b} \left[\binom{b}{c} \sum_{d=0}^{c} \left[\binom{c}{d} \dots \left[\sum_{n_l=0}^{n_{l-1}} \left[\binom{n_{l-1}}{n_l} (x-la)^{n_l} a^{n_{l-1}-n_l} \right] \right] a^{c-d} \right] a^{b-c} \right] a^{r-b} \right]}^{l \text{ sums}}$$

falling sums

$$(x+a)^r = \sum_{b=0}^{r} \left[\binom{r}{b} x^b a^{r-b} \right]$$

$$(x+a)^r = \sum_{b=0}^{r-1} \left[\binom{r}{b} x^b a^{r-b} \right] + \binom{r}{r} x^r a^0$$

$$((x-a)+a)^r = x^r = \sum_{b=0}^{r-1} \left[\binom{r}{b} (x-a)^b a^{r-b} \right] + \binom{r}{r} (x-a)^r a^0$$

the same reassoning applys here as in the last case:
"we can subtitute the defintion of x^r into $(x-a)^b$ or a^{r-b}
$(x-a)^b$ is a good substitution as it the nested power term will shrink by a
after $\frac{x}{a}$ expansions the inner power term will be 0 "

$$x^r = \sum_{b=0}^{r-1}\left[\binom{r}{b}(x-a)^b a^{r-b}\right] + \binom{r}{r}(x-a)^r a^0$$

$$(x-a)^b = \sum_{c=0}^{b-1}\left[\binom{r}{b}((x-a)-a)^c a^{b-c}\right] + \binom{b}{b}((x-a)-a)^b a^0$$

$$x^r = \sum_{b=0}^{r-1}\left[\binom{r}{b}\left[\sum_{c=0}^{b-1}\left[\binom{r}{b}((x-a)-a)^c a^{b-c}\right] + \binom{b}{b}((x-a)-a)^b a^0\right] a^{r-b}\right] + \binom{r}{r}(x-a)^r a^0$$

$$\overbrace{\sum_{b=0}^{r-1}\left[\binom{r}{b}\left[\sum_{c=0}^{b-1}\left[\binom{r}{b}\right.\right.\right.}^{l \text{ sums}}$$

$$\left.\left.\left.\cdots\left[\sum_{n_l=0}^{n_{l-1}-1}\left[\binom{n_{l-1}}{n_l}(x-la)^{n_l}a^{n_{l-1}-n_l}\right] + \binom{n_{l-1}}{n_{l-1}}(x-la)^{n_{l-1}}a^0\right]\cdots\right.\right.\right.$$

$$\left.\left.\left.a^{b-c}\right] + \binom{b}{b}(x-2a)^b a^0\right] a^{r-b}\right] + \binom{r}{r}(x-a)^r a^0$$

$$x^r =$$

$$a^0 = 1 \text{ for all } a \in \mathbb{R}$$

$$\binom{r}{r} = 1 \text{ for all } r \in \mathbb{Z}^+$$

if we restrict our formula with these we can simplify it

according to our restriciton $a^0 = 1$ and $\binom{r}{r} = 1$

$$x^r = \overbrace{\sum_{b=0}^{r-1}\left[\binom{r}{b}\left[\sum_{c=0}^{b-1}\left[\binom{r}{b}\cdots\left[\sum_{n_l=0}^{n_{l-1}-1}\left[\binom{n_{l-1}}{n_l}(x-la)^{n_l}a^{n_{l-1}-n_l}\right] + (x-la)^{n_{l-1}}\right]\cdots a^{b-c}\right] + (x-2a)^b\right]a^{r-b}\right] + (x-a)^r}^{l \text{ sums}}$$

the sum can be factored into a sum of falling sums The papers theory holds if the restrictions hold:

$$a^0 = 1 \text{ for all } a \in \mathbb{R}$$

$\binom{r}{r} = 1$ for all $r \in \mathbb{Z}^+$ Since we can freely choose a only the restriction on r remains

“I have a theory, all x^n’s can be written as a sum of nested sums with falling upper bounds “

all x^r’s can be written as a sum of nested sums with falling upper bounds for all $r \in \mathbb{Z}^+$ also x must have a divisor d in $\mathbb{R}$ where $\left(\frac{x}{d} = l, l \in \mathbb{Z}^+\right)$, if it has this divisor it can be written a nested sum with l sums for all $a \in \mathbb{R} \mid r \in \mathbb{Z}^+ \mid d$ in $\mathbb{R}$ where $\left(\frac{x}{d} = l, l \in \mathbb{Z}^+\right) \Rightarrow$

$$x^r = \overbrace{\sum_{b=0}^{r-1}\left[\binom{r}{b}\left[\sum_{c=0}^{b-1}\left[\binom{r}{b}\cdots\left[\sum_{n_l=0}^{n_{l-1}-1}\left[\binom{n_{l-1}}{n_l}(x-la)^{n_l}a^{n_{l-1}-n_l}\right]\right]\cdots a^{b-c}\right]\right]a^{r-b}\right]}^{l \text{ sums}}$$

$$+\overbrace{\sum_{b=0}^{r-1}\left[\binom{r}{b}\left[\sum_{c=0}^{b-1}\left[\binom{r}{b}\cdots\left[\sum_{n_{l-1}=0}^{n_{l-2}-1}[(x-la)^{n_{l-1}}]\right]\cdots a^{b-c}\right]\right]a^{r-b}\right]}^{l-1 \text{ sums}}$$

$$+\ldots+\overbrace{\sum_{b=0}^{r-1}\left[\binom{r}{b}[(x-2a)^b]a^{r-b}\right]}^{1 \text{ sums}}$$

$$+\overbrace{(x-a)^r}^{0 \text{ sums}}$$

Falling sums (l,r upper bound)

$$\sum_{b=0}^{r}\left[\binom{r}{b}x^b a^{r-b}\right](x+a)^r = \sum_{b=0}^{r-1}\left[\binom{r}{b}x^b a^{r-b}\right] + \binom{r}{r}x^r a^0 \text{ let } x' = x-a, \text{let } a' = a((x-a)+a)^r = \sum_{b=0}^{r-1}\left[\binom{r}{b}(x-a)^b a^{r-b}\right] + \binom{r}{r}$$

the previous falling sums resulted from expanding the inner (x-a)^b, if we expand the outer (x-a)^r we get a diffrent falling sum

$$((x-a)+a)^r = x^r = \sum_{b=0}^{r-1}\left[\binom{r}{b}(x-a)^b a^{r-b}\right] + \binom{r}{r}(x-a)^r a^0$$

$$x^r = \sum_{b=0}^{r-1}\left[\binom{r}{b}(x-a)^b a^{r-b}\right] + \binom{r}{r}\left[\sum_{b=0}^{r-1}\left[\binom{r}{b}((x-a)-a)^b a^{r-b}\right] + \binom{r}{r}((x-a)-a)^r a^0\right]a^0$$

$$x^r = \overbrace{\sum_{b=0}^{r-1}\left[\binom{r}{b}(x-a)^b a^{r-b}\right] + \binom{r}{r}\left[\sum_{b=0}^{r-1}\left[\binom{r}{b}((x-a)-a)^b a^{r-b}\right] + \binom{r}{r}\right.}^{l \text{ sums}}$$

$$\left.\cdots\left[\sum_{b=0}^{r-1}\left[\binom{r}{b}(x-la)^b a^{r-b}\right] + \binom{r}{r}(x-la)^r a^0\right]\cdots a^0\right]a^0$$

$$x^r = \overbrace{\sum_{b=0}^{r-1}\left[\binom{r}{b}(x-a)^b a^{r-b}\right]}^{l \text{ sums}}$$

$$+\binom{r}{r}a^0\sum_{b=0}^{r-1}\left[\binom{r}{b}(x-2a)^b a^{r-b}\right]$$

$$+\binom{r}{r}^2 a^{(0)^2}\sum_{b=0}^{r-1}\left[\binom{r}{b}(x-3a)^b a^{r-b}\right]$$

$$+\cdots+\binom{r}{r}^{l-1} a^{(0)^{l-1}}\sum_{b=0}^{r-1}\left[\binom{r}{b}(x-la)^b a^{r-b}\right]$$

$$x^r = +\binom{r}{r}^{l}(x-la)^r a^{(0)^l}$$

$$x^r = \sum_{b=0}^{l-1}\left[\binom{r}{r}^b a^{(0)^b}\sum_{c=0}^{r-1}\left[\binom{r}{c}(x-(b+1)a)^c a^{r-c}\right]\right] + \binom{r}{r}^{l}(x-la)^r a^{(0)^l}$$

note: for the simplifed sum aboe to work binom(r,r)^(0) and a^(0)^(0) must simplify to 1, so a^0 is implictly assumed to be 1 for that sum to work. you could factor out first term of the sum to make it pure

if we add restrictions the sum simplifies

$$a^0 = 1 \text{ for all } a \in \mathbb{R}$$

$$\binom{r}{r} = 1 \text{ for all } r \in \mathbb{Z}^+$$

$$\text{for all } a \in \mathbb{R} \mid r \in \mathbb{Z}^+ \mid l \in \mathbb{Z}^+$$

$$x^r = \sum_{b=0}^{l-1}\left[\sum_{c=0}^{r-1}\left[\binom{r}{c}(x-(b+1)a)^c a^{r-c}\right]\right] + (x-la)^r$$

falling sums, free upper bounds

if we just keep exapnding the new outer term derived from the falling l,r formula with the falling l,r formula we get more nested sums

for all $a \in \mathbb{R} \mid r \in \mathbb{Z}^+ \mid l \in \mathbb{Z}^+$

$$x^r = \sum_{b=0}^{l-1} \left[\sum_{c=0}^{r-1} \left[\binom{r}{c} (x-(b+1)a)^c a^{r-c} \right] \right] + (x-la)^r$$

$$x^r = \sum_{b=0}^{l-1} \left[\sum_{c=0}^{r-1} \left[\binom{r}{c} (x-(b+1)a)^c a^{r-c} \right] \right] + \sum_{b=0}^{l-1} \left[\sum_{c=0}^{r-1} \left[\binom{r}{c} ((x-la)-(b+1)a)^c a^{r-c} \right] \right] + ((x-la)-la)^r$$

$$x^r = \sum_{d=0}^{m-1} \left[\sum_{b=0}^{l-1} \left[\sum_{c=0}^{r-1} \left[\binom{r}{c} ((x-dla)-(b+1)a)^c a^{r-c} \right] \right] \right] + (x-mla)^r$$

$$x^r = \sum_{b_3=0}^{n_3-1} \left[\sum_{b_2=0}^{n_2-1} \left[\sum_{b_1=0}^{n_1-1} \left[\sum_{b_0=0}^{r-1} \left[\binom{r}{b_0} (((x-b_3n_2n_1n_0)-b_2n_1n_0)-(b_1+1)n_0)^{b_0} n_0^{r-b_0} \right] \right] \right] \right] + (x-n_3n_2n_1n_0)^r$$

if we expand n times

$$x^r = \sum_{b_n=0}^{n_n-1} \left[\dots \sum_{b_4=0}^{n_4-1} \left[\sum_{b_3=0}^{n_3-1} \left[\sum_{b_2=0}^{n_2-1} \left[\sum_{b_1=0}^{n_1-1} \left[\sum_{b_0=0}^{r-1} \left[\right. \right. \right. \right. \right. \right.$$

$$\left. \binom{r}{b_0} ((((\dots(x-b_nn_{n-1}\dots n_3n_2n_1n_0)\dots - b_4n_3n_2n_1n_0) - b_3n_2n_1n_0) - b_2n_1n_0) - (b_1+1)n_0)^{b_0} n_0^{r-b_0} \right.$$

$$\left. \left. \left. \left. \left. \left. \right] \right] \right] \right] \right] \right] + (x-n_n\dots n_4n_3n_2n_1n_0)^r$$

this can be cleaned up if we avoid $((x-a)+a)^r = x^r$ substituion

$$(x+a)^r = \sum_{b_0=0}^{r-1} \left[x^{b_0} a^{r-b_0} \right] + x^r$$

$$x^r = -\sum_{b_0=0}^{r-1} \left[x^{b_0} a^{r-b_0} \right] + (x+a)^r$$

$$x^r = -\sum_{b_0=0}^{r-1} \left[x^{b_0} a^{r-b_0} \right] + \left[-\sum_{b_0=0}^{r-1} \left[(x+a)^{b_0} a^{r-b_0} \right] + ((x+a)+a)^r \right]$$

$$x^r = \sum_{b_1=0}^{n_1-1} \left[-\sum_{b_0=0}^{r-1} \left[(x+b_1a)^{b_0} a^{r-b_0} \right] \right] + ((x+n_1a)^r$$

$$x^r = \sum_{b_1=0}^{n_1-1} \left[-\sum_{b_0=0}^{r-1} \left[(x+b_1n_0)^{b_0} n_0^{r-b_0} \right] \right] + ((x+n_1n_0)^r$$

factoring out the −1 we get:

for all $a \in \mathbb{R} \mid r \in \mathbb{Z}^+ \mid l \in \mathbb{Z}^+$

$$x^r = -1 \sum_{b_n=0}^{n_n-1} \left[\dots \sum_{b_2=0}^{n_2-1} \left[\sum_{b_1=0}^{n_1-1} \left[\sum_{b_0=0}^{r-1} \left[(((x+b_nn_n\dots n_1n_0)+b_2n_1n_0)+b_1n_0)^{b_0} n_0^{r-b_0} \right] \right] \right] + ((x+n_n\dots n_2n_1n_0)^r$$