

Contingency Tables

Contingency Tables

| trigger_type |                        | Response      |        |         | Total |
|--------------|------------------------|---------------|--------|---------|-------|
|              |                        | Accommodation | Reject | Unclear |       |
| C            | Count                  | 1.00          | 17.00  | 16.00   | 34.00 |
|              | Standardized residuals | −0.40         | −1.72  | 1.93    |       |
| F            | Count                  | 0.00          | 24.00  | 9.00    | 33.00 |
|              | Standardized residuals | −1.44         | 1.61   | −1.05   |       |
| T            | Count                  | 3.00          | 20.00  | 9.00    | 32.00 |
|              | Standardized residuals | 1.86          | 0.12   | −0.90   |       |
| Total        | Count                  | 4.00          | 61.00  | 34.00   | 99.00 |
|              |                        |               |        |         |       |

Chi-Squared Tests

|                | Value | df | p    |
|----------------|-------|----|------|
| X <sup>2</sup> | 7.540 | 4  | .110 |
| N              | 99    |    |      |

*Note.* Continuity correction is available only for 2x2 tables.

Nominal

|                 | Value <sup>a</sup> |
|-----------------|--------------------|
| Phi-coefficient | NaN                |
| Cramer's V      | 0.195              |

<sup>a</sup> Phi coefficient is only available for 2 by 2 contingency Tables

| probability |                        | Response      |        |         | Total |
|-------------|------------------------|---------------|--------|---------|-------|
|             |                        | Accommodation | Reject | Unclear |       |
| H           | Count                  | 4.00          | 25.00  | 21.00   | 50.00 |
|             | Standardized residuals | 2.02          | −2.40  | 1.62    |       |
| L           | Count                  | 0.00          | 36.00  | 13.00   | 49.00 |
|             | Standardized residuals | −2.02         | 2.40   | −1.62   |       |
| Total       | Count                  | 4.00          | 61.00  | 34.00   | 99.00 |
|             |                        |               |        |         |       |

Chi-Squared Tests

|                | Value | df | p    |
|----------------|-------|----|------|
| X <sup>2</sup> | 7.857 | 2  | .020 |
| N              | 99    |    |      |

Note. Continuity correction is available only for 2x2 tables.

Nominal

|                 | Value <sup>a</sup> |
|-----------------|--------------------|
| Phi-coefficient | NaN                |
| Cramer's V      | 0.282              |

<sup>a</sup> Phi coefficient is only available for 2 by 2 contingency Tables

Contingency Tables

| embedding_context |                        | Response      |        |         | Total |
|-------------------|------------------------|---------------|--------|---------|-------|
|                   |                        | Accommodation | Reject | Unclear |       |
| N                 | Count                  | 2.00          | 27.00  | 18.00   | 47.00 |
|                   | Standardized residuals | 0.10          | −0.81  | 0.79    |       |
| Q                 | Count                  | 2.00          | 34.00  | 16.00   | 52.00 |
|                   | Standardized residuals | −0.10         | 0.81   | −0.79   |       |
| Total             | Count                  | 4.00          | 61.00  | 34.00   | 99.00 |
|                   |                        |               |        |         |       |

Chi-Squared Tests

|                | Value | df | p    |
|----------------|-------|----|------|
| X <sup>2</sup> | 0.670 | 2  | .715 |
| N              | 99    |    |      |

Note. Continuity correction is available only for 2x2 tables.

Nominal

|                 | Value <sup>a</sup> |
|-----------------|--------------------|
| Phi-coefficient | NaN                |
| Cramer's V      | 0.082              |

<sup>a</sup> Phi coefficient is only available for 2 by 2 contingency Tables

Contingency Tables

|                    |                        | Response      |        |         |       |
|--------------------|------------------------|---------------|--------|---------|-------|
| political_distance |                        | Accommodation | Reject | Unclear | Total |
| C                  | Count                  | 3.00          | 29.00  | 17.00   | 49.00 |
|                    | Standardized residuals | 1.04          | −0.49  | 0.07    |       |
| F                  | Count                  | 1.00          | 32.00  | 17.00   | 50.00 |
|                    | Standardized residuals | −1.04         | 0.49   | −0.07   |       |
| Total              | Count                  | 4.00          | 61.00  | 34.00   | 99.00 |
|                    |                        |               |        |         |       |

Chi-Squared Tests

|                | Value | df | p    |
|----------------|-------|----|------|
| X <sup>2</sup> | 1.138 | 2  | .566 |
| N              | 99    |    |      |

Note. Continuity correction is available only for 2x2 tables.

Nominal

| Value <sup>a</sup> |       |
|--------------------|-------|
| Phi-coefficient    | NaN   |
| Cramer's V         | 0.107 |

<sup>a</sup> Phi coefficient is only available for 2 by 2 contingency Tables