

# Complex Ecological Networks

1

Mathilde Besson <sup>1,2</sup> Eva Delmas\* <sup>1,2</sup> Timothée Poisot <sup>1,2</sup> Dominique Gravel <sup>2,3</sup>

2

**1:** Université de Montréal, Département de Sciences Biologiques; **2:** Québec Centre for Biodiversity Sciences;

3

**3:** Université de Sherbrooke, Département de Biologie

4

Introduction to ecological network

5

**Keywords:** Assemblage - Biodiversity Ecosystem Function - Communities - Ecological mechanisms - Graph theory -  
Network structure - Regional species pool - Stability - -

6

7



8

*This work is licensed under a Creative Commons Attribution 4.0 Unported License.*

9

10

February 23, 2018

11

# 1 Glossary

Adjacency matrix: square matrix representing species interactions. If two species $i$ and $j$ interact, the intersection of the matrix at $i, j$ will be 1, and if no interaction.	13 14
Assembly rules: Ecological processes that lead to a specific species' composition of a community, <i>e.g.</i> competition, predator-prey interactions, arrival history, etc.	15 16
<b>Bipartite / Unipartite network</b> The entire representation of the adjacency matrix offers an <i>unipartite</i> network representation, where the hierarchy between nodes and their position into the network is not always visible ( <i>Figure 1</i> ). On the contrary, a bipartite or k-partite network is a hierarchical representation of the network ( <i>Figure 2</i> ), where nodes are separated depending on their position or function into the network ( <i>e.g.</i> pollinator-plant as bipartite network).	17 18 19 20 21
Ecological interactions: Every type of contact between two species that alters the abundance, biomass and/or behavior of one or both species. Interactions can be trophic, mutualistic or antagonistic, directed or undirected, weighted or unweighted.	22 23 24
Ecosystem functioning: Biotic and abiotic processes that regulate ecosystem, allowing energy and matter flux between trophic levels and between ecosystems, <i>e.g.</i> biogeochemical cycles.	25 26
Graph theory: Mathematical framework used to model the relationship between the objects of a network	27
Network structure: General shape of a network. It is commonly measured using connectance, link distribution, general architecture (nestedness and modularity), etc.	28 29
Nodes/Links, Vertices/Edges: Following graph theory, species are represented as nodes (or vertices), and interactions between them are represented by links (or edges).	30 31
Phylogenetic signal: tendency of phylogenetically close species to have similar traits.	32

## 2 Nomenclature 33

$N$  population size 34

$r$  growth rate 35

$A$  adjacency matrix 36

$\alpha$  interaction strength 37

## 3 38

### 3.1 Introduction 39

In ecological systems (*e.g.* communities), interactions between components (*e.g.* species) are organized in ways that are non-random and constrained by sets of rules (see section X). The organization of these interactions drives or changes some properties of the community, such as its stability, productivity, or ability to resist extinctions, which eventually feedback on the system organization. This constant interplay between interaction organization and system functions results in signatures on the system structure (*e.g.* invariance in key features of the system structure, see section X). Detecting and analyzing these signatures gives us information on the system itself. Studying the structure of ecological systems is necessary to gain insights on the fundamental rules and processes that govern ecosystem formation, maintenance and functioning.

The way interactions are organized can be studied by investigating the structure of ecological networks. Designed to analyze the structure of a studied system, the *graph theory* is a mathematical framework which seemed to be largely appropriate to answer ecological questions. Every system can be abstracted by a *graph*, a representation of the system components organization (*Figure 1a*). These components are called *nodes* and are linked together by *edges*. These combinations form the structure of the studied system. In an ecological system, nodes can be individuals, populations, communities or landscape patches and edges can represent trophic links, energetic flux, etc, every kind of interaction. Both nodes and edges can carry additional information such as weight (*e.g.* species abundance, intensity of the gene flow between two populations, etc.), location in space and

time, edges direction and sign and nodes labels (*e.g.* species identity). Specific informations can be attached to edges, modifying the characteristics of the graph. Graphs can be *directed* (*i.e* interaction goes from A to B) or *undirected*, *weighted* (*i.e* different strength of interaction among the network) or *unweighted* (Figures 1 and 2). This information is summarized mathematically in the *adjacency matrix* (typically named *A*, see definition in glossary) (Figure 1b). In this paper, for simplicity, time and space, we decided to mostly focus on *Species Interaction Networks* (SIN). Ecological system such as landscape, genetic or nutrient networks are not represented here, but they can be studying following the same framework defined further [ref].

Describing and understanding the structure of species interaction networks is an active, and growing, field of ecological research. In this contribution, we will give an overview of some of the most prominent findings and areas of research from the last decade. Starting from a discussion of some invariant properties of the structure of species interaction networks, we will then discuss how this structure affects ecological properties. We will follow by a discussion of the ways ecological networks can be studied under familiar concepts from ecological theory, and finally how this approach scales up to larger temporal or spatial scales.

[Figure 1 about here.]

### 3.2 Invariants in ecological networks

One striking particularity of ecological networks is their consistency: even though they depict interactions between different organisms across all sorts of ecosystems, they all tend to look the same (Jordano, Bascompte and Olesen, 2003). Remarkably, even when interactions among species themselves vary (see section x), the overall network structure tends to remain unchanged (Kemp, Evans, Augustyn and Ellis, 2017). Most ecological networks have a very specific degree distribution (Williams, 2011) (Figure 1d), whereby most species have a small number of interactions, and a small proportions of species have a large number of interactions. In food webs, which represent interactions between preys and their predators, there is a well-described relationship between the number of species and the number of interactions: the number of interactions ( $L$ ) increases proportionally to the number of species ( $S$ ) raised to some exponent, or  $L \propto S^k$ . Martinez (1992) suggested that this exponent is approximately equal to 2, *i.e.* the number of interactions is proportional to the squared number of species. Brose, Ostling, Harrison and Martinez (2004) show that this general relationship holds

even across space: it is possible to estimate how many interactions a species will establish across its entire range. In some other instances, networks may differ on some aspect of their structure, despite obeying to a shared underlying principle. For example, Fortuna, Stouffer and Olesen *et al.* (2010) show that in networks with a low connectance (*Figure 1c*), nestedness (the degree to which the diet of specialists and generalists overlaps – *Figure 2*) and modularity (the tendency of species to form densely aggregated clusters – *Figure 2*) are positively correlated. In networks with higher connectance, this became the opposite: networks with a large number of interactions were either nested (and not modular) or modular (and not nested). In the recent years, it emerged that many aspects of network structure covary with connectance (Poisot and Gravel, 2014; Chagnon, 2015): this suggests that simply knowing how many species there are, and how many interactions they establish, is already very informative about the network structure.

[Figure 2 about here.]

Another remarkable generality of network structure is the distribution of particular shapes of interconnection between three-species subsets. Milo (2002) found that networks (not just ecological but other types of networks such as neuronal or electronical networks as well) can be characterized by the over or under representation of some of these three-species subset, which they called motifs (*Figure 1e*). Motifs can be more broadly defined as being particular shapes of interconnection between three or more nodes in networks at a frequency significantly higher than those found in randomized networks. Three-species motifs represent the simplest building blocks of networks, and more importantly the typical interaction found in communities. As such, they offer the possibility to integrate and test theories developed on simple modules (*e.g.* omnivory, McCann, Hastings and Huxel (1998), Holt (1997)) in larger, more realistic networks. Food webs, for example, are characterized by an over representation of linear food chains and omnivory and an under representation of apparent and exploitative competition (Bascompte and Melián, 2005; Camacho, Stouffer and Amaral, 2007). Stouffer and Bascompte (2010) found that this distribution promotes stability in food webs, with over-represented motifs being more stable in isolation and correlated with higher stability in large realistic communities, and conversely. Motifs can also be used to characterize species role in networks. From the 13 different three-species motifs emerge 30 unique positions for species to occupy in these motifs, representing how the species is embedded in its community. The different positions a species will occupy, and the frequency with which it will occupy these different positions in networks are called species motif role (Stouffer, Sales-Pardo, Sirer and Bascompte, 2012).

These roles have been shown to be evolutionary conserved in food webs (Stouffer, Sales-Pardo, Sirer and Bascompte, 2012) and to have less variability in time than expected in host-parasitoids bipartite networks (Baker, Kaartinen, Roslin and Stouffer, 2015).

Another invariant network property lies in species phylogeny. Phylogeny is a key determinant of ecological network structure, allowing the understanding of species position and interactions into the community. Phylogenetically close species indeed inherit traits from their common ancestors (*e.g.* body size, habitat, defensive strategy, metabolic type, phenology), increasing their propensity to interact with the same group of species or with similar species. This conservatism of interactions has been found to hold across different types of interactions such as antagonistic or mutualistic interactions (Fontaine and Thébault, 2015). However, every species role, such as host or parasite in antagonistic interaction, prey or predator in food web and plant or pollinator/seed disperser in mutualistic interaction, do not provide the same links structure. For instance, closely related host tend to share parasites, while closely related parasites, because of competition for resources, tend to have different hosts species (Krasnov, Fortuna and Mouillot *et al.*, 2012). The conservatism of interactions is consequently unequal all over the network. Following the logic that closely related species interact with the same group of species, Rezende, Albert, Fortuna and Bascompte (2009) shown that phylogenetic structure of ecological networks explains almost entirely the formation and composition of modules in the network, and the connections between them. For the same reasons that conservatism of interactions is asymmetrical, the link between phylogenetic signal and module composition is different depending on the species role (Krasnov, Fortuna and Mouillot *et al.*, 2012), and species that are modules connector are usually phylogenetically close. Cattin, Bersier and Banašek-Richter *et al.* (2004) also found, using a niche-hierarchic model, that diet is constrained by the phylogenetic origin of consumers. The nested structure of trophic networks, generated by the diet structuration, is then influenced by the phylogenetic signal of interacting species and the compatibility of their traits. In contrast, the nested structure of mutualistic networks would be a consequence of trait complementary between species (Rezende, Jordano and Bascompte, 2007). For now, mechanisms underlying the nestedness-phylogeny relationship remain to be further investigated. Moreover, because of species plasticity, phylogeny alone does not fully the structure and evolution of ecological networks.

### 3.3 From structure to properties

136

The relationship between ecological network structure and stability remains a contemporary topic of discussion. 137  
First, MacArthur (1955) and Elton (1958) observed in natural ecosystems that diverse communities have a more 138  
stable dynamic than simple ones. Using a mathematical model based on random ecological networks, May 139  
(1972) undermined this hypothesis. Taking into account not only species diversity but also the connectance and 140  
the interaction strengths in the network, he found that, contrarily to the way of thinking at this time, diversity 141  
was destabilizing communities. This kick in the anthill was the beginning of a prolific complexity-stability 142  
debate, highly oriented on trophic networks (McCann, 2000; see Allesina and Tang, 2015). Two different 143  
approaches of the stability have emerged: one based on the general complexity-stability relationship and 144  
dynamics among species in communities and the second one based on the communities ability to resist biotic 145  
and abiotic changes. The both use different notions and definitions of stability, inducing different ways to 146  
study it (see ???). Despite their dissimilarities, these approaches are not totally independent and have allowed 147  
highlighting that species diversity has no direct influence on communities stability. However, the structure of 148  
ecological network such as interactions distribution and strength seems to play a crucial role (Yodzis, 1981). 149  
The links distribution of ecological networks follows a power-law distribution (Montoya and Solé, 2002), 150  
meaning that few species are highly connected to the rest of the community and a large number of species are 151  
weakly connected to others. This organization combined with the myriad of weak interactions found across 152  
ecological networks, called the weak interaction effect (Berlow, 1999), buffers species variations and then 153  
stabilizes the entire community (Bascompte, Melián and Sala, 2005; Jacquet, Moritz and Morissette *et al.*, 154  
2016). Other parameters, such as the predator-prey body-mass ratio (Emmerson and Raffaelli, 2004; Brose, 155  
Jonsson and Berlow *et al.*, 2006) or network architecture (Montoya, Pimm and Solé, 2006; Thébault and 156  
Fontaine, 2010), determine the distribution and strength of interactions and then contribute to the stability of 157  
ecological networks. 158

Perturbations in ecological communities, generally caused by landscape fragmentation, habitat loss, species 159  
loss or invasion, induce a decrease of species diversity. This loss of diversity is explained by extinctions of 160  
species due directly or indirectly to a first species loss (*i.e.* secondary extinctions or cascade of extinctions). 161  
These extinctions are used to measure the robustness of ecological communities and to explore what happen 162  
when a species is removed or changed in a network. The use of dynamic-based models led to highlight the fact 163

that the probability of secondary extinctions increases with the community size (Lundberg, Ranta and Kaitala, 2008), and decreases with the network connectance (Dunne, Williams and Martinez, 2002). Then, the focus on species removal have allowed to understand that the loss of an highly connected species, also called hub, induces a higher rate of secondary extinctions than the loss of a random, weakly connected species (Solé and Montoya, 2001). However, even if a species is weakly directly connected, if it represents a highway of energy-flow in the network (carbon, nitrogen or biomass), its loss will have similar impact in term of secondary extinctions than the loss of an hug (Allesina and Bodini, 2004). The network architecture also affects the community response to perturbations. For instance, thanks to their structural properties (high nestedness and connectance, Jordano, Bascompte and Olesen (2003)), mutualistic networks persist longer than randomly structured networks (Memmott, Waser and Price, 2004 ; Fortuna and Bascompte, 2006). On the other hand, presence of modules in the community structure limits propagation of perturbations across the rest of the network and then secondary extinctions (Stouffer and Bascompte, 2010).

The consequences of the erosion of biodiversity for ecosystem functioning has been for almost three decades a central problematic for ecologists. While the hypothesis that an increase in species diversity results in an increased productivity dates back to Darwin (Darwin, 1859), the emergence of experimental ecology and the shift from observation in natural systems to the quantification of ecology has made possible to develop a quite general theory for what is now called the biodiversity-ecosystem functioning (BEF) relationship. In a trophic group (*i.e.* a group of species that all belong to the same trophic level, *e.g.* producers or herbivores), the loss of diversity results in a loss of efficiency to capture the shared resource compartment (Loreau, 2010) (*e.g.* nutrients for producers, or producers for herbivores). This leading to a decrease in productivity or other index of functioning. Yet, when the trophic group under focus is coupled to other(s), the action of diversity on functioning is more variable (Duffy, Cardinale and France *et al.*, 2007). This makes the BEF relationship unpredictable in real-world communities, composed of several trophic groups that are virtually never differentiable – as intraguild predation and omnivory blurr the frontier between levels. The multiplicity of the factors influencing the BEF relationship calls for a more general framework that allows the integration of the theories developped for trophic group and for simple modules or sub-systems. By mapping transfer of biomass and energy and/or constraints on organism through the different compartments that compose a natural community, ecological networks – and food webs in particular – offer the possibility to perform this integration. Analyses performed on simulated food-webs with unchanged diversity have already shown that interactions, and more specifically



their structure, have a significative influence on functioning (Thebault and Loreau, 2003; Thébault, Huber and Loreau, 2007; Poisot, Mouquet and Gravel, 2013). The structure of interactions translates the distribution of different types of properties important for ecosystem functioning, such as the presence of omnivory, the generality of species, the modularity of the food-web, etc.

### 3.4 Linking interactions to ecological mechanisms

It is worth remembering that ecological interactions are the direct expression of ecological mechanisms. A pollinator is able to effectively reach the nectar in a plant because the traits of the two organisms match, because they have compatible phenologies, and because they occur in the same environment. A virus can infect its host because it is able to attach to the cell surface, effectively penetrate it, and hijack the cellular machinery to its benefit. Interactions that are not allowed because trait values do not match have been called “forbidden links” (Olesen, Bascompte and Dupont *et al.*, 2011). This prompted a search for “linkage rules” (Bartomeus, 2013) in ecological networks, *i.e.* the relationships that must exist between traits borne by two organisms in order for an interaction between them to exist. These can be identified from existing data on traits and interactions (Bartomeus, Gravel and Tylianakis *et al.*, 2016), and then used to generate realistic ecological networks (Crea, Ali and Rader, 2015). González-Varo and Traveset (2016) point out that interactions are happening between individuals: this requires to consider how the traits are distributed at the individual scale, but also how different behaviors may allow organisms to overcome some of the forbidden links. Although traits are an important part of what makes interactions happen, they are only relevant insofar as the organisms are able to encounter one another. The importance of neutral dynamics (*i.e.* how abundances of different species can determine the probability that they can interact, based on how often they would bump into one another by chance) is, somewhat counter-intuitively, great. Canard, Mouquet and Marescot *et al.* (2012) reveals that simulating food web dynamics by using only population abundances to predict interactions yields realistic food webs. In a host-parasite system, local abundances has also been identified as a key predictor of species interactions (Canard, Mouquet and Mouillot *et al.*, 2014). Speaking more broadly, because interactions emerge from all of these ecological mechanisms, there is a need to develop a deeper understanding of their variability (Poisot, Stouffer and Gravel, 2015). Beyond the fundamental advance that this represents, this would allow to predict interactions based on external information (Morales-Castilla, Matias, Gravel and Araújo, 2015).

The realization of an interaction between individuals from the same or different populations within a community 220  
also have ecological consequences as it modifies the dynamics of at least one of the interacting populations, 221  
and through it, emerging properties. If we consider for instance a population A, its dynamics is not the same 222  
when it multiplies in isolation – where it can grow exponentially if resources are unlimited (Malthus, 1798) or 223  
logistically otherwise (Verhulst, 1938) – or when it is embedded in a real-world community, composed of 224  
several species interacting with one another through different mechanisms (Chesson and Kuang, 2008). It can 225  
lose individuals to predation, have parasitism increase its death rate and at the same time see its establishment 226  
eased through facilitation. It then becomes necessary to account for interactions when studying the dynamic 227  
a population, community stability or ecosystem functioning. But the effect of interactions on populations 228  
dynamics is not always straightforward, both in terms of directionality and intensity, as there is different types 229  
of interactions and multiple factors influence their probability of occurrence and strength. Since the seminal 230  
work of May (1972), the analysis of these effects has been a prolific field of ecology, feeding in particular the 231  
famous “complexity-stability debate” (see Allesina and Tang (2015) for an overview). Including interactions in 232  
population dynamics analyses can be done by using model of the following general form: 233

$$\frac{1}{N_i} \frac{d}{dt} N_i = r_i \times \sum_j A_{ij} \alpha_{i,j} N_j$$

wherein the adjacency matrix  $A$  ( $n * n$ ), list the realized interactions in a given community composed of  $n$  234  
species.  $A_{ij} \neq 0$  when species  $i$  and  $j$  interact and 0 otherwise.  $\alpha_{ij}$  quantifies the strength of the interaction. 235  
This equation model populations abundance  $N$  but can easily be adapted to model biomass flows by replacing 236  
populations’ abundances by their biomasses  $B_i$  (see for instance Williams, Brose and Martinez (2007)). 237

Ecological networks are also spatially and temporally variable (Trøjelsgaard and Olesen, 2016). There are two 238  
drivers to this variability: changes in species composition, and changes in the way these species interact (Poisot, 239  
Canard and Mouillot *et al.*, 2012). Changes in species alone are able to generate variation in network properties 240  
(Havens, 1992). Spatial variation in network structure can also reflect deep-time constraints; for example, 241  
Dalsgaard, Trøjelsgaard and Martín González *et al.* (2013) reveal that historical climate change trends have a 242  
signature on the nestedness and modularity of pollination networks. Even when the same species are present, 243  
interactions between them can vary. Carstensen, Sabatino, Trøjelsgaard and Morellato (2014) and Trøjelsgaard, 244  
Jordano, Carstensen and Olesen (2015) investigated this phenomenon in mutualistic networks. Interaction 245

turnover results from variations in partner fidelity (some species pairs are extremely closely associated), but also from variations in the local environment in which the species interact. Interestingly, and as mentioned in section x, networks overwhelmingly tend to conserve their structure even when interactions within them change. Díaz-Castelazo, Guimarães and Jordano *et al.* (2010) surveyed a pollination network over 10 years, and found important species turnover during this period. Nevertheless, the network retained its structure because species were replaced by their functional equivalent; a generalist pollinator often succeeded to another generalist pollinator. Conversely, species tend to retain their role in different communities: Baker, Kaartinen, Roslin and Stouffer (2015) show that species keep occupying the same position in the network across space, regardless of the species they interact with at every location.

### 3.5 From the regional species pool to local structured communities

Describing the different local communities that occur at macroecological scales through their ecological networks represent an additional layer of information compared to simple species lists. As such, ecological networks are a powerful tool to shed new light on the processes underlying species distribution (Cazelles, Araújo, Mouquet and Gravel, 2016). Until recently, the prevailing idea was that at large scales, the role of biotic interactions was very small compared to that of abiotic conditions, and thought to only be important locally (Pearson and Dawson, 2003; Boulangeat, Gravel and Thuiller, 2012). Empirical observations of species-environment relationship are then used to understand species physiological tolerance to environmental conditions and potentially predict their range under different scenarios of climate change (e.g. Araújo, Thuiller and Pearson, 2006). While these climate envelope models provide a useful approximation of species potential distribution (Pearson, Dawson, Berry and Harrison, 2002), there is mounting evidence that biotic interactions – both positive and negative – play a critical role in shaping communities not only at local scales (Boulangeat, Gravel and Thuiller, 2012), but also at macroecological scales (Davis, Lawton, Shorrocks and Jenkinson, 1998; Araújo and Luoto, 2007; Heikkinen, Luoto and Virkkala *et al.*, 2007; Gotelli, Graves and Rahbek, 2010; Araújo, Rozenfeld, Rahbek and Marquet, 2011). So far, the role of interactions in shaping species distribution is mainly estimated from co-occurrence data, used to build joint species distribution models (JSDM) (Pollock, Tingley and Morris *et al.*, 2014). But there are limitations to this approach. For instance, it does not allow to distinguish between co-occurrence caused by biotic interactions and correlated responses to unmeasured environmental variables (Pollock, Tingley and Morris *et al.*, 2014). Conversely, the lack of

association between species is no evidence of absence of interaction (Cazelles, Araújo, Mouquet and Gravel, 2016). To move from empirical-based species distribution models (SDM) toward theory-driven SDM, further work is needed. In particular, developing methods allowing to include prior information about the underlying ecological network when building (J)SDM could help shedding light on the the fundamental processes underlying species distribution and thus making more accurate predictions (Cazelles, Araújo, Mouquet and Gravel, 2016). Additionally, Poisot, Guéveneux-Julien and Fortin *et al.* (2017) recently showed that biotic interactions respond to environmental conditions on their own, independently of species.

Ecological networks also offer an ideal framework to study the conditions for the maintenance of biodiversity in communities through species coexistence. Gause (1934) predicted that species that shared similar ecologies could not live together in the same area. This competitive exclusion principle states that the the strongest competitor will eventually come to dominate the other species and drive them to local extinction. This stands in contradiction with the existence of ecological communities containing species that overlap in some extent in their resources or consumers. Phytoplanktonic communities are a paradigmatic example of this paradox (Hutchinson, 1961), as they exhibit a high biodiversity while species are competing for a limited number of shared resources (e.g. light, nitrate). The use of consumer-resources models has allowed to highlight some mechanisms improving species coexistence (Chesson, 2000). These mechanisms are based on species traits that either decrease fitness differences (equalizing mechanisms) and/or increase niche differentiation between species (stabilizing mechanisms). The coupling of this type of model with the representation of ecological communities as their underlying network of interactions has brought new perspective on species coexistence, as it is allowing to integrate these mechanisms in large realistic communities. Using this methodological framework, Martinez, Williams and Dunne (2006) showed that the global non-random structure of the food webs improved community persistence. The distribution of motifs in food webs (Stouffer and Bascompte, 2010, see section ‘Invariants in ecological networks’) as well as species’ role within motifs (Stouffer, Sales-Pardo, Sirer and Bascompte, 2012) are related to community persistence. In mutualistic networks, the nested structure has been shown to minimize competition relatively to competition (Bastolla, Fortuna and Pascual-García *et al.*, 2009; Sugihara and Ye, 2009). In these networks, the asymmetry of dependences – the fact that when one species *A* depends strongly on another species *B* as resource for food or pollination, the other species (*B*) only weakly depends on *A* – also increase persistence (Bascompte, Jordano and Olesen, 2006). This type of approach also allowed to highlight the interplay between traits and structure. As an example, Brose, Williams

and Martinez (2006) showed that the allometric scaling of metabolic rates of species improve community persistence when the organization of the food webs is such that predator–prey body mass ratios are different from zero.

Ecologists have also questioned the way communities are formed and the hypothetical set of rules embedding this assembly. Diamond (1975) defined emblematic rules to understand community structure and assembly. In this continuity, network framework allows to explore in detail processes influencing ecological communities assembly. Capitán, Cuesta and Bascompte (2009), for instance, have retraced the pathway of the community assembly process through an assembly graph, based on graph theory. It allows to follow step by step every possible path in community assembly from, for instance, 0 to 21 species among 3 trophic levels, and highlight underlying mechanisms. For food webs especially, mechanistic models such as niche model (Williams and Martinez, 2000) and the cascade model (Cohen, 1989) originally constructed to understand networks structure, have actually be used to understand community assembly and the impact of invasion. Using also network framework, (Verdú and Valiente-Banuet, 2008) found that nested community provides generalists species which facilitate the presence of other species into the network. At the same time, thanks to an experimental network study, (Olesen, Bascompte, Elberling and Jordano, 2008) have observed that new arrival species tend to interact more easily with already well-connected or generalist species. These kind of results could let us think about the Drake’s controversial idea Drake (1991) that species arrival history would be a *important* factor driving community assembly (Drake, 1991). Thanks to community network, Campbell, Yang, Albert and Shea (2011) shown that history assembly process is and important factor for mutualistic networks. Community assembly have however, a myriad of different drivers, such as dispersion, interaction strength and phylogeny distance between species composing communities (Montoya and Solé, 2003; Kraft, Cornwell, Webb and Ackerly, 2007; Maherali and Klironomos, 2007; Leibold, Chase and Ernest, 2017). Based on these drivers, distinct types of models have been developed to predict community assembly dynamics (Tilman, 2004; Gravel, Canham, Beaudet and Messier, 2006; Souza, Bezerra and Longhi, 2016). In one hand, niche-based theory models use coexistence theory and niche differentiation. In the other hand, neutral theory models are based on species dynamics (migration, extinction and speciation) under stochastic processes. Theses two types of model are actually complementary, offering processes explanation at the metacommunity level (niche theory) and at the phylogenetic level (neutral theory) [*ref*]. Network framework in community assembly have brought the field one step further and makes links between other ecological fields, such as disassembly prediction (see

Bascompte and Stouffer, 2009) or co-evolutionary processes (Nuismer, Jordano and Bascompte, 2013) much more easier.

### 3.6 Conclusion

As networks and graph theory allowed to understand breakdown into electricity system in United States or the structure and functioning of social network, it is also a powerful tool to investigate ecological questions. As long as the studying system contains interactions, links or connections, the graph theory provides a perfectly adapted simple framework to characterize complicated networks such as ecological networks. Indices such as connectance, degree distribution of network topology serve as basic measurements to describe systems. Using theses indices, this framework facilitates comparison between different ecological networks. And the relatively important number of network studies leads to a myriads of ways to sample, analyze and interpret them (see Delmas, Besson and Brice *et al.*, 2017).

Studying ecological networks have however a larger purpose than just their description and classification. Basic measurements are correlated to several environmental factors and network analysis appears to be helpful in different ecological fields. As we seen through this paper, it can be used to study dynamics of ecological systems and their responses to changes, according to their stability over time or the BEF relationships in the system. It also highlight the understanding of mechanisms underlying ecological properties such as community assembly, coexistence and species distribution. Network studies were a key to reveal relationships between different properties of ecological network such as trait and structure.

### References

- Allesina, S. and Bodini, A. (2004) ‘Who dominates whom in the ecosystem? Energy flow bottlenecks and cascading extinctions’, *Journal of Theoretical Biology*, 230(3), pp. 351–358. doi: 10.1016/j.jtbi.2004.05.009.
- Allesina, S. and Tang, S. (2015) ‘The stability–complexity relationship at age 40: a random matrix perspective’, *Population Ecology*, 57(1), pp. 63–75. doi: 10.1007/s10144-014-0471-0.
- Araújo, M. B. and Luoto, M. (2007) ‘The importance of biotic interactions for modelling species distribu-

- tions under climate change', *Global Ecology and Biogeography*, 16(6), pp. 743–753. doi: 10.1111/j.1466- 356  
8238.2007.00359.x. 357
- Araújo, M. B., Rozenfeld, A., Rahbek, C. and Marquet, P. A. (2011) 'Using species co-occurrence networks to 358  
assess the impacts of climate change', *Ecography*, 34(6), pp. 897–908. doi: 10.1111/j.1600-0587.2011.06919.x. 359
- Araújo, M. B., Thuiller, W. and Pearson, R. G. (2006) 'Climate warming and the decline of amphibians and 360  
reptiles in Europe', *Journal of Biogeography*, 33(10), pp. 1712–1728. doi: 10.1111/j.1365-2699.2006.01482.x. 361
- Baker, N. J., Kaartinen, R., Roslin, T. and Stouffer, D. B. (2015) 'Species' roles in food webs show fidelity 362  
across a highly variable oak forest', *Ecography*, 38(2), pp. 130–139. doi: 10.1111/ecog.00913. 363
- Bartomeus, I. (2013) 'Understanding linkage rules in plant-pollinator networks by using hierarchical models 364  
that incorporate pollinator detectability and plant traits', *PloS one*, 8(7), p. e69200. Available at: <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0069200> (Accessed: 21 February 2017). 365  
366
- Bartomeus, I., Gravel, D. and Tylianakis, J. M. *et al.* (2016) 'A common framework for identifying linkage 367  
rules across different types of interactions', *Functional Ecology*, 30(12), pp. 1894–1903. Available at: <http://onlinelibrary.wiley.com/doi/10.1111/1365-2435.12666/full> (Accessed: 21 February 2017). 368  
369
- Bascompte, J. and Melián, C. J. (2005) 'Simple trophic modules for complex food webs', *Ecology*, 86(11), pp. 370  
2868–2873. doi: 10.1890/05-0101. 371
- Bascompte, J. and Stouffer, D. B. (2009) 'The assembly and disassembly of ecological networks', *Philosophical 372  
Transactions of the Royal Society B: Biological Sciences*, 364(1524), pp. 1781–1787. doi: 10.1098/rstb.2008.0226. 373
- Bascompte, J., Jordano, P. and Olesen, J. M. (2006) 'Asymmetric Coevolutionary Networks Facilitate Biodiver- 374  
sity Maintenance', *Science*, 312(5772), pp. 431–433. doi: 10.1126/science.1123412. 375
- Bascompte, J., Melián, C. J. and Sala, E. (2005) 'Interaction strength combinations and the overfishing of a 376  
marine food web', *Proceedings of the National Academy of Sciences of the United States of America*, 102(15), 377  
pp. 5443–5447. doi: 10.1073/pnas.0501562102. 378
- Bastolla, U., Fortuna, M. A. and Pascual-García, A. *et al.* (2009) 'The architecture of mutualistic networks min- 379

imizes competition and increases biodiversity’, *Nature*, 458(7241), pp. 1018–1020. doi: 10.1038/nature07950. 380

Berlow, E. L. (1999) ‘Strong effects of weak interactions in ecological communities’, *Nature*, 398(6725), pp. 381  
330–334. doi: 10.1038/18672. 382

Boulangeat, I., Gravel, D. and Thuiller, W. (2012) ‘Accounting for dispersal and biotic interactions to disen- 383  
tangle the drivers of species distributions and their abundances’, *Ecology Letters*, 15(6), pp. 584–593. doi: 384  
10.1111/j.1461-0248.2012.01772.x. 385

Brose, U., Jonsson, T. and Berlow, E. L. *et al.* (2006) ‘Consumer–Resource Body-Size Relationships in Natural 386  
Food Webs’, *Ecology*, 87(10), pp. 2411–2417. doi: 10.1890/0012-9658(2006)87[2411:CBRINF]2.0.CO;2. 387

Brose, U., Ostling, A., Harrison, K. and Martinez, N. D. (2004) ‘Unified spatial scaling of species and their 388  
trophic interactions’, *Nature*, 428(6979), pp. 167–171. doi: 10.1038/nature02297. 389

Brose, U., Williams, R. J. and Martinez, N. D. (2006) ‘Allometric scaling enhances stability in complex food 390  
webs’, *Ecology Letters*, 9(11), pp. 1228–1236. doi: 10.1111/j.1461-0248.2006.00978.x. 391

Camacho, J., Stouffer, D. and Amaral, L. (2007) ‘Quantitative analysis of the local structure of food webs’, 392  
*Journal of Theoretical Biology*, 246(2), pp. 260–268. doi: 10.1016/j.jtbi.2006.12.036. 393

Campbell, C., Yang, S., Albert, R. and Shea, K. (2011) ‘A network model for plant–pollinator community assem- 394  
bly’, *Proceedings of the National Academy of Sciences*, 108(1), pp. 197–202. doi: 10.1073/pnas.1008204108. 395

Canard, E. F., Mouquet, N. and Mouillot, D. *et al.* (2014) ‘Empirical evaluation of neutral interactions in 396  
host-parasite networks’, *The American Naturalist*, 183(4), pp. 468–479. doi: 10.1086/675363. 397

Canard, E., Mouquet, N. and Marescot, L. *et al.* (2012) ‘Emergence of Structural Patterns in Neutral Trophic 398  
Networks’, *PLOS ONE*, 7(8), p. e38295. doi: 10.1371/journal.pone.0038295. 399

Capitán, J. A., Cuesta, J. A. and Bascompte, J. (2009) ‘Statistical Mechanics of Ecosystem Assembly’, *Physical 400  
Review Letters*, 103(16), p. 168101. doi: 10.1103/PhysRevLett.103.168101. 401

Carstensen, D. W., Sabatino, M., Trøjelsgaard, K. and Morellato, L. P. C. (2014) ‘Beta Diversity of Plant- 402  
Pollinator Networks and the Spatial Turnover of Pairwise Interactions’, *PLoS ONE*, 9(11), p. e112903. doi: 403



10.1371/journal.pone.0112903.

404

Cattin, M.-F., Bersier, L.-F. and Banašek-Richter, C. *et al.* (2004) ‘Phylogenetic constraints and adaptation explain food-web structure’, *Nature*, 427(6977), pp. 835–839. doi: 10.1038/nature02327.

405  
406

Cazelles, K., Araújo, M. B., Mouquet, N. and Gravel, D. (2016) ‘A theory for species co-occurrence in interaction networks’, *Theoretical Ecology*, 9(1), pp. 39–48. doi: 10.1007/s12080-015-0281-9.

407  
408

Chagnon, P.-L. (2015) ‘Characterizing topology of ecological networks along gradients: the limits of metrics’ standardization’, *Ecological Complexity*, 22, pp. 36–39. Available at: <http://www.sciencedirect.com/science/article/pii/S1476945X15000070> (Accessed: 21 February 2017).

409  
410  
411

Chesson, P. (2000) ‘Mechanisms of Maintenance of Species Diversity’, *Annual Review of Ecology and Systematics*, 31(1), pp. 343–366. doi: 10.1146/annurev.ecolsys.31.1.343.

412  
413

Chesson, P. and Kuang, J. J. (2008) ‘The interaction between predation and competition’, *Nature*, 456(7219), pp. 235–238. doi: 10.1038/nature07248.

414  
415

Cohen, J. E. (1989) ‘Food webs and community structure’, *Perspectives in ecological theory*, pp. 181–202.

416

Crea, C., Ali, R. A. and Rader, R. (2015) ‘A new model for ecological networks using species-level traits’, *Methods in Ecology and Evolution*. Available at: <http://onlinelibrary.wiley.com/doi/10.1111/2041-210X.12471/pdf> (Accessed: 21 February 2017).

417  
418  
419

Dalsgaard, B., Trøjelsgaard, K. and Martín González, A. M. *et al.* (2013) ‘Historical climate-change influences modularity and nestedness of pollination networks’, *Ecography*, 36(12), pp. 1331–1340. doi: 10.1111/j.1600-0587.2013.00201.x.

420  
421  
422

Darwin, C. (1859) *On the Origin of Species by Means of Natural Selection, or Preservation of Favoured Races in the Struggle for Life*. John Murray, London.

423  
424

Davis, A. J., Lawton, J. H., Shorrocks, B. and Jenkinson, L. S. (1998) ‘Individualistic species responses invalidate simple physiological models of community dynamics under global environmental change’, *Journal*

425  
426

- of Animal Ecology*, 67(4), pp. 600–612. doi: 10.1046/j.1365-2656.1998.00223.x. 427
- Delmas, E., Besson, M. and Brice, M.-H. *et al.* (2017) ‘Analyzing ecological networks of species interactions’, 428  
pp. –. doi: 10.1101/112540. 429
- Diamond, J. M. (1975) ‘Assembly of species communities’, in *Ecology and evolution of communities*. Harvard 430  
University Press, Cambridge, Massachusetts, US: M. L. Cody and J. M. Diamond, editors, pp. 342–444. 431
- Díaz-Castelazo, C., Guimarães, P. R. and Jordano, P. *et al.* (2010) ‘Changes of a mutualistic network over time: 432  
reanalysis over a 10-year period’, *Ecology*, 91(3), pp. 793–801. Available at: [http://www.jstor.org/stable/](http://www.jstor.org/stable/25661111) 433  
**25661111** (Accessed: 21 February 2017). 434
- Drake, J. A. (1991) ‘Community-Assembly Mechanics and the Structure of an Experimental Species Ensemble’, 435  
*The American Naturalist*, 137(1), pp. 1–26. doi: 10.1086/285143. 436
- Duffy, J. E., Cardinale, B. J. and France, K. E. *et al.* (2007) ‘The functional role of biodiversity in ecosystems: 437  
incorporating trophic complexity’, *Ecology Letters*, 10(6), pp. 522–538. doi: 10.1111/j.1461-0248.2007.01037.x. 438
- Dunne, J. A., Williams, R. J. and Martinez, N. D. (2002) ‘Network structure and biodiversity loss in food 439  
webs: robustness increases with connectance’, *Ecology Letters*, 5(4), pp. 558–567. doi: 10.1046/j.1461- 440  
0248.2002.00354.x. 441
- Elton, C. C. (1958) ‘The Reasons for Conservation’, in *The Ecology of Invasions by Animals and Plants*. 442  
Springer Netherlands, pp. 143–153. doi: 10.1007/978-94-009-5851-7\_8. 443
- Emmerson, M. C. and Raffaelli, D. (2004) ‘Predator–prey body size, interaction strength and the stability of a 444  
real food web’, *Journal of Animal Ecology*, 73(3), pp. 399–409. doi: 10.1111/j.0021-8790.2004.00818.x. 445
- Fontaine, C. and Thébault, E. (2015) ‘Comparing the conservatism of ecological interactions in plant–pollinator 446  
and plant–herbivore networks’, *Population Ecology*, 57(1), pp. 29–36. doi: 10.1007/s10144-014-0473-y. 447
- Fortuna, M. A. and Bascompte, J. (2006) ‘Habitat loss and the structure of plant-animal mutualistic net- 448  
works: Mutualistic networks and habitat loss’, *Ecology Letters*, 9(3), pp. 281–286. doi: 10.1111/j.1461- 449

0248.2005.00868.x.

450

Fortuna, M. A., Stouffer, D. B. and Olesen, J. M. *et al.* (2010) ‘Nestedness versus modularity in ecological  
networks: two sides of the same coin?’, *Journal of Animal Ecology*, 79(4), pp. 811–817. doi: 10.1111/j.1365-  
2656.2010.01688.x.

451

452

453

Gause, G. F. (1934) ‘Experimental analysis of Vito Volterra’s mathematical theory of the struggle for existence’,  
*Science*, 79(2036), pp. 16–17. Available at: [http://faculty.jsd.claremont.edu/dmcfarlane/bio146mcfarlane/  
papers/Gause\\_paramecium.pdf](http://faculty.jsd.claremont.edu/dmcfarlane/bio146mcfarlane/papers/Gause_paramecium.pdf) (Accessed: 23 August 2017).

454

455

456

González-Varo, J. P. and Traveset, A. (2016) ‘The Labile Limits of Forbidden Interactions’, *Trends in Ecology  
& Evolution*, 31(9), pp. 700–710. doi: 10.1016/j.tree.2016.06.009.

457

458

Gotelli, N. J., Graves, G. R. and Rahbek, C. (2010) ‘Macroecological signals of species interactions in  
the Danish avifauna’, *Proceedings of the National Academy of Sciences*, 107(11), pp. 5030–5035. doi:  
10.1073/pnas.0914089107.

459

460

461

Gravel, D., Canham, C. D., Beaudet, M. and Messier, C. (2006) ‘Reconciling niche and neutrality: the continuum  
hypothesis’, *Ecology Letters*, 9(4), pp. 399–409. doi: 10.1111/j.1461-0248.2006.00884.x.

462

463

Havens, K. (1992) ‘Scale and structure in natural food webs’, *Science (New York, N.Y.)*, 257(5073), pp.  
1107–1109. doi: 10.1126/science.257.5073.1107.

464

465

Heikkinen, R. K., Luoto, M. and Virkkala, R. *et al.* (2007) ‘Biotic interactions improve prediction of boreal bird  
distributions at macro-scales’, *Global Ecology and Biogeography*, 16(6), pp. 754–763. doi: 10.1111/j.1466-  
8238.2007.00345.x.

466

467

468

Holt, R. D. (1997) ‘Community modules’, A.C. Gange, V.K. Brown (Eds.), *Multitrophic interactions in  
Terrestrial Ecosystems, 6th Symposium of the British Ecological Society, Blackwell Science*, pp. 333–350.

469

470

Hutchinson, G. E. (1961) ‘The Paradox of the Plankton’, *The American Naturalist*, 95(882), pp. 137–145. doi:  
10.1086/282171.

471

472

Jacquet, C., Moritz, C. and Morissette, L. *et al.* (2016) ‘No complexity–stability relationship in empirical

473

- ecosystems', *Nature Communications*, 7, p. 12573. doi: 10.1038/ncomms12573. 474
- Jordano, P., Bascompte, J. and Olesen, J. M. (2003) 'Invariant properties in coevolutionary networks of 475  
plant–animal interactions', *Ecology Letters*, 6(1), pp. 69–81. doi: 10.1046/j.1461-0248.2003.00403.x. 476
- Kemp, J. E., Evans, D. M., Augustyn, W. J. and Ellis, A. G. (2017) 'Invariant antagonistic network structure 477  
despite high spatial and temporal turnover of interactions', *Ecography*, pp. n/a–n/a. doi: 10.1111/ecog.02150. 478
- Kraft, N., Cornwell, W., Webb, C. and Ackerly, D. (2007) 'Trait Evolution, Community Assembly, and the 479  
Phylogenetic Structure of Ecological Communities.', *The American Naturalist*, 170(2), pp. 271–283. doi: 480  
10.1086/519400. 481
- Krasnov, B. R., Fortuna, M. A. and Mouillot, D. *et al.* (2012) 'Phylogenetic Signal in Module Composition and 482  
Species Connectivity in Compartmentalized Host-Parasite Networks', *The American Naturalist*, 179(4), pp. 483  
501–511. doi: 10.1086/664612. 484
- Leibold, M. A., Chase, J. M. and Ernest, S. K. M. (2017) 'Community assembly and the functioning of 485  
ecosystems: how metacommunity processes alter ecosystems attributes', *Ecology*, 98(4), pp. 909–919. doi: 486  
10.1002/ecy.1697. 487
- Loreau, M. (2010) 'Linking biodiversity and ecosystems: towards a unifying ecological theory', *Philosophical 488  
Transactions of the Royal Society B: Biological Sciences*, 365(1537), pp. 49–60. doi: 10.1098/rstb.2009.0155. 489
- Lundberg, P., Ranta, E. and Kaitala, V. (2008) 'Species loss leads to community closure', *Ecology Letters*, 3(6), 490  
pp. 465–468. doi: 10.1111/j.1461-0248.2000.00170.x. 491
- MacArthur, R. (1955) 'Fluctuations of Animal Populations and a Measure of Community Stability', *Ecology*, 492  
36(3), p. 533. doi: 10.2307/1929601. 493
- Maherali, H. and Klironomos, J. N. (2007) 'Influence of Phylogeny on Fungal Community Assembly and 494  
Ecosystem Functioning', *Science*, 316(5832), pp. 1746–1748. doi: 10.1126/science.1143082. 495
- Malthus, T. R. (1798) 'An Essay on the Principle of Population, as it Affects the Future Improvement of 496

- Society: With Remarks on the Speculations of Mr. Godwin, Mr. Condorcet, and Other Writers.’ 497
- Martinez, N. D. (1992) ‘Constant Connectance in Community Food Webs’, *The American Naturalist*, 139(6), 498  
pp. 1208–1218. Available at: <http://www.jstor.org/stable/2462337> (Accessed: 21 February 2017). 499
- Martinez, N. D., Williams, R. J. and Dunne, J. A. (2006) ‘Diversity, complexity, and persistence in large model 500  
ecosystems’, *Ecological networks: linking structure to dynamics in food webs*, pp. 163–185. Available at: [https:// 501  
books.google.com/books?hl=en&lr=&id=YpQRDAAAQBAJ&oi=fnd&pg=PA163&dq=%22spatial+ 502  
e%EF%AC%80ects,+and+evolutionary%22+%22of+diversity+and+complexity+in+the+functioning+ 503  
and+stability+of%22+%22\(Hutchinson+1959\).+However,+the+apparent+inevitability+of+this%22+ 504  
%22diversity/complexity+and+stability+\(for+review,+see+Dunne+et+al.%22+&ots=K4c5g6-yd-& 505  
sig=QXohTQ7JTQ88WJRhNBOYrGb640E](https://books.google.com/books?hl=en&lr=&id=YpQRDAAAQBAJ&oi=fnd&pg=PA163&dq=%22spatial+e%EF%AC%80ects,+and+evolutionary%22+%22of+diversity+and+complexity+in+the+functioning+and+stability+of%22+%22(Hutchinson+1959).+However,+the+apparent+inevitability+of+this%22+%22diversity/complexity+and+stability+(for+review,+see+Dunne+et+al.%22+&ots=K4c5g6-yd-&sig=QXohTQ7JTQ88WJRhNBOYrGb640E) (Accessed: 11 May 2017). 506
- May, R. M. (1972) ‘Will a Large Complex System be Stable?’, *Nature*, 238(5364), pp. 413–414. doi: 507  
10.1038/238413a0. 508
- McCann, K. S. (2000) ‘The diversity–stability debate’, *Nature*, 405(6783), pp. 228–233. doi: 10.1038/35012234. 509
- McCann, K., Hastings, A. and Huxel, G. R. (1998) ‘Weak trophic interactions and the balance of nature’, 510  
*Nature*, 395(6704), pp. 794–798. doi: 10.1038/27427. 511
- Memmott, J., Waser, N. M. and Price, M. V. (2004) ‘Tolerance of pollination networks to species extinc- 512  
tions’, *Proceedings of the Royal Society B: Biological Sciences*, 271(1557), pp. 2605–2611. doi: 10.1098/r- 513  
spb.2004.2909. 514
- Milo, R. (2002) ‘Network Motifs: Simple Building Blocks of Complex Networks’, *Science*, 298(5594), pp. 515  
824–827. doi: 10.1126/science.298.5594.824. 516
- Montoya, J. M. and Solé, R. V. (2002) ‘Small World Patterns in Food Webs’, *Journal of Theoretical Biology*, 517  
214(3), pp. 405–412. doi: 10.1006/jtbi.2001.2460. 518
- Montoya, J. M. and Solé, R. V. (2003) ‘Topological properties of food webs: from real data to community 519  
assembly models’, *Oikos*, 102(3), pp. 614–622. doi: 10.1034/j.1600-0706.2003.12031.x. 520
- Montoya, J. M., Pimm, S. L. and Solé, R. V. (2006) ‘Ecological networks and their fragility’, *Nature*, 442(7100), 521

pp. 259–264. doi: 10.1038/nature04927. 522

Morales-Castilla, I., Matias, M. G., Gravel, D. and Araújo, M. B. (2015) ‘Inferring biotic interactions from 523  
proxies’, *Trends in Ecology & Evolution*, 30(6), pp. 347–356. doi: 10.1016/j.tree.2015.03.014. 524

Nuismer, S. L., Jordano, P. and Bascompte, J. (2013) ‘COEVOLUTION AND THE ARCHITECTURE 525  
OF MUTUALISTIC NETWORKS: COEVOLVING NETWORKS’, *Evolution*, 67(2), pp. 338–354. doi: 526  
10.1111/j.1558-5646.2012.01801.x. 527

Olesen, J. M., Bascompte, J. and Dupont, Y. L. *et al.* (2011) ‘Missing and forbidden links in mutualistic 528  
networks’, *Proceedings of the Royal Society of London B: Biological Sciences*, 278(1706), pp. 725–732. 529  
Available at: <http://rspb.royalsocietypublishing.org/content/278/1706/725.short> (Accessed: 21 February 530  
2017). 531

Olesen, J. M., Bascompte, J., Elberling, H. and Jordano, P. (2008) ‘TEMPORAL DYNAMICS IN A POLLI- 532  
NATION NETWORK’, *Ecology*, 89(6), pp. 1573–1582. doi: 10.1890/07-0451.1. 533

Pearson, R. G. and Dawson, T. P. (2003) ‘Predicting the impacts of climate change on the distribution of 534  
species: are bioclimate envelope models useful?’, *Global Ecology and Biogeography*, 12(5), pp. 361–371. doi: 535  
10.1046/j.1466-822X.2003.00042.x. 536

Pearson, R. G., Dawson, T. P., Berry, P. M. and Harrison, P. A. (2002) ‘SPECIES: A Spatial Evaluation of 537  
Climate Impact on the Envelope of Species’, *Ecological Modelling*, 154(3), pp. 289–300. doi: 10.1016/S0304- 538  
3800(02)00056-X. 539

Poisot, T. and Gravel, D. (2014) ‘When is an ecological network complex? Connectance drives degree 540  
distribution and emerging network properties’, *PeerJ*, 2, p. e251. doi: 10.7717/peerj.251. 541

Poisot, T., Canard, E. and Mouillot, D. *et al.* (2012) ‘The dissimilarity of species interaction networks’, *Ecology 542  
Letters*, 15(12), pp. 1353–1361. doi: 10.1111/ele.12002. 543

Poisot, T., Guéveneux-Julien, C. and Fortin, M.-J. *et al.* (2017) ‘Hosts, parasites and their interactions respond 544  
to different climatic variables: POISOT *et al.*’, *Global Ecology and Biogeography*, 26(8), pp. 942–951. doi: 545

10.1111/geb.12602. 546

Poisot, T., Mouquet, N. and Gravel, D. (2013) 'Trophic complementarity drives the biodiversity-ecosystem 547  
functioning relationship in food webs', *Ecology Letters*, 16(7), pp. 853–861. doi: 10.1111/ele.12118. 548

Poisot, T., Stouffer, D. B. and Gravel, D. (2015) 'Beyond species: why ecological interaction networks vary 549  
through space and time', *Oikos*, 124(3), pp. 243–251. doi: 10.1111/oik.01719. 550

Pollock, L. J., Tingley, R. and Morris, W. K. *et al.* (2014) 'Understanding co-occurrence by modelling species 551  
simultaneously with a Joint Species Distribution Model (JSDM)', *Methods in Ecology and Evolution*, 5(5), pp. 552  
397–406. doi: 10.1111/2041-210X.12180. 553

Rezende, E. L., Albert, E. M., Fortuna, M. A. and Bascompte, J. (2009) 'Compartments in a marine food 554  
web associated with phylogeny, body mass, and habitat structure', *Ecology Letters*, 12(8), pp. 779–788. doi: 555  
10.1111/j.1461-0248.2009.01327.x. 556

Rezende, E. L., Jordano, P. and Bascompte, J. (2007) 'Effects of phenotypic complementarity and phy- 557  
logeny on the nested structure of mutualistic networks', *Oikos*, 116(11), pp. 1919–1929. doi: 10.1111/j.0030- 558  
1299.2007.16029.x. 559

Solé, R. V. and Montoya, M. (2001) 'Complexity and fragility in ecological networks', *Proceedings of the 560  
Royal Society of London B: Biological Sciences*, 268(1480), pp. 2039–2045. doi: 10.1098/rspb.2001.1767. 561

Souza, A. F., Bezerra, A. D. and Longhi, S. J. (2016) 'Quasi-neutral community assembly: Evidence from niche 562  
overlap, phylogenetic, and trait distribution analyses of a subtropical forest in South America', *Perspectives in 563  
Plant Ecology, Evolution and Systematics*, 23, pp. 1–10. doi: 10.1016/j.ppees.2016.09.006. 564

Stouffer, D. B. and Bascompte, J. (2010) 'Understanding food-web persistence from local to global scales', 565  
*Ecology Letters*, 13(2), pp. 154–161. doi: 10.1111/j.1461-0248.2009.01407.x. 566

Stouffer, D. B., Sales-Pardo, M., Sirer, M. I. and Bascompte, J. (2012) 'Evolutionary Conservation of Species' 567  
Roles in Food Webs', *Science*, 335(6075), pp. 1489–1492. doi: 10.1126/science.1216556. 568

Sugihara, G. and Ye, H. (2009) 'Complex systems: Cooperative network dynamics', *Nature*, 458(7241), pp. 569

979–980. doi: 10.1038/458979a. 570

Thebault, E. and Loreau, M. (2003) ‘Food-web constraints on biodiversity-ecosystem functioning relationships’, 571  
*Proceedings of the National Academy of Sciences*, 100(25), pp. 14949–14954. doi: 10.1073/pnas.2434847100. 572

Thébault, E. and Fontaine, C. (2010) ‘Stability of Ecological Communities and the Architecture of Mutualistic 573  
and Trophic Networks’, *Science*, 329(5993), pp. 853–856. doi: 10.1126/science.1188321. 574

Thébault, E., Huber, V. and Loreau, M. (2007) ‘Cascading extinctions and ecosystem functioning: contrasting 575  
effects of diversity depending on food web structure’, *Oikos*, 116(1), pp. 163–173. doi: 10.1111/j.2006.0030- 576  
1299.15007.x. 577

Tilman, D. (2004) ‘Niche tradeoffs, neutrality, and community structure: A stochastic theory of resource 578  
competition, invasion, and community assembly’, *Proceedings of the National Academy of Sciences of the 579  
United States of America*, 101(30), pp. 10854–10861. doi: 10.1073/pnas.0403458101. 580

Trøjelsgaard, K. and Olesen, J. M. (2016) ‘Ecological networks in motion: micro- and macroscopic variability 581  
across scales’, *Functional Ecology*, 30(12), pp. 1926–1935. doi: 10.1111/1365-2435.12710. 582

Trøjelsgaard, K., Jordano, P., Carstensen, D. W. and Olesen, J. M. (2015) ‘Geographical variation in mutualistic 583  
networks: similarity, turnover and partner fidelity’, *Proceedings of the Royal Society B: Biological Sciences*, 584  
282(1802), pp. 20142925–20142925. doi: 10.1098/rspb.2014.2925. 585

Verdú, M. and Valiente-Banuet, A. (2008) ‘The Nested Assembly of Plant Facilitation Networks Prevents 586  
Species Extinctions’, *The American Naturalist*, 172(6), pp. 751–760. doi: 10.1086/593003. 587

Verhulst, P. (1938) ‘Notice sur la loi que la population suit dans son accroissement’, *Curr. Math. Phys*, 10. 588

Williams, R. J. (2011) ‘Biology, Methodology or Chance? The Degree Distributions of Bipartite Ecological 589  
Networks’, *PLoS One*. Edited by J. Langowski, 6(3), p. e17645. doi: 10.1371/journal.pone.0017645. 590

Williams, R. J. and Martinez, N. D. (2000) ‘Simple rules yield complex food webs’, *Nature*, 404(6774), pp. 591  
180–183. doi: 10.1038/35004572. 592

Williams, R. J., Brose, U. and Martinez, N. D. (2007) ‘Homage to Yodzis and Innes 1992: scaling up feeding- 593




based population dynamics to complex ecological networks', in *From energetics to ecosystems: the dynamics* 594  
*and structure of ecological systems*. Springer, pp. 37–51. Available at: [http://link.springer.com/content/pdf/](http://link.springer.com/content/pdf/10.1007/978-1-4020-5337-5_2.pdf) 595  
[10.1007/978-1-4020-5337-5\\_2.pdf](http://link.springer.com/content/pdf/10.1007/978-1-4020-5337-5_2.pdf) (Accessed: 26 May 2016). 596


Yodzis, P. (1981) 'The stability of real ecosystems', *Nature*, 289(5799), pp. 674–676. doi: 10.1038/289674a0. 597

## List of Figures

		598
-1	Figure 1: Graphical representation of a fictional ecological network (a), where species are represented by circles and their directed interactions by arrows. The representation is build from the adjacency matrix (b). In an unipartite representation as this one, each species is represented both as a column and a row. 1 indicates an interaction between two species (e.g. the green square in (b)), and 0 indicates the absence of interaction. This matrix allows to calculate network characteristics such as the connectance (c) and the degree distribution (d). (c) represents the level of connection into the network and is calculated as shown in the figure. (d) represents the distribution of interaction per species. The circles size is relative to the amount of interactions a species have (d1). This distribution is non-random and generally follows a power-law distribution (d2). The network can be split into subnets composed of 3 species, called motif (e). Among the 13 different possible motifs, we only represented the most commonly found in natural communities. . . . .	599 600 601 602 603 604 605 606 607 608 609 27 610
0	Figure 2: Network topology, example of a fictional plant-pollinator network. (a) shows a perfectly nested network, where specialists pollinators are visiting plants embedded into the diet of more generalist pollinators. (b) shows a perfectly modular network, where sub-groups of species interact more strongly with each over than with the rest of the network. (c) shows a random network. Two representations are possibles. Top: Bipartite representation using nodes and edges ; Bottom: Ordered interaction matrix. Here, we used striped yellow squares instead of 1 for presence of interaction and empty squares in absence of interaction. . . . .	611 612 613 614 615 616 28 617



**Figure -1** Figure 1: Graphical representation of a fictional ecological network (a), where species are represented by circles and their directed interactions by arrows. The representation is build from the adjacency matrix (b). In an unipartite representation as this one, each species is represented both as a column and a row. 1 indicates an interaction between two species (e.g. the green square in (b)), and 0 indicates the absence of interaction. This matrix allows to calculate network characteristics such as the connectance (c) and the degree distribution (d). (c) represents the level of connection into the network and is calculated as shown in the figure. (d) represents the distribution of interaction per species. The circles size is relative to the amount of interactions a species have (d1). This distribution is non-random and generally follows a power-law distribution (d2). The network can be split into subnets composed of 3 species, called motif (e). Among the 13 different possible motifs, we only represented the most commonly found in natural communities.



**Figure 0** Figure 2: Network topology, example of a fictional plant-pollinator network. (a) shows a perfectly nested network, where specialists pollinators are visiting plants embedded into the diet of more generalist pollinators. (b) shows a perfectly modular network, where sub-groups of species interact more strongly with each other than with the rest of the network. (c) shows a random network. Two representations are possible. Top: Bipartite representation using nodes and edges; Bottom: Ordered interaction matrix. Here, we used striped yellow squares instead of 1 for presence of interaction and empty squares in absence of interaction.