

# The coevolutionary mosaic of bat-betacoronaviruses spillover risk

Norma Forero Rocio Munoz<sup>1,2</sup>, Renata L. Muylaert<sup>3</sup>, Stephanie N. Seifert<sup>4</sup>, Gregory F. Albery<sup>5</sup>, Colin J. Carlson<sup>5</sup>, Timothée Poisot<sup>1,2,‡</sup>

<sup>1</sup> Université de Montréal; <sup>2</sup> Québec Centre for Biodiversity Sciences; <sup>3</sup> Molecular Epidemiology and Public Health Laboratory, Hopkirk Research Institute, Massey University, Palmerston North, New Zealand; <sup>4</sup> Paul G. Allen School for Global Health, Washington State University, Pullman, WA, United States; <sup>5</sup> ???

‡ These authors contributed equally to the work

## Correspondance to:

Timothée Poisot — timothee.poisot@umontreal.ca

Coming soon

## Keywords:

bats  
betacoronavirus  
disease ecology  
geographic mosaic theory of coevolution  
phylogenetic diversity  
viral sharing  
SARS-CoV-2

Spillover risk is not unidimensional. From the standpoint of an animal community, *i.e.* a pool of suitable hosts, it is driven by a multiplicity of factors (Plowright et al. 2017). The global richness of hosts is one such component commonly mentioned/analysed (see *e.g.* Anthony et al. 2017 for coronaviruses), but there is an argument to be made that species who are not competent (or know) hosts of a specific virus genus may not factor into this (Plowright et al. 2015), or that species who are assumed to share viruses at different rates should be weighted accordingly (Albery et al. 2020). In mammals, key functional traits (for which phylogeny is a reasonable proxy) are determinants of the spillover potential (Olival et al. 2017); these include, notably, body mass and affinity for urban environments (Albery et al. 2022). Finally, especially when the pool of potential hosts spans the entire globe, there may be local host pools that are highly unique; not having been observed in other locations, these can act on the overall risk either by providing novel contact opportunities, reflecting unique host-environment combinations (Engering et al. 2013), or facilitating rapid evolutionary changes in specialism of their pathogens (Agosta et al. 2010). In the specific case of generalist pathogens, there is conceptual and empirical support to the idea that these community- level mechanisms are even more important in driving the overall risk (Power and Mitchell 2004).

Bats are important reservoir hosts for different classes of microorganisms (Chu 2008, Donaldson 2010, Li 2010), some of which can threaten human health. Chiropterans emerged around 64 million years ago and are one of the most diverse mammalian orders, with an estimated richness of more than 14000 species (Peixoto et al. 2018, Simmons and Cirranello 2020). They exhibit a broad variety of habitat use, behaviour, and feeding strategies, resulting in their playing an essential role in the delivery of several ecosystem services tied to important ecosystem-derived benefits (Kasso and Balakrishnan 2013). For example, over two-thirds of bats are known to be either obligate or facultative insectivorous mammals, therefore playing an important role in the regulation of insect pests that can affect crops (Williams-Guillén et al. 2008, Voigt and Kingston 2016), and vectors of diseases that put a risk on human health (Gonsalves et al. 2013a, b). Because bats are globally distributed and have a long evolutionary history, phylogeographic and biogeographic approaches are required to shed light on the extant distribution of coevolutionary processes between bats and the pathogens they carry. Not all areas in which bats, viruses, and humans are co-occurring are facing a risk of spillover towards human populations, and the areas in which this risk exists may not be facing risks of the same nature and magnitude.

In this paper, we examine the biogeographic structure of bats-betacoronaviruses associations, based on a curated dataset of known and recently discovered hosts. This work is important both as a description

of the bats-betacoronavirus complex, but also because more broadly, bats are known reservoirs for a variety of emerging viruses and pathogens (Calisher et al. 2006, Melaun et al. 2014), making balancing the needs for bat conservation and disease prevention a potentially difficult act and a source of human-wildlife conflicts, especially in more densely populated areas (Stone et al. 2015, Rego et al. 2015). By drawing on concepts from the Geographic Mosaic Theory of Coevolution (Thompson 2005), we turn these associations into a spatially explicit additive mapping of zoonotic risk components, which reveals extreme heterogeneity of risk at the global scale; furthermore, we identify the Amazon and South-Eastern Asia as hotspots of phylogenetic distinctiveness of betacoronaviruses (Anthony et al. 2017); surprisingly, current data suggest that viral sharing between hosts is high in the Amazon and low in South-Eastern Asia, which has the potential to result in different evolutionary dynamics between these two regions.

## 1

---

## Methods

**1.1. Known betacoronavirus hosts** We downloaded the data on bats hosts of betacoronaviruses assembled by Becker et al. (2022) from <https://www.viralemergence.org/betacov> on Apr. 2022, and filtered it to “known” hosts (established before the emergence of SARS-CoV-2) and “novel” hosts (confirmed through sampling since the emergence of SARS-CoV-2). The original database was assembled by a combination of data mining and literature surveys, including automated alerts on the “bats” and “coronavirus” keywords to identify novel empirical evidence of bats-betacoronaviruses associations.

**1.2. Bats occurrences** We downloaded the rangemap of every extant bat species that was either classified as an empirically documented host of beta-coronaviruses from the previous step, according to recent IUCN data (IUCN 2021). The range maps were subsequently rasterized using the `rasterize` function from GDAL (Rouault et al. 2022) at a resolution of approximately **TK TP**. For every pixel in the resulting raster where at least one bat host of betacoronavirus was present, we extract the species pool (list of all bat species), which was used to calculate the following risk assessment components: phylogenetic diversity, bat compositional uniqueness, and predicted viral sharing risk.

**1.3. Bats phyogeography** For every pixel, we measured Faith’s Phylogenetic Diversity (Faith 1992) based on a recent synthetic tree with robust time calibration, covering about 6000 mammalian species (Upham et al. 2019). Faith’s PD measures the sum of unique branches from an arbitrary root to a set of tips, and comparatively larger values indicate a more phylogenetic diverse species pool. We measured phylogenetic diversity starting from the root of the entire tree (and not from Chiroptera); this bears no consequences on the resulting values, since all branches leading up to Chiroptera are only counted one per species pool, and (as we explain when describing the assembly of the composite risk map), all individual risk components are ranged in [0,1]. This measure incorporates a richness component, which we chose not to correct for; the interpretation of the phylogenetic diversity is therefore a weighted species richness, that accounts for phylogenetic over/under-dispersal in some places.

**1.4. Bats compositional uniqueness** For every species pool, we measured its Local Contribution to Beta-Diversity (Legendre and De Cáceres 2013); LCBD works from a species-data matrix (traditionally noted as  $\mathbf{Y}$ ), where species are rows and sites are columns, and a value of 1 indicates occurrence. We extracted the  $\mathbf{Y}$  matrix assuming that every pixel represents a unique location, and following best practices (Legendre and Condit 2019) transformed it using Hellinger’s distance to account for unequal bat richness at different pixels. The correction of raw community data is particularly important for two reasons: first, it prevents the artifact of richer sites having higher importance; second, it removes the effect of overall species richness, which is already incorporated in the phylogenetic diversity component. High values of LCBD indicate that the pixel has a community that is on average more dissimilar in species composition than what is expected knowing the entire matrix, i.e. a more unique community. Recent results by Dansereau et al. (2022) shows that LCBD measures are robust with regards to spatial scale, and are therefore applicable at the global scale.

**1.5. Viral sharing between hosts** For all bat hosts of betacoronaviruses, we extracted their predicted viral sharing network (Albery et al. 2020). This network stores pairwise values of viral community

similarity. To project viral sharing values into a single value for every pixel, we averaged the pairwise scores. High values of the average sharing propensity means that this specific extant bat assemblage is likely to be proficient at exchanging viruses.

**1.6. Composite risk map** To visualize the aggregated risk at the global scale, we combine the three individual risk components (phylogenetic diversity, compositional uniqueness, and viral sharing) using an additive color model (Seekell et al. 2018). In this approach, every risk component gets assigned a component in the RGB color model (phylogenetic diversity is green, compositional uniqueness is red, and viral sharing is blue). In order to achieve a valid RGB measure, all components are re-scaled to the [0,1] interval, so that a pixel with no sharing, no phylogenetic diversity, and no compositional uniqueness is black, and a pixel with maximal values for each is white. This additive model conveys both the intensity of the overall risk, but also the nature of the risk as colors diverge towards combinations of values for three risk components. Out of the possible combinations, the most risky in terms of rapid diversification and spillover potential is high phylogenetic diversity and low viral sharing (Gomulkiewicz et al. 2000, Cavender-Bares et al. 2009), in that this allows multiple independent host-virus coevolutionary dynamics to take place in the same location. In the colorimetric space, this correspond to yellow – because the HSV space is more amenable to calculations for feature extraction (see e.g. Keke et al. 2010), we measured the risk level by calculating the angular distance of the hue of each pixel to a reference value of 60, and weighted this risk level by the value component. Specifically, given a pixel with colorimetric coordinates  $(h, s, v)$ , its ranged weighted risk value is

$$v \times \left[ 1 - \frac{|\text{atan}(\cos(\text{rad}(h)), \sin(\text{rad}(h))) - X|}{2\pi} \right],$$

where  $X$  is  $\text{atan}(\cos(\text{rad}(60)), \sin(\text{rad}(60)))$ , a constant approximately equal to 0.5235.

**1.7. Viral phyogeography and evolutionary diversification** We used the following query to pull all betacoronavirus sequence data from the GenBank Nucleotide database except SARS-CoV-2; (“Betacoronavirus”[Organism] OR betacoronavirus[All Fields]) NOT (“Severe acute respiratory syndrome coronavirus 2”[Organism] OR sars-cov-2[All Fields]). We added a single representative sequence for SARS-CoV-2 and manually curated to remove sequences without the RNA-dependent RNA polymerase (RdRp) sequence or that contained words indicating recombinant or laboratory strains including “patent,” “mutant,” “GFP,” and “recombinant.” We filtered over-represented taxa including betacoronavirus 1, hCoV-OC43, Middle East respiratory syndrome coronavirus, Murine hepatitis virus, and hCoV-HKU1. Curated betacoronavirus RdRp sequences were then aligned using MAFFT v 1.4.0 (**Katoh and Standley 2013**, parameters in text?) and a maximum likelihood tree reconstructed in IQ-TREE v 1.6.12 (Nguyen et al. 2015) with ModelFinder (Kalyaanamoorthy et al. 2017) ultrafast bootstrap approximation (Hoang et al. 2018) and the following parameters (**STEPH WILL ADD**, parameters in text?).

We first tested the hypothesis that hotspots of viral diversification would track hotspots of bat diversification. To do so, we plotted the number of known bat hosts (specifically only those included in the phylogeny, so there was a 1:1 correspondence between data sources) against the “mean evolutionary distinctiveness” of the associated viruses. To calculate this, we derived the fair proportions evolutionary distinctiveness (Isaac et al. 2007) for each of the viruses in the tree, then averaged these at the bat species level, projected these values onto their geographic distributions, and averaged across every bat found in a given pixel. As such, this can be thought of as a map of the mean evolutionary distinctiveness of the known viral community believed to be associated with a particular subset of bats present.

**1.8. Co-distribution of hosts and viral hotspots** Subsequently, we tested the hypothesis that the biogeography of bat betacoronaviruses should track the biogeography of their hosts. To test this idea, we loosely adapted a method from (Kreft and Jetz 2007, 2010), who proposed a phylogenetic method for the delineation of animal biogeographic regions. In their original method, a distance matrix - where each row or column represents a geographic raster’s grid cell, and the dissimilarity values are the “beta diversity similarity” of their community assemble - undergoes non-metric multidimensional scaling (NMDS); the first two axes of the NMDS are projected geographically using a four-color bivariate map. Here, we build on this idea with an entirely novel methodology. First, we measure the phylogenetic distance between the different viruses in the betacoronavirus tree by using the cophenetic function in ape (Paradis and Schliep 2019); subsequently, we take a principal components analysis of that distance matrix (readily interchangeable for NMDS in this case) to project the viral tree into an n-dimensional

space. We then take the first two principal components and, as with the evolutionary distinctiveness analysis, aggregated these to a mean host value and projected them using a four-color bivariate map.

## 2

---

### Results and discussion


**2.1. Host richness does not predict virus distinctiveness** Bats are found worldwide and are both one of the most diverse groups among mammals (**Moratelli & Calisher, 2015**), and one of the main animal reservoir for different strains of betacoronaviruses (Drexler et al. 2014). This has attracted attention to areas where high diversity of bats, and therefore presumably high diversity of betacoronaviruses, can be an important issue for human health (Calisher et al. 2006, Moratelli and Calisher 2015). By overlaying the IUCN rangempas for confirmed bat hosts of betacoronaviruses [fig. 1, top], we see that the main hotspots of host richness are primarily South-Eastern Asia, parts of Southern Europe, and to a lesser extent parts of Africa in the -25-0 range of latitudes. The description of host richness is an important first step towards understanding risk, as previous research (**Anthony et al., 2017; Mollentze & Streicker, 2020**) states that locally diverse bat communities could maintain more viruses and hence, a higher probability of having a pathogen that could represent a risk for human health.

Nevertheless, locally diverse and virus-rich bat communities could represent an increased risk of spillover under climate change through the creation of novel interactions (**Ice ice berg berg**), and therefore the diversity of betacoronavirus strains should similarly be accounted for. In fig. 1 (bottom), we contrast the evolutionary distinctiveness of bats and viruses – this reveals a slightly different portrait than bat richness alone. Chiropterans can be classified, from a macro-evolutionary standpoint, as microchiroptera and macrochiroptera, where macrochiroptera have an older history from an evolutionary perspective compared to macrochiroptera (Teeling et al. 2005, Springer 2013). Specifically, we would expect that the so-called “New World” group of bats, being more evolutionary distinct, would also have evolutionary distinct viruses. Indeed fig. 1 (bottom) reveals it to be the case, and this region harbors a distinct bat-betacoronavirus complex. By contrast, South-Eastern Asia has a lot of non-evolutionary distinct bats, but evolutionary-distinct viruses.


It is noteworthy that outside of South America, viral evolutionary distinctiveness does not accurately tracks host diversity, with some areas having over-distinct viruses (southern China but, oddly, not the rest of southeast Asia). There are a number of likely explanations. First, given the richness of bats in southeast Asia, many betacoronaviruses likely remain to be discovered in this region. Indeed, global predictions by (**Becker?**) highlight that southeast Asia is a likely hotspot of unconfirmed hosts of betacoronaviruses, which would likely result in additional viral discoveries. This idea is unsurprising given the growing realization, around the emergence of SARS-CoV-2, that a unique lineage of similar viruses are widespread in bats but still mostly undescribed. The most distinct bats/betacoronavirus complex is found in South America, a region with a comparatively lower number of hosts; this matches with the isolation through variance of the host group, and may highlight a different co-evolutionary dynamic. Alternatively, this distinctiveness hotspot may be a product of under-sampling: South-America is one of the places where the fewest betacoronaviruses have been discovered (Anthony et al. 2017, Olival et al. 2017, Allen et al. 2017), resulting in sparser phylogenetic tree, thereby artificially inflating distinctiveness. Adding more viruses would bring the distinctiveness of known sequences down.

**2.2. The phylogeographic regions of hosts and their viruses overlap** Despite the difference in evolutionary distinctiveness globally, there are reasons to expect that the phylogeography of bats and betacoronaviruses should show some degree of congruence. High density of hosts sharing the same virus (albeit possibly different strains) can drive or result from evolution of the bat antiviral immune system, resulting in spatially distinct immunological responses, as evidenced in several bat species (Banerjee et al. 2020). Immune characteristics that allow bats to be better adapted to infection by emerging viruses (Gorbunova et al., 2020; Irving et al., 2021) may be related to a wide variety of diets (Jones et al., 2022; Moreno Santillán et al., 2021; Banerjee et al., 2020; Schneeberger et al., 2013), themselves likely to be driven by spatial effect, especially at the local scale – bats, indeed, occupy a variety of environments.


In fig. 2, we show a projection of the phylogeographic signal of bats (top) and viruses (bottom) in space; the distinct groupings (represented by different colors symbolizing positions in the subspace formed by the first two axes of the PCoA) are essentially equivalent between the two groups, and can be coarsely delineated as southeast Asia, Eurasia above a northing of 25, and Africa and south America. These results suggest that, although the evolutionary distinctiveness of the bat/betacoronavirus complex varies




**Figure 1** Top panel: relative diversity of known bat hosts of betacoronaviruses. This map shows that the region with the largest number of possible hosts is South-Eastern Asia. Bottom panel: congruence between the evolutionary distinctiveness of the hosts (grey to blue) and the viruses (grey to red). By contrast to the richness map, this reveals that South America has the most evolutionary distinct hosts *and* viruses, whereas South-Eastern Asia has mostly distinct viruses. This is congruent with known results about New World bats being evolutionary distinct, and suggests that they similarly have distinct viruses.



**Figure 2** This is the legend of the figure...






**Figure 3** This is the legend of the figure...

spatially, the system shows an important degree of spatial consistency, with a reduced number of bioregions. Available information describing the spillover of zoonotic betacoronaviruses of bat origin where data was available before and up through the COVID-19 pandemic puts spillover events of SARS-CoV-2 in Wuhan, China; SARS-CoV in XXX based on the presence of closest known viruses circulating in nature, and a nearby location where serological (antibody) evidence has indicated human exposure to SARS-like viruses (Wang et al. 2018 *Virologica Sinica*); MERS-CoV in XXX based on index cases available from a recently-published compendium of cases (Ramshaw et al. 2019). For the latest event, most if not all index cases are presumed to be camel-to-human transmission, and the precise origin point (if it exists) of MERS-CoV in bats is uncertain. Recent recombinant canine coronavirus spillover events in Haiti ([ref](#)) and Europe ([ref](#)) are not relevant here, as bats' involvement in these cycles of transmission have been supposed to be non-existent.

**2.3. Coevolution-informed spillover risk is different in space** As host richness, joint distinctiveness, or phylogeographic structure suggest that the bat/betacoronavirus complex is globally fragmented enough to give rise to both different levels of risk (as evidenced by the spatial location of spillover events) and different types of co-evolutionary dynamics, we turn to the Geographic Mosaic Theory of Coevolution [REF](#) to provide a measure of risk accounting for multiple processes. In fig. 3, we overlapped three components of spillover risk: viral sharing, *i.e.* the chance that two bats will share viruses overall; Local Contribution to Beta Diversity, *i.e.* the fact that a bat community is compositionally unique compared to the average compositional similarity across the entire system; finally, host phylogenetic diversity, *i.e.* how dispersed the bats in a location are within the tree of life. This approach leads to the definition of broad biogeographic regions of risk, where the same color represents the same type of risk. By way of contrast to figures fig. 1 and fig. 2, these regions do not necessarily overlap with previous spatial partitions of the bat/betacoronavirus complex.

From the perspective of spillover risk, the most important combination of factors is a high phylogenetic diversity of hosts with low viral sharing; this, essentially, means that very different betacoronavirus could co-exist within the same place. This is particularly the case given that betacoronaviruses often evolve and even achieve host shifts through recombination, which requires the co-occurrence of sufficiently distinct viruses to be a major driver of emergence. In Fig. xx, this corresponds to yellow to pale green areas, which are essentially limited to South-Eastern Asia, and to some part of Sub-Saharan Africa. Adopting a geographic mosaic theory perspective on risk, other regions of the world are of lesser concern.

Available data on bat betacoronavirus spillover into humans (TP overlay on the figure) is limited and circumstantial at best for these purposes, but our risk maps suggest that the areas predicted by prior expectations about host biogeography correspond loosely to those where previous emergence events have been recorded. Areas with high bat diversity and high turnover may facilitate the evolutionary radiation of viruses, matching previous findings that the diversification of bat coronaviruses is driven



**Figure 4** This is the legend of the figure...

largely by host shifts (inter-genus or higher levels of cross-species transmission) and, to a lesser degree, cospeciation and sharing (intra-genus cross-species transmission; Anthony et al. 2017). This diversification - while not an actual risk factor for spillover itself - likely increases the random chance of a virus with the raw genomic components required for the potential to infect humans.


**2.4. Human occupancy drives different levels of effective risk globally** Based on the previous result, we extracted the yellow component of the risk map (TP add methods), to provide a single measure of risk varying between 0 and 1. This measure is presented in Fig. xxA. However, this maps the potential risk, which must be weighed by the potential for contacts with humans. As a proxy for this measure, we used the proportion of built/urban land from the EarthEnv dataset: this is a reasonable proxy for the density of humans per unit area, which increases the probability of pathogen spread more widely (Hazarie et al., 2021). Since human activity is required to amplify the frequency of virus encounters and thus create areas of viral amplification, mapping the potential risk against measures of land use is required to generate a more actionable assessment of risk. This map is presented in Fig. xxB. Most of South America and Europe are at low risk, as although densely populated, settlements tend to be in areas with lower potential risk. However, this mapping reveals that South-East Asia, the Indian sub-continent, and parts of sub-Saharan Africa, are at high risk due to the overlap between built areas and bat communities representing more opportunities for cross-species transmission of betacoronaviruses.

3

## Conclusion

Our study focuses largely on the biogeography of hosts. Yet, we know that viruses with high host plasticity, that is, the ability of a given virus to adapt to various taxonomic orders and ecological groups (Kreuder Johnson et al., 2015); are more likely to amplify viral spillover, followed by secondary human-to-human transmission, and geographical spread (Hazarie et al., 2021). High viral host plasticity is an especially important trait for RNA viruses such as betacov (Kreuder Johnson et al., 2015; Haddad et al., 2021). Indeed, our analysis of viral sequences reveals that Latin America is a hotspot of viral distinctiveness, suggesting that this part of the bats-betacov system may be undergoing independent evolutionary dynamics (related species sharing viruses that are different from the rest of the global pool). The other hotspot of viral distinctiveness is S.E. Asia, in which richness is high but sharing is low; this suggests a different type of evolutionary dynamics (unrelated viruses coevolving with evolutionarily distinct hosts, generating high diversity locally).

Driven by the need to understand the ecological factors involved in the emergence of viral pathogens, we spatially mapped bat-betacoronavirus interactions worldwide, using (i) a database of known betacov hosts (Becker et al., 2020), and (ii) range maps for the hosts according to IUCN (IUCN 2021). To



**Figure 5** This is the legend of the figure...

reflect the fact that the risk posed by viruses has many ecological origins, we quantified the phylogenetic diversity of hosts, their compositional uniqueness, and the expected viral sharing. Because these components of risk matter when contrasted to human density, we compared them to a proxy, namely the proportion of each pixel that is covered by urban or built land. This provides a synthetic risk map, allowing to identifying of hotspots where the bat-betacoronavirus system may originate viruses in humans. SE Asia is one of the regions with the highest risk since, according to our results, several of its conditions could increase the risk of transmission of the virus.

Species richness, therefore, is not a sufficient measure of viral risk. This is exemplified in our results, where both South America and South-Eastern Asia have a high species richness of betacov hosts, but only the latter region has a high risk. Specifically, because previous studies propose that Asia is important when it comes to understanding the evolutionary origin of various mammalian taxa (Beard C K, 1988).

There are several factors that drive changes in the diversity of bats (Alves et al., 2018), but human activities' effects on the ecosystem (like modifications of land use) could significantly decrease it. Therefore, it can be suggested that changes in the diversity of betacovs in bats are linked to their biogeographic variation, and human population density and other anthropogenic factors are decisive moderators for its implications in public health. With the increase of contact between humans and potential hosts, we also increase the risk of emergence of novel diseases (Johnson et al., 2020), as previous studies on RNA viruses suggest the importance of host phylogeography at the time of virus dispersal (Gryseels et al., 2017).

This diversity of hosts and how the exchange of viruses occurs between species, is largely affected by the different environmental changes, as the case of sarbecovirus bats reservoirs (Muylaert et al., 2021) where they are affected by the area of the cave or the alteration of the forest, which could result in modifications of host distribution. Additionally, our results highlight the importance of Asia as a betacov hotspot, which is consistent with recent studies (Muylaert et al., 2021), where projections on this area suggest that new future events of sarbecovirus viral exchange might be easily spread among species or humans.

One of these scenarios where interaction between bats and humans can occur can be seed dispersal in tropical agroecosystems. It opens the discussion of whether the fruits thrown by bats not only disperse seeds but could also be a source of indirect interaction between viruses of bat origin and humans (Deshpande et al., 2022). This represents a challenge for conservation strategies and disease ecology since we have areas with potential zoonotic viruses and bat-human interaction. However, it must still be taken into account the quantification of real exposure from several scenarios, where there can be directly or indirectly bat - human interaction.

This probability involves multiple factors, among which the relatedness of hosts (which can make the jumps easier (**Longdon et al., 2011; Mollentze et al., 2020; Wolfe et al., 2007**), and the overall tendency of hosts within a locality to share viruses, which may limit viral diversity because of within-host

competition (Leeks et al., 2018; Sallinen et al., 2020). All things considered, the richness of known betacoronaviruses hosts is not a sufficient predictor of spillover risk, and we should move towards a more coevolutionary understanding of this system.

Considering whether viruses easily adapted to multiple hosts have lower virulence on these hosts, or lower ability to jump to hosts with different immune characteristics, should yield valuable additional predictors for the total risk of spillover.

**Acknowledgements:** We acknowledge that this study was conducted on land within the traditional unceded territory of the Saint Lawrence Iroquoian, Anishinabewaki, Mohawk, Huron-Wendat, and Omàmiwininiwak nations. This work was supported by funding to the Viral Emergence Research Initiative (VERENA) consortium including NSF BII 2021909 and a grant from Institut de Valorisation des Données (IVADO). This research was enabled in part by support provided by Calcul Québec ([www.calculquebec.ca](http://www.calculquebec.ca)) and Compute Canada ([www.computecanada.ca](http://www.computecanada.ca)). NF is funded by the NSERC BIOS<sup>2</sup> CREATE program. TP and NF are funded by the Courtois Foundation. RLM was supported by Bryce Carmine and Anne Carmine (née Percival), through the Massey University Foundation.

---

## References

- Agosta, S. J. et al. 2010. How specialists can be generalists: Resolving the “parasite paradox” and implications for emerging infectious disease. - *Zoologia* (Curitiba) 27: 151–162.
- Albery, G. F. et al. 2020. Predicting the global mammalian viral sharing network using phylogeography. - *Nature Communications* 11: 2260.
- Albery, G. F. et al. 2022. Urban-adapted mammal species have more known pathogens. in press.
- Allen, T. et al. 2017. Global hotspots and correlates of emerging zoonotic diseases. - *Nature Communications* in press.
- Anthony, S. J. et al. 2017. Global patterns in coronavirus diversity. - *Virus Evolution* in press.
- Banerjee, A. et al. 2020. Novel Insights Into Immune Systems of Bats. - *Frontiers in Immunology* 11: 26.
- Becker, D. J. et al. 2022. Optimising predictive models to prioritise viral discovery in zoonotic reservoirs. - *The Lancet Microbe* in press.
- Calisher, C. H. et al. 2006. Bats: Important Reservoir Hosts of Emerging Viruses. - *Clinical Microbiology Reviews* 19: 531–545.
- Cavender-Bares, J. et al. 2009. The merging of community ecology and phylogenetic biology. - *Ecol. Lett.* 12: 693–715.
- Dansereau, G. et al. 2022. Evaluating ecological uniqueness over broad spatial extents using species distribution modelling. - *Oikos* n/a: e09063.
- Drexler, J. F. et al. 2014. Ecology, evolution and classification of bat coronaviruses in the aftermath of SARS. - *Antiviral Research* 101: 45–56.
- Engering, A. et al. 2013. Pathogen–host–environment interplay and disease emergence. - *Emerging Microbes & Infections* 2: e5.
- Faith, D. P. 1992. Conservation evaluation and phylogenetic diversity. - *Biological Conservation* 61: 1–10.
- Gomulkiewicz, R. et al. 2000. Hot Spots, Cold Spots, and the Geographic Mosaic Theory of Coevolution. - *The American Naturalist* 156: 156–174.
- Gonsalves, L. et al. 2013a. Do mosquitoes influence bat activity in coastal habitats? - *Wildlife Research* 40: 10–24.
- Gonsalves, L. et al. 2013b. Mosquito Consumption by Insectivorous Bats: Does Size Matter? - *PLOS ONE* 8: e77183.

- Isaac, N. J. B. et al. 2007. Mammals on the EDGE: Conservation Priorities Based on Threat and Phylogeny. - PLOS ONE 2: e296.
- IUCN 2021. The IUCN Red List of Threatened Species.
- Kasso, M. and Balakrishnan, M. 2013. Ecological and Economic Importance of Bats (Order Chiroptera). - ISRN Biodiversity 2013: e187415.
- Keke, S. et al. 2010. Study on skin color image segmentation used by Fuzzy-c-means arithmetic. - 2010 Seventh International Conference on Fuzzy Systems and Knowledge Discovery 2: 612–615.
- Kreft, H. and Jetz, W. 2007. Global patterns and determinants of vascular plant diversity. - Proceedings of the National Academy of Sciences 104: 5925–5930.
- Kreft, H. and Jetz, W. 2010. A framework for delineating biogeographical regions based on species distributions. - Journal of Biogeography 37: 2029–2053.
- Legendre, P. and De Cáceres, M. 2013. Beta diversity as the variance of community data: Dissimilarity coefficients and partitioning (H Morlon, Ed.). - Ecology Letters 16: 951–963.
- Legendre, P. and Condit, R. 2019. Spatial and temporal analysis of beta diversity in the Barro Colorado Island forest dynamics plot, Panama. - Forest Ecosystems 6: 7.
- Melaun, C. et al. 2014. Bats as Potential Reservoir Hosts for Vector-Borne Diseases. - In: Kliment, S. and Mehlhorn, H. (eds), Bats (Chiroptera) as Vectors of Diseases and Parasites: Facts and Myths. Parasitology Research Monographs. Springer, pp. 25–61.
- Moratelli, R. and Calisher, C. H. 2015. Bats and zoonotic viruses: Can we confidently link bats with emerging deadly viruses? - Memórias do Instituto Oswaldo Cruz 110: 1–22.
- Olival, K. J. et al. 2017. Host and viral traits predict zoonotic spillover from mammals. - Nature 546: 646–650.
- Paradis, E. and Schliep, K. 2019. Ape 5.0: An environment for modern phylogenetics and evolutionary analyses in R. - Bioinformatics 35: 526–528.
- Peixoto, F. P. et al. 2018. A synthesis of ecological and evolutionary determinants of bat diversity across spatial scales. - BMC ecology 18: 18.
- Plowright, R. K. et al. 2015. Ecological dynamics of emerging bat virus spillover. - Proceedings of the Royal Society B: Biological Sciences 282: 20142124.
- Plowright, R. K. et al. 2017. Pathways to zoonotic spillover. - Nature Reviews Microbiology 15: 502–510.
- Power, A. G. and Mitchell, C. E. 2004. Pathogen Spillover in Disease Epidemics. - The American Naturalist 164: S79–S89.
- Rego, K. M. da C. et al. 2015. Assessing human-bat interactions around a protected area in northeastern Brazil. - Journal of Ethnobiology and Ethnomedicine 11: 80.
- Rouault, E. et al. 2022. GDAL/OGR Geospatial Data Abstraction software Library. - Zenodo.
- Seekell, D. A. et al. 2018. A geography of lake carbon cycling. - Limnology and Oceanography Letters 3: 49–56.
- Simmons, N. B. and Cirranello, A. L. 2020. Bat Species of the World: A taxonomic and geographic database.
- Springer, M. S. 2013. Phylogenetics: Bats United, Microbats Divided. - Current Biology 23: R999–R1001.
- Stone, E. et al. 2015. Managing Conflict between Bats and Humans: The Response of Soprano Pipistrelles (*Pipistrellus pygmaeus*) to Exclusion from Roosts in Houses. - PLoS ONE 10: e0131825.
- Teeling, E. C. et al. 2005. A Molecular Phylogeny for Bats Illuminates Biogeography and the Fossil Record. - Science (New York, N.Y.) 307: 580–584.
- Thompson, J. N. 2005. The Geographic Mosaic of Coevolution. - University Of Chicago Press.
- Upham, N. S. et al. 2019. Inferring the mammal tree: Species-level sets of phylogenies for questions in ecology, evolution, and conservation. - PLOS Biology 17: e3000494.

2016. Bats in the Anthropocene: Conservation of Bats in a Changing World (CC Voigt and T Kingston, Eds.). - Springer International Publishing.

Williams-Guillén, K. et al. 2008. Bats Limit Insects in a Neotropical Agroforestry System. - Science 320: 70–70.