

The prediction of species interaction networks is facilitated by graph embedding and transfer learning despite data limitations

Tanya Strydom^{1,2,‡} Salomé Bouskila^{1,‡} Francis Banville^{1,3,2} Ceres Barros⁴ Dominique Caron^{5,2}
Maxwell J Farrell⁶ Marie-Josée Fortin⁶ Victoria Hemming⁷ Benjamin Mercier^{3,2} Laura
J. Pollock^{5,2} Rogini Runghen⁸ Giulio V. Dalla Riva⁹ Timothée Poisot^{1,2,‡}

¹ Département de Sciences Biologiques, Université de Montréal, Montréal, Canada ² Quebec Centre for Biodiversity Science, Montréal, Canada ³ Département de Biologie, Université de Sherbrooke, Sherbrooke, Canada ⁴ Department of Forest Resources Management, University of British Columbia, Vancouver, B.C., Canada ⁵ Department of Biology, McGill University, Montréal, Canada ⁶ Department of Ecology & Evolutionary Biology, University of Toronto, Toronto, Canada ⁷ Department of Forest and Conservation Sciences, University of British Columbia, Vancouver, Canada ⁸ Centre for Integrative Ecology, School of Biological Sciences, University of Canterbury, Canterbury, New Zealand ⁹ School of Mathematics and Statistics, University of Canterbury, Canterbury, New Zealand

‡ These authors contributed equally to the work

Correspondance to:

Timothée Poisot — timothee.poisot@umontreal.ca

1. Metawebs, i.e. networks of potential interactions within a species pool, are a powerful abstraction to understand how large-scales species interaction networks are structured.
2. Because metawebs are typically expressed at large spatial and taxonomic scales, assembling them is a tedious and costly process; predictive methods can help circumvent the limitations in data deficiencies, by providing 'draft' metawebs.
3. One way to improve the predictive ability is to maximize the information used for prediction, by using graph embeddings rather than the list of species interactions. Graph embedding is an emerging field in machine learning that holds great potential for ecological problems.
4. In this perspective, we outline how the challenges associated with inferring metawebs line-up with the advantages of graph embeddings; furthermore, because metawebs are inherently spatial objects, we discuss how the choice of the species pool has consequences on the reconstructed network, but also embeds hypotheses about which human-made boundaries are ecologically meaningful.

1 Being able to infer *potential* interactions could be the catalyst for significant breakthroughs in our ability
2 to start thinking about species interaction networks over large spatial scales (Hortal et al., 2015).
3 Understanding species interactions holds enormous potential to not only understand and more rapidly
4 learn about species interactions and metawebs, but also how changes in management of a single species
5 may impact non-target species. In a recent overview of the field of ecological network prediction, Strydom,
6 Catchen, et al. (2021) identified two challenges of interest to the prediction of interactions at large scales.
7 First, there is a relative scarcity of relevant data in most places globally – paradoxically, this restricts our
8 ability to infer interactions for locations where inference is perhaps the least required (and leaves us
9 unable to make inference in regions without interaction data); second, accurate predictions often demand
10 accurate predictors, and the lack of methods that can leverage small amount of *accurate* data is a serious
11 impediment to our global predictive ability. In most places, our most reliable biodiversity knowledge is
12 that of a species pool (*i.e.* a set of potentially interacting species in a given area): through the analysis of
13 databases like GBIF or IUCN, it is possible to construct a list of species in a region of interest; but inferring
14 the potential interactions between these species is difficult.

15 Following the definition of Dunne (2006), a metaweb is the ecological network analogue to the species
16 pool; specifically, it inventories all *potential* interactions between species for a spatially delimited area (and
17 so captures the γ diversity of interactions). The metaweb is not a prediction of the network at a specific
18 point within the spatial area it covers: it will have a different structure, notably by having a larger
19 connectance (see *e.g.* Wood et al., 2015) and complexity (see *e.g.* Galiana et al., 2022), from any of these
20 local networks. These local networks (which capture the α diversity of interactions) are a subset of the
21 metaweb's species and their interactions, and have been called “metaweb realizations” (Poisot et al., 2015).
22 Differences between local networks and their metawebs are due to chance, species abundance and
23 co-occurrence, local environmental conditions, and local distribution of functional traits, among others.
24 Yet, recent results by Saravia et al. (2021) strongly suggest that the local realizations only respond weakly
25 to local conditions: instead, they reflect constraints inherited by the structure of their metaweb. This
26 establishes the metaweb structure as the core goal of predictive network ecology, as it is a required
27 information to accurately produce downscaled, local predictions.

28 Because the metaweb represents the joint effect of functional, phylogenetic, and macroecological
29 processes (Morales-Castilla et al., 2015), it holds valuable ecological information. Specifically, it is the
30 “upper bounds” on what the composition of the local networks, given the local species pool, can be (see

31 e.g. McLeod et al., 2021); this information can help evaluate the ability of ecological assemblages to
32 withstand the effects of, for example, climate change (Fricke et al., 2022). These local networks may be
33 reconstructed given an appropriate knowledge of local species composition and provide information on
34 the structure of food webs at finer spatial scales. This has been done for example for tree-galler-parasitoid
35 systems (Gravel et al., 2018), fish trophic interactions (Albouy et al., 2019), tetrapod trophic interactions
36 (Braga et al., 2019; O'Connor et al., 2020), and crop-pest networks (Grünig et al., 2020). In this
37 contribution, we highlight the power in viewing (and constructing) metawebs as *probabilistic* objects in
38 the context of rare interactions, discuss how a family of machine learning tools (graph embeddings and
39 transfer learning) can be used to overcome data limitations to metaweb inference, and highlight how the
40 use of metawebs introduces important questions for the field of network ecology.

41 **The metaweb is an inherently probabilistic object**

42 Treating interactions as probabilistic (as opposed to binary) events is a more nuanced and realistic way to
43 represent them. Dallas et al. (2017) suggested that most links in ecological networks are cryptic, *i.e.*
44 uncommon or hard to observe. This argument echoes Jordano (2016): sampling ecological interactions is
45 difficult because it requires first the joint observation of two species, and then the observation of their
46 interaction. In addition, it is generally expected that weak or rare links to be more prevalent in networks
47 than common or rare links (Csermely, 2004), compared to strong, persistent links; this is notably the case
48 in food chains, wherein many weaker links are key to the stability of a system (Neutel et al., 2002). In the
49 light of these observations, we expect to see an over-representation of low-probability interactions under a
50 model that accurately predicts interaction probabilities. Yet the original metaweb definition, and indeed
51 most past uses of metawebs, was based on the presence/absence of interactions. Moving towards
52 *probabilistic* metawebs, by representing interactions as Bernoulli events (see e.g. Poisot et al., 2016), offers
53 the opportunity to weigh these rare interactions appropriately. The inherent plasticity of interactions is
54 important to capture: there have been documented instances of food webs undergoing rapid
55 collapse/recovery cycles over short periods of time (e.g. Pedersen et al., 2017). These considerations
56 emphasize why metaweb predictions should focus on quantitative (preferentially probabilistic)
57 predictions; this should constrain the suite of appropriate models.

58 Yet it is important to recall that a metaweb is intended as a catalogue of all potential interactions, which is

59 then filtered (Morales-Castilla et al., 2015). In a sense, that most ecological interactions are elusive can call
60 for a slightly different approach to sampling: once the common interactions are documented, the effort
61 required in documenting each rare interaction will increase exponentially. Recent proposals suggest that
62 machine learning algorithms, in these situations, can act as data generators (Hoffmann et al., 2019): high
63 quality observational data can generate the core rules underpinning the network structure, and be
64 supplemented with synthetic data coming from predictive models, increasing the volume of information
65 available for inference. Indeed, Strydom, Catchen, et al. (2021) suggested that knowing the metaweb may
66 render the prediction of local networks easier, because it fixes an “upper bound” on which interactions
67 can exist. In this context a probabilistic metaweb represents an aggregation of informative priors on the
68 interactions, elusive information with the potential to boost our predictive ability (Bartomeus et al., 2016).

69 [Figure 1 about here.]

70 **Graph embedding offers promises for the inference of potential** 71 **interactions**

72 Graph embedding (fig. 1) is a varied family of machine learning techniques aiming to transform nodes and
73 edges into a vector space (Arsov & Mirceva, 2019), usually of a lower dimension, whilst maximally
74 retaining key properties of the graph (Yan et al., 2005). Ecological networks are an interesting candidate
75 for the widespread application of embeddings, as they tend to possess a shared structural backbone
76 (Bramon Mora et al., 2018), which hints at structural invariants that can be revealed at lower dimensions.
77 Indeed, food webs are inherently low-dimensional objects, and can be adequately represented with less
78 than ten dimensions (Braga et al., 2019; Eklöf et al., 2013). Simulation results by Botella et al. (2022)
79 suggest that there is no best method to identify architectural similarities between networks, and that
80 multiple approaches need to be tested and compared to the network descriptor of interest. This matches
81 previous, more general results on graph embedding, which suggest that the choice of embedding
82 algorithm matters for the results (Goyal & Ferrara, 2018). In tbl. 1, we present a selection of common
83 graph embedding methods, alongside examples of their use to predict species interactions; most of these
84 methods rely either on linear algebra, or on pseudo-random walks on graphs.

85 One prominent family of approaches we do not discuss in the present manuscript is Graph Neural

86 Networks (GNN; Zhou et al., 2020); GNN are, in a sense, a method to embed a graph into a dense
87 subspace, but belong to the family of deep learning methods, which has its own set of practices (see *e.g.*
88 Goodfellow et al., 2016). An important issue with methods based on deep learning is that because their
89 parameter space is immense, the sample size of the data fed into them must be similarly large (typically
90 thousands of instances). This is a requirement for the model to converge correctly during training, but this
91 assumption is unlikely to be met given the size of datasets about currently available metawebs (or single
92 time/location species interaction networks). This data volume requirement is mostly absent from the
93 techniques we list below. Furthermore, GNN still have some challenges related to their shallow structure,
94 and concerns related to scalability (see Gupta et al., 2021 for a review), which are mostly absent from the
95 methods listed in tbl. 1. Assuming that the uptake of next-generation biomonitoring techniques does
96 indeed deliver larger datasets on species interactions (Bohan et al., 2017), there is a potential for GNN to
97 become an applicable embedding/predictive technique in the coming years.

98 [Table 1 about here.]

99 The popularity of graph embedding techniques in machine learning is more than the search for structural
100 invariants: graphs are discrete objects, and machine learning techniques tend to handle continuous data
101 better. Bringing a sparse graph into a continuous, dense vector space (Xu, 2020) opens up a broader variety
102 of predictive algorithms, notably of the sort that are able to predict events as probabilities (Murphy, 2022).
103 Furthermore, the projection of the graph itself is a representation that can be learned; Runghen et al.
104 (2021), for example, used a neural network to learn the embedding of a network in which not all
105 interactions were known, based on nodes metadata. This example has many parallels in ecology (see
106 fig. 2), in which node metadata can be given by phylogeny or functional traits. Rather than directly
107 predicting biological rules (see *e.g.* Pichler et al., 2020 for an overview), which may be confounded by the
108 sparse nature of graph data, learning embeddings works in the low-dimensional space that maximizes
109 information about the network structure. This approach is further justified by the observation, for
110 example, that the macro-evolutionary history of a network is adequately represented by some graph
111 embeddings (RDPG; see Dalla Riva & Stouffer, 2016). In a recent publication, Strydom, Bouskila, et al.
112 (2021) have used an embedding (based on RDPG) to project a metaweb of trophic interactions between
113 European mammals, and transferred this information to mammals of Canada by using the phylogenetic
114 distance between related clades to infer the values in the latent sub-space into which the metaweb was

115 projected. By performing the RDPG step on re-constructed value, this approach yields a probabilistic
116 trophic metaweb for mammals of Canada based on knowledge of European species, despite a limited (\approx
117 5%) taxonomic overlap.

118 Graph embeddings *can* serve as a dimensionality reduction method. For example, RDPG (Strydom,
119 Bouskila, et al., 2021) and t-SVD (Poisot et al., 2021) typically embed networks using fewer dimensions
120 than the original network (the original network has as many dimensions as species, and as many
121 informative dimensions as trophically unique species; Strydom, Dalla Riva, et al., 2021). But this is not
122 necessarily the case – indeed, one may perform a PCA (a special case of SVD) to project the raw data into
123 a subspace that improves the efficacy of t-SNE (Maaten, 2009). There are many dimensionality reductions
124 (Anowar et al., 2021) that can be applied to an embedded network should the need for dimensionality
125 reduction (for example for data visualisation) arise. In brief, many graph embeddings *can* serve as
126 dimensionality reduction steps, but not all do, neither do all dimensionality reduction methods provide
127 adequate graph embedding capacities.

128 [Figure 2 about here.]

129 **An illustration of metaweb embedding**

130 In this section, we illustrate the embedding of a collection of bipartite networks collected by **REF**, using
131 truncated Singular Value Decomposition (t-SVD). The code to reproduce this example (as a Jupyter
132 notebook) is available as supplementary material. The resulting (binary) metaweb \mathcal{M} has 2131 interactions
133 between 206 parasites and 121 hosts (the same metaweb was used for the illustration of interaction
134 inference through deep learning in Strydom, Catchen, et al., 2021), and its adjacency matrix has full rank
135 (*i.e.* it represents a space with 121 dimensions). All analyses were done using Julia (Bezanson et al., 2017)
136 version 1.7.2, *Makie.jl* (Danisch & Krumbiegel, 2021), and *EcologicalNetworks.jl* (Poisot et al., 2019).

137 **The metaweb embeds both ecological hypotheses and practices**

138 The goal of metaweb inference is to provide information about the interactions between species at a large
139 spatial scale. But as Herbert (1965) rightfully pointed out, “[y]ou can’t draw neat lines around planet-wide

140 problems”; any inference of a metaweb at large scales must contend with several novel, and interwoven,
141 families of problems. In this section, we list some of the most pressing research priorities (*i.e.* problems
142 that can be addressed with subsequent data analysis or simulations), as well as issues related to the
143 application of these methods at the science-policy interface.

144 The first open research problem is the taxonomic and spatial limit of the metaweb to embed and transfer.
145 If the initial metaweb is too narrow in scope, notably from a taxonomic point of view, the chances of
146 finding another area with enough related species (through phylogenetic relatedness or similarity of
147 functional traits) to make a reliable inference decreases; this would likely be indicated by large confidence
148 intervals during estimation of the values in the low-rank space, meaning that the representation of the
149 original graph is difficult to transfer to the new problem. In addition, other problems can arise due to
150 non-overlapping trait distributions in the known and unknown species. Alternatively a metaweb is too
151 large (taxonomically), then the resulting embeddings would need to represent interactions between
152 taxonomic groups that are not present in the new location. This would lead to a much higher variance in
153 the starting dataset, and to under-dispersal in the target dataset, resulting in the potential under or over
154 estimation of the strength of new predicted interactions. The lack of well documented metawebs is
155 currently preventing the development of more concrete guidelines. The question of phylogenetic
156 relatedness and dispersal is notably true if the metaweb is assembled in an area with mostly endemic
157 species (*e.g.* a system that has undergone recent radiation and might not have an analogous system with
158 which to draw knowledge from), and as with every predictive algorithm, there is room for the application
159 of our best ecological judgement. Because this problem relates to dispersal of species in the geographic or
160 phylogenetic space, it can certainly be approached through assessing the performance of embedding
161 transfer in simulated starting/target species pools.

162 The second series of problems relate to determining which area should be used to infer the new metaweb
163 in, as this determines the species pool that must be used. Metawebs can be constructed by assigning
164 interactions in a list of species within geographic boundaries. The upside of this approach is that
165 information at the country level is likely to be required for biodiversity assessments, as countries set goals
166 at the national level (Buxton et al., 2021), and as quantitative instruments are designed to work at these
167 scales (Turak et al., 2017); specific strategies are often enacted at smaller scales, nested within a specific
168 country (Ray et al., 2021). But there is no guarantee that these boundaries are meaningful. In fact, we do
169 not have a satisfying answer to the question of “where does a food web stop?”; the most promising

170 solutions involve examining the spatial consistency of network area relationships (Fortin et al., 2021; see
171 e.g. Galiana et al., 2018, 2019, 2021), which is impossible in the absence of enough information about the
172 network itself. This suggests that inferred metawebs should be further downscaled to allow for *a posteriori*
173 analyses. The methodology for metaweb downscaling is currently limited, and it is likely that the
174 sustained effort to characterize the spatial dependency of food web structure will lead to more prescriptive
175 guidelines about the need for prediction downscaling.

176 The final family of problems relates less to ecological methods than to the praxis of ecological research.
177 Operating under the context of national divisions, in large parts of the world, reflects nothing more than
178 the legacy of settler colonialism, which drives a disparity in available ecological data. Applying any
179 embedding to biased data does not debias them, but instead embeds these very same biases, propagating
180 them to the machine learning models using embeddings to make predictions. Indeed, the use of ecological
181 data is not an apolitical act (Nost & Goldstein, 2021), as data infrastructures tend to be designed to answer
182 questions within national boundaries (therefore placing contingencies on what is available to be
183 embedded), and their use often draws upon and reinforces territorial statecraft. As per Machen & Nost
184 (2021), this is particularly true when the output of “algorithmic thinking” (e.g. relying on machine
185 learning to generate knowledge) can be re-used for governance (e.g. enacting conservation decisions at the
186 national scale). As information on species interaction networks structure is increasingly leveraged as a
187 tool to guide conservation actions (see e.g. recent discussions for food-web based conservation; Eero et al.,
188 2021; Naman et al., 2022; Stier et al., 2017), the need to appraise and correct biases that are unwittingly
189 propagated to algorithms when embedded from the original data is paramount. Predictive approaches
190 deployed at the continental scale, no matter their intent, originate in the framework that contributed to
191 the ongoing biodiversity crisis (Adam, 2014) and reinforced environmental injustice (Choudry, 2013;
192 Domínguez & Luoma, 2020). Particularly on Turtle Island and other territories that were traditionally
193 stewarded by Indigenous people, these approaches should be replaced (or at least guided and framed) by
194 Indigenous principles of land management (Eichhorn et al., 2019; No’kmaq et al., 2021), as part of an
195 “algorithm-in-the-loop” approach. Human-algorithm interactions are notoriously difficult and can yield
196 adverse effect (Green & Chen, 2019; Stevenson & Doleac, 2021), suggesting the need to systematically
197 study them for the specific purpose of biodiversity governance, as well as to improve the algorithmic
198 literacy of decision makers. As we see artificial intelligence/machine learning being increasingly
199 mobilized to generate knowledge that is lacking for conservation decisions (e.g. Lamba et al., 2019;

200 Mosebo Fernandes et al., 2020) and drive policy decisions (Weiskopf et al., 2022), our discussion of these
201 tools need to go beyond the technical and statistical, and into the governance consequences they can have.

202 **Acknowledgements:** We acknowledge that this study was conducted on land within the traditional
203 unceded territory of the Saint Lawrence Iroquoian, Anishinabewaki, Mohawk, Huron-Wendat, and
204 Omàmiwininiwak nations. TP, TS, DC, and LP received funding from the Canadian Institute for Ecology
205 & Evolution. FB is funded by the Institute for Data Valorization (IVADO). TS, SB, and TP are funded by a
206 donation from the Courtois Foundation. CB was awarded a Mitacs Elevate Fellowship no. IT12391, in
207 partnership with fRI Research, and also acknowledges funding from Alberta Innovates and the Forest
208 Resources Improvement Association of Alberta. M-JF acknowledges funding from NSERC Discovery
209 Grant and NSERC CRC. RR is funded by New Zealand's Biological Heritage Ngā Koiora Tuku Iho National
210 Science Challenge, administered by New Zealand Ministry of Business, Innovation, and Employment. BM
211 is funded by the NSERC Alexander Graham Bell Canada Graduate Scholarship and the FRQNT master's
212 scholarship. LP acknowledges funding from NSERC Discovery Grant (NSERC RGPIN-2019-05771). TP
213 acknowledges financial support from NSERC through the Discovery Grants and Discovery Accelerator
214 Supplement programs. MJF is supported by an NSERC PDF and an RBC Post-Doctoral Fellowship

215 **Conflict of interest:** The authors have no conflict interests to disclose

216 **Authors' contributions:** TS, and TP conceived the ideas discussed in the manuscript. All authors
217 contributed to writing and editing the manuscript.

218 **Data availability:** There is no data associated with this manuscript.

219 References

- 220 Adam, R. (2014). *Elephant treaties: The Colonial legacy of the biodiversity crisis*. UPNE.
- 221 Albouy, C., Archambault, P., Appeltans, W., Araújo, M. B., Beauchesne, D., Cazelles, K., Cirtwill, A. R.,
222 Fortin, M.-J., Galiana, N., Leroux, S. J., Pellissier, L., Poisot, T., Stouffer, D. B., Wood, S. A., & Gravel, D.
223 (2019). The marine fish food web is globally connected. *Nature Ecology & Evolution*, 3(8, 8),
224 1153–1161. <https://doi.org/10.1038/s41559-019-0950-y>
- 225 Anowar, F., Sadaoui, S., & Selim, B. (2021). Conceptual and empirical comparison of dimensionality
226 reduction algorithms (PCA, KPCA, LDA, MDS, SVD, LLE, ISOMAP, LE, ICA, t-SNE). *Computer*

227 *Science Review*, 40, 100378. <https://doi.org/10.1016/j.cosrev.2021.100378>

228 Arsov, N., & Mirceva, G. (2019). *Network Embedding: An Overview*. <http://arxiv.org/abs/1911.11726>

229 Bartomeus, I., Gravel, D., Tylianakis, J. M., Aizen, M. A., Dickie, I. A., & Bernard-Verdier, M. (2016). A
 230 common framework for identifying linkage rules across different types of interactions. *Functional*
 231 *Ecology*, 30(12), 1894–1903.
 232 <http://onlinelibrary.wiley.com/doi/10.1111/1365-2435.12666/full>

233 Bezanson, J., Edelman, A., Karpinski, S., & Shah, V. (2017). Julia: A Fresh Approach to Numerical
 234 Computing. *SIAM Review*, 59(1), 65–98. <https://doi.org/10.1137/141000671>

235 Bohan, D. A., Vacher, C., Tamaddoni-Nezhad, A., Raybould, A., Dumbrell, A. J., & Woodward, G. (2017).
 236 Next-Generation Global Biomonitoring: Large-scale, Automated Reconstruction of Ecological
 237 Networks. *Trends in Ecology & Evolution*. <https://doi.org/10.1016/j.tree.2017.03.001>

238 Botella, C., Dray, S., Matias, C., Miele, V., & Thuiller, W. (2022). An appraisal of graph embeddings for
 239 comparing trophic network architectures. *Methods in Ecology and Evolution*, 13(1), 203–216.
 240 <https://doi.org/10.1111/2041-210X.13738>

241 Braga, J., Pollock, L. J., Barros, C., Galiana, N., Montoya, J. M., Gravel, D., Maiorano, L., Montemaggiore,
 242 A., Ficetola, G. F., Dray, S., & Thuiller, W. (2019). Spatial analyses of multi-trophic terrestrial vertebrate
 243 assemblages in Europe. *Global Ecology and Biogeography*, 28(11), 1636–1648.
 244 <https://doi.org/10.1111/geb.12981>

245 Bramon Mora, B., Gravel, D., Gilarranz, L. J., Poisot, T., & Stouffer, D. B. (2018). Identifying a common
 246 backbone of interactions underlying food webs from different ecosystems. *Nature Communications*,
 247 9(1), 2603. <https://doi.org/10.1038/s41467-018-05056-0>

248 Buxton, R. T., Bennett, J. R., Reid, A. J., Shulman, C., Cooke, S. J., Francis, C. M., Nyboer, E. A., Pritchard,
 249 G., Binley, A. D., Avery-Gomm, S., Ban, N. C., Beazley, K. F., Bennett, E., Blight, L. K., Bortolotti, L. E.,
 250 Camfield, A. F., Gadallah, F., Jacob, A. L., Naujokaitis-Lewis, I., ... Smith, P. A. (2021). Key
 251 information needs to move from knowledge to action for biodiversity conservation in Canada.
 252 *Biological Conservation*, 256, 108983. <https://doi.org/10.1016/j.biocon.2021.108983>

253 Chen, D., Xue, Y., Fink, D., Chen, S., & Gomes, C. P. (2017). *Deep Multi-species Embedding*. 3639–3646.
 254 <https://www.ijcai.org/proceedings/2017/509>

255 Chen, H., Perozzi, B., Hu, Y., & Skiena, S. (2017). *HARP: Hierarchical Representation Learning for*
 256 *Networks*. <http://arxiv.org/abs/1706.07845>

257 Choudry, A. (2013). Saving biodiversity, for whom and for what? Conservation NGOs, complicity,
 258 colonialism and conquest in an era of capitalist globalization. In *NGOization: Complicity,*
 259 *contradictions and prospects* (pp. 24–44). Bloomsbury Publishing.

260 Cieslak, M. C., Castelfranco, A. M., Roncalli, V., Lenz, P. H., & Hartline, D. K. (2020). T-Distributed
 261 Stochastic Neighbor Embedding (t-SNE): A tool for eco-physiological transcriptomic analysis. *Marine*
 262 *Genomics*, 51, 100723. <https://doi.org/10.1016/j.margen.2019.100723>

263 Csermely, P. (2004). Strong links are important, but weak links stabilize them. *Trends in Biochemical*
 264 *Sciences*, 29(7), 331–334. <https://doi.org/10.1016/j.tibs.2004.05.004>

265 Dalla Riva, G. V., & Stouffer, D. B. (2016). Exploring the evolutionary signature of food webs' backbones
 266 using functional traits. *Oikos*, 125(4), 446–456. <https://doi.org/10.1111/oik.02305>

267 Dallas, T., Park, A. W., & Drake, J. M. (2017). Predicting cryptic links in host-parasite networks. *PLOS*
 268 *Computational Biology*, 13(5), e1005557. <https://doi.org/10.1371/journal.pcbi.1005557>

269 Danisch, S., & Krumbiegel, J. (2021). Makie.jl: Flexible high-performance data visualization for Julia.
 270 *Journal of Open Source Software*, 6(65), 3349. <https://doi.org/10.21105/joss.03349>

271 Domínguez, L., & Luoma, C. (2020). Decolonising Conservation Policy: How Colonial Land and
 272 Conservation Ideologies Persist and Perpetuate Indigenous Injustices at the Expense of the
 273 Environment. *Land*, 9(3, 3), 65. <https://doi.org/10.3390/land9030065>

274 Dunne, J. A. (2006). The Network Structure of Food Webs. In J. A. Dunne & M. Pascual (Eds.), *Ecological*
 275 *networks: Linking structure and dynamics* (pp. 27–86). Oxford University Press.

276 Eero, M., Dierking, J., Humborg, C., Undeman, E., MacKenzie, B. R., Ojaveer, H., Salo, T., & Köster, F. W.
 277 (2021). Use of food web knowledge in environmental conservation and management of living
 278 resources in the Baltic Sea. *ICES Journal of Marine Science*, 78(8), 2645–2663.
 279 <https://doi.org/10.1093/icesjms/fsab145>

280 Eichhorn, M. P., Baker, K., & Griffiths, M. (2019). Steps towards decolonising biogeography. *Frontiers of*
 281 *Biogeography*, 12(1), 1–7. <https://doi.org/10.21425/F5FBG44795>

282 Eklöf, A., Jacob, U., Kopp, J., Bosch, J., Castro-Urgal, R., Chacoff, N. P., Dalsgaard, B., de Sassi, C., Galetti,

283 M., Guimarães, P. R., Lomáscolo, S. B., Martín González, A. M., Pizo, M. A., Rader, R., Rodrigo, A.,
 284 Tylianakis, J. M., Vázquez, D. P., & Allesina, S. (2013). The dimensionality of ecological networks.
 285 *Ecology Letters*, 16(5), 577–583. <https://doi.org/10.1111/ele.12081>

286 Fortin, M.-J., Dale, M. R. T., & Brimacombe, C. (2021). Network ecology in dynamic landscapes.
 287 *Proceedings of the Royal Society B: Biological Sciences*, 288(1949), rspb.2020.1889, 20201889.
 288 <https://doi.org/10.1098/rspb.2020.1889>

289 Fricke, E. C., Ordonez, A., Rogers, H. S., & Svenning, J.-C. (2022). The effects of defaunation on plants’
 290 capacity to track climate change. *Science*.
 291 <https://www.science.org/doi/abs/10.1126/science.abk3510>

292 Galiana, N., Barros, C., Braga, J., Ficetola, G. F., Maiorano, L., Thuiller, W., Montoya, J. M., & Lurgi, M.
 293 (2021). The spatial scaling of food web structure across European biogeographical regions. *Ecography*,
 294 *n/a*(*n/a*). <https://doi.org/10.1111/ecog.05229>

295 Galiana, N., Hawkins, B. A., & Montoya, J. M. (2019). The geographical variation of network structure is
 296 scale dependent: Understanding the biotic specialization of host–parasitoid networks. *Ecography*,
 297 42(6), 1175–1187. <https://doi.org/10.1111/ecog.03684>

298 Galiana, N., Lurgi, M., Bastazini, V. A. G., Bosch, J., Cagnolo, L., Cazelles, K., Claramunt-López, B., Emer,
 299 C., Fortin, M.-J., Grass, I., Hernández-Castellano, C., Jauker, F., Leroux, S. J., McCann, K., McLeod, A.
 300 M., Montoya, D., Mulder, C., Osorio-Canadas, S., Reverté, S., ... Montoya, J. M. (2022). Ecological
 301 network complexity scales with area. *Nature Ecology & Evolution*, 1–8.
 302 <https://doi.org/10.1038/s41559-021-01644-4>

303 Galiana, N., Lurgi, M., Claramunt-López, B., Fortin, M.-J., Leroux, S., Cazelles, K., Gravel, D., & Montoya,
 304 J. M. (2018). The spatial scaling of species interaction networks. *Nature Ecology & Evolution*, 2(5),
 305 782–790. <https://doi.org/10.1038/s41559-018-0517-3>

306 Goodfellow, I., Bengio, Y., & Courville, A. (2016). *Deep learning*. MIT Press.

307 Goyal, P., & Ferrara, E. (2018). Graph embedding techniques, applications, and performance: A survey.
 308 *Knowledge-Based Systems*, 151, 78–94. <https://doi.org/10.1016/j.knosys.2018.03.022>

309 Gravel, D., Baiser, B., Dunne, J. A., Kopelke, J.-P., Martinez, N. D., Nyman, T., Poisot, T., Stouffer, D. B.,
 310 Tylianakis, J. M., Wood, S. A., & Roslin, T. (2018). Bringing Elton and Grinnell together: A quantitative

framework to represent the biogeography of ecological interaction networks. *Ecography*, 0(0).
<https://doi.org/10.1111/ecog.04006>

Green, B., & Chen, Y. (2019). Disparate Interactions: An Algorithm-in-the-Loop Analysis of Fairness in Risk Assessments. *Proceedings of the Conference on Fairness, Accountability, and Transparency*, 90–99.
<https://doi.org/10.1145/3287560.3287563>

Grover, A., & Leskovec, J. (2016). Node2vec: Scalable Feature Learning for Networks. *Proceedings of the 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*, 855–864.
<https://doi.org/10.1145/2939672.2939754>

Grünig, M., Mazzi, D., Calanca, P., Karger, D. N., & Pellissier, L. (2020). Crop and forest pest metawebs shift towards increased linkage and suitability overlap under climate change. *Communications Biology*, 3(1, 1), 1–10. <https://doi.org/10.1038/s42003-020-0962-9>

Gupta, A., Matta, P., & Pant, B. (2021). Graph neural network: Current state of Art, challenges and applications. *Materials Today: Proceedings*, 46, 10927–10932.
<https://doi.org/10.1016/j.matpr.2021.01.950>

Herbert, F. (1965). *Dune* (1st ed.). Chilton Book Company.

Hinton, G., & Roweis, S. T. (2002). Stochastic neighbor embedding. *NIPS*, 15, 833–840.

Hoffmann, J., Bar-Sinai, Y., Lee, L. M., Andrejevic, J., Mishra, S., Rubinstein, S. M., & Rycroft, C. H. (2019). Machine learning in a data-limited regime: Augmenting experiments with synthetic data uncovers order in crumpled sheets. *Science Advances*, 5(4), eaau6792.
<https://doi.org/10.1126/sciadv.aau6792>

Hortal, J., de Bello, F., Diniz-Filho, J. A. F., Lewinsohn, T. M., Lobo, J. M., & Ladle, R. J. (2015). Seven Shortfalls that Beset Large-Scale Knowledge of Biodiversity. *Annual Review of Ecology, Evolution, and Systematics*, 46(1), 523–549. <https://doi.org/10.1146/annurev-ecolsys-112414-054400>

Jordano, P. (2016). Sampling networks of ecological interactions. *Functional Ecology*, 30(12), 1883–1893.
<https://doi.org/10.1111/1365-2435.12763>

Lamba, A., Cassey, P., Segaran, R. R., & Koh, L. P. (2019). Deep learning for environmental conservation. *Current Biology*, 29(19), R977–R982. <https://doi.org/10.1016/j.cub.2019.08.016>

Maaten, L. van der. (2009). Learning a Parametric Embedding by Preserving Local Structure. *Proceedings*

339 of the Twelfth International Conference on Artificial Intelligence and Statistics, 384–391.
 340 <https://proceedings.mlr.press/v5/maaten09a.html>

341 Machen, R., & Nost, E. (2021). Thinking algorithmically: The making of hegemonic knowledge in climate
 342 governance. *Transactions of the Institute of British Geographers*, 46(3), 555–569.
 343 <https://doi.org/10.1111/tran.12441>

344 McLeod, A., Leroux, S. J., Gravel, D., Chu, C., Cirtwill, A. R., Fortin, M.-J., Galiana, N., Poisot, T., & Wood,
 345 S. A. (2021). Sampling and asymptotic network properties of spatial multi-trophic networks. *Oikos*,
 346 *n/a*(*n/a*). <https://doi.org/10.1111/oik.08650>

347 Melnyk, K., Klus, S., Montavon, G., & Conrad, T. O. F. (2020). GraphKKE: Graph Kernel Koopman
 348 embedding for human microbiome analysis. *Applied Network Science*, 5(1), 96.
 349 <https://doi.org/10.1007/s41109-020-00339-2>

350 Morales-Castilla, I., Matias, M. G., Gravel, D., & Araújo, M. B. (2015). Inferring biotic interactions from
 351 proxies. *Trends in Ecology & Evolution*, 30(6), 347–356.
 352 <https://doi.org/10.1016/j.tree.2015.03.014>

353 Mosebo Fernandes, A. C., Quintero Gonzalez, R., Lenihan-Clarke, M. A., Leslie Trotter, E. F., & Jokar
 354 Arsanjani, J. (2020). Machine Learning for Conservation Planning in a Changing Climate.
 355 *Sustainability*, 12(18, 18), 7657. <https://doi.org/10.3390/su12187657>

356 Murphy, K. P. (2022). *Probabilistic machine learning: An introduction*. MIT Press. probml.ai

357 Naman, S. M., White, S. M., Bellmore, J. R., McHugh, P. A., Kaylor, M. J., Baxter, C. V., Danehy, R. J.,
 358 Naiman, R. J., & Puls, A. L. (2022). Food web perspectives and methods for riverine fish conservation.
 359 *WIREs Water*, *n/a*(*n/a*), e1590. <https://doi.org/10.1002/wat2.1590>

360 Narayanan, A., Chandramohan, M., Venkatesan, R., Chen, L., Liu, Y., & Jaiswal, S. (2017). *Graph2vec:*
 361 *Learning Distributed Representations of Graphs*. <http://arxiv.org/abs/1707.05005>

362 Neutel, A.-M., Heesterbeek, J. A. P., & de Ruiter, P. C. (2002). Stability in Real Food Webs: Weak Links in
 363 Long Loops. *Science*, 296(5570), 1120–1123. <https://doi.org/10.1126/science.1068326>

364 No'kmaq, M., Marshall, A., Beazley, K. F., Hum, J., Joudry, shalan, Papadopoulos, A., Pictou, S., Rabesca,
 365 J., Young, L., & Zurba, M. (2021). “Awakening the sleeping giant”: Re-Indigenization principles for
 366 transforming biodiversity conservation in Canada and beyond. *FACETS*, 6(1), 839–869.

367 Nost, E., & Goldstein, J. E. (2021). A political ecology of data. *Environment and Planning E: Nature and*
368 *Space*, 25148486211043503. <https://doi.org/10.1177/25148486211043503>

369 O'Connor, L. M. J., Pollock, L. J., Braga, J., Ficetola, G. F., Maiorano, L., Martinez-Almoyna, C.,
370 Montemaggiore, A., Ohlmann, M., & Thuiller, W. (2020). Unveiling the food webs of tetrapods across
371 Europe through the prism of the Eltonian niche. *Journal of Biogeography*, 47(1), 181–192.
372 <https://doi.org/10.1111/jbi.13773>

373 Pedersen, E. J., Thompson, P. L., Ball, R. A., Fortin, M.-J., Gouhier, T. C., Link, H., Moritz, C., Nenzen, H.,
374 Stanley, R. R. E., Taranu, Z. E., Gonzalez, A., Guichard, F., & Pepin, P. (2017). Signatures of the
375 collapse and incipient recovery of an overexploited marine ecosystem. *Royal Society Open Science*, 4(7),
376 170215. <https://doi.org/10.1098/rsos.170215>

377 Perozzi, B., Al-Rfou, R., & Skiena, S. (2014). DeepWalk: Online learning of social representations.
378 *Proceedings of the 20th ACM SIGKDD International Conference on Knowledge Discovery and Data*
379 *Mining*, 701–710. <https://doi.org/10.1145/2623330.2623732>

380 Pichler, M., Boreux, V., Klein, A.-M., Schleuning, M., & Hartig, F. (2020). Machine learning algorithms to
381 infer trait-matching and predict species interactions in ecological networks. *Methods in Ecology and*
382 *Evolution*, 11(2), 281–293. <https://doi.org/10.1111/2041-210X.13329>

383 Poisot, T., Belisle, Z., Hoebeke, L., Stock, M., & Szefer, P. (2019). EcologicalNetworks.jl - analysing
384 ecological networks. *Ecography*. <https://doi.org/10.1111/ecog.04310>

385 Poisot, T., Cirtwill, A. R., Cazelles, K., Gravel, D., Fortin, M.-J., & Stouffer, D. B. (2016). The structure of
386 probabilistic networks. *Methods in Ecology and Evolution*, 7(3), 303–312.
387 <https://doi.org/10.1111/2041-210X.12468>

388 Poisot, T., Ouellet, M.-A., Mollentze, N., Farrell, M. J., Becker, D. J., Albery, G. F., Gibb, R. J., Seifert, S. N.,
389 & Carlson, C. J. (2021). *Imputing the mammalian virome with linear filtering and singular value*
390 *decomposition*. <http://arxiv.org/abs/2105.14973>

391 Poisot, T., Stouffer, D. B., & Gravel, D. (2015). Beyond species: Why ecological interaction networks vary
392 through space and time. *Oikos*, 124(3), 243–251. <https://doi.org/10.1111/oik.01719>

393 Ramasamy, D., & Madhow, U. (2015). Compressive spectral embedding: Sidestepping the SVD. In C.
394 Cortes, N. Lawrence, D. Lee, M. Sugiyama, & R. Garnett (Eds.), *Advances in neural information*

processing systems (Vol. 28). Curran Associates, Inc. <https://proceedings.neurips.cc/paper/2015/file/4f6ffe13a5d75b2d6a3923922b3922e5-Paper.pdf>

Ray, J. C., Grimm, J., & Olive, A. (2021). The biodiversity crisis in Canada: Failures and challenges of federal and sub-national strategic and legal frameworks. *FACETS*, 6, 1044–1068. <https://doi.org/10.1139/facets-2020-0075>

Runghen, R., Stouffer, D. B., & Dalla Riva, G. V. (2021). *Exploiting node metadata to predict interactions in large networks using graph embedding and neural networks*. <https://doi.org/10.1101/2021.06.10.447991>

Saravia, L. A., Marina, T. I., Kristensen, N. P., De Troch, M., & Momo, F. R. (2021). Ecological network assembly: How the regional metaweb influences local food webs. *Journal of Animal Ecology*, n/a(n/a). <https://doi.org/10.1111/1365-2656.13652>

Stevenson, M. T., & Doleac, J. L. (2021). *Algorithmic Risk Assessment in the Hands of Humans* [SSRN Scholarly Paper]. <https://doi.org/10.2139/ssrn.3489440>

Stier, A. C., Samhouri, J. F., Gray, S., Martone, R. G., Mach, M. E., Halpern, B. S., Kappel, C. V., Scarborough, C., & Levin, P. S. (2017). Integrating Expert Perceptions into Food Web Conservation and Management. *Conservation Letters*, 10(1), 67–76. <https://doi.org/10.1111/conl.12245>

Strydom, T., Bouskila, S., Banville, F., Barros, C., Caron, D., Farrell, M. J., Fortin, M.-J., Hemming, V., Mercier, B., Pollock, L., Runghen, R., Riva, G. V. D., & Poisot, T. (2021). *Food web reconstruction through phylogenetic transfer of low-rank network representation*. <https://doi.org/10.32942/osf.io/y7sdz>

Strydom, T., Catchen, M. D., Banville, F., Caron, D., Dansereau, G., Desjardins-Proulx, P., Forero-Muñoz, N. R., Higino, G., Mercier, B., Gonzalez, A., Gravel, D., Pollock, L., & Poisot, T. (2021). A roadmap towards predicting species interaction networks (across space and time). *Philosophical Transactions of the Royal Society B: Biological Sciences*, 376(1837), 20210063. <https://doi.org/10.1098/rstb.2021.0063>

Strydom, T., Dalla Riva, G. V., & Poisot, T. (2021). SVD Entropy Reveals the High Complexity of Ecological Networks. *Frontiers in Ecology and Evolution*, 9. <https://doi.org/10.3389/fevo.2021.623141>

Tang, J., Qu, M., Wang, M., Zhang, M., Yan, J., & Mei, Q. (2015). LINE: Large-scale Information Network Embedding. *Proceedings of the 24th International Conference on World Wide Web*, 1067–1077.

423 <https://doi.org/10.1145/2736277.2741093>

424 Turak, E., Brazill-Boast, J., Cooney, T., Drielsma, M., DelaCruz, J., Dunkerley, G., Fernandez, M., Ferrier,
 425 S., Gill, M., Jones, H., Koen, T., Leys, J., McGeoch, M., Mihoub, J.-B., Scanes, P., Schmeller, D., &
 426 Williams, K. (2017). Using the essential biodiversity variables framework to measure biodiversity
 427 change at national scale. *Biological Conservation*, 213, 264–271.
 428 <https://doi.org/10.1016/j.biocon.2016.08.019>

429 Wang, D., Cui, P., & Zhu, W. (2016). Structural Deep Network Embedding. *Proceedings of the 22nd ACM*
 430 *SIGKDD International Conference on Knowledge Discovery and Data Mining*, 1225–1234.
 431 <https://doi.org/10.1145/2939672.2939753>

432 Wang, S., Arroyo, J., Vogelstein, J. T., & Priebe, C. E. (2021). Joint Embedding of Graphs. *IEEE*
 433 *Transactions on Pattern Analysis and Machine Intelligence*, 43(4), 1324–1336.
 434 <https://doi.org/10.1109/TPAMI.2019.2948619>

435 Wardeh, M., Baylis, M., & Blagrove, M. S. C. (2021). Predicting mammalian hosts in which novel
 436 coronaviruses can be generated. *Nature Communications*, 12(1, 1), 780.
 437 <https://doi.org/10.1038/s41467-021-21034-5>

438 Weiskopf, S. R., Harmáčková, Z. V., Johnson, C. G., Londoño-Murcia, M. C., Miller, B. W., Myers, B. J. E.,
 439 Pereira, L., Arce-Plata, M. I., Blanchard, J. L., Ferrier, S., Fulton, E. A., Harfoot, M., Isbell, F., Johnson,
 440 J. A., Mori, A. S., Weng, E., & Rosa, I. M. D. (2022). Increasing the uptake of ecological model results in
 441 policy decisions to improve biodiversity outcomes. *Environmental Modelling & Software*, 149, 105318.
 442 <https://doi.org/10.1016/j.envsoft.2022.105318>

443 Wood, S. A., Russell, R., Hanson, D., Williams, R. J., & Dunne, J. A. (2015). Effects of spatial scale of
 444 sampling on food web structure. *Ecology and Evolution*, 5(17), 3769–3782.
 445 <https://doi.org/10.1002/ece3.1640>

446 Xu, M. (2020). *Understanding graph embedding methods and their applications*.
 447 <http://arxiv.org/abs/2012.08019>

448 Yan, S., Xu, D., Zhang, B., & Zhang, H.-J. (2005). Graph embedding: A general framework for
 449 dimensionality reduction. *2005 IEEE Computer Society Conference on Computer Vision and Pattern*
 450 *Recognition (CVPR'05)*, 2, 830–837 vol. 2. <https://doi.org/10.1109/CVPR.2005.170>

- 451 Young, S. J., & Scheinerman, E. R. (2007). Random Dot Product Graph Models for Social Networks. In A.
452 Bonato & F. R. K. Chung (Eds.), *Algorithms and Models for the Web-Graph* (pp. 138–149). Springer.
453 https://doi.org/10.1007/978-3-540-77004-6_11
- 454 Zhou, J., Cui, G., Hu, S., Zhang, Z., Yang, C., Liu, Z., Wang, L., Li, C., & Sun, M. (2020). Graph neural
455 networks: A review of methods and applications. *AI Open*, 1, 57–81.
456 <https://doi.org/10.1016/j.aiopen.2021.01.001>




Figure 1: Overview of the embedding process. A network (A), represented as its adjacency matrix (B), is converted into a lower-dimensional object (C) where nodes, subgraphs, or edges have specific values (see tbl. 1). For the purposes of prediction, this low-dimensional object encodes feature vectors for *e.g.* the nodes. Embedding also allows to visualize the structure in the data differently (D), much like with a principal component analysis.




Figure 2: From a low-dimensional feature vector (see fig. 1), it is possible to develop predictive approaches. Nodes in an ecological network are species, for which we can leverage phylogenetic relatedness (*e.g.* Strydom, Bouskila, et al., 2021) or functional traits to fill the values of additional species we would like to project in this space (here, I, J, K, and L) from the embedding of known species (here, A, B, C, and D). Because embeddings can be projected back to a graph, this allows us to reconstruct a network with these new species. This approach constitutes an instance of transfer learning.

Table 1: Overview of some common graph embedding approaches, by time of publication, alongside examples of their use in the prediction of species interactions. These methods have not yet been routinely used to predict species interactions; most examples that we identified were either statistical associations, or analogues to joint species distribution models. ^a: statistical interactions; ^b: joint-SDM-like approach.

Method	Embedding approach	Reference	Application in species interactions
tSNE	nodes through statistical divergence	Hinton & Roweis (2002)	Cieslak et al. (2020) ^a
RDPG	graph through SVD	Young & Scheinerman (2007)	Poisot et al. (2021); Dalla Riva & Stouffer (2016)
DeepWalk	graph walk	Perozzi et al. (2014)	Wardeh et al. (2021)
FastEmbed	graph through PCA/SVD analogue	Ramasamy & Madhow (2015)	
LINE	nodes through statistical divergence	Tang et al. (2015)	
SDNE	nodes through auto-encoding	D. Wang et al. (2016)	
node2vec	nodes embedding	Grover & Leskovec (2016)	
graph2vec	sub-graph embedding	Narayanan et al. (2017)	
DMSE	joint nodes embedding	D. Chen et al. (2017)	D. Chen et al. (2017) ^b
HARP	nodes through a meta-strategy	H. Chen et al. (2017)	
GraphKKE	graph embedding	Melnyk et al. (2020)	Melnyk et al. (2020) ^a
Joint methods	multiple graphs	S. Wang et al. (2021)	