OCTOBER 26, 2014

Macroevolution of ecological networks

Timothée Poisot







$$\frac{\mathrm{d}}{\mathrm{dt}}x = \frac{1}{2}\mu\sigma^2 N^\star(x)\frac{\partial}{\partial x'}\omega(x', x)$$


$1-p$


$1-p$
ϵ


$1-p$
ϵ
$1-\epsilon$


$1-p$
ϵ
$1-\epsilon$
p
ϵ
$1-\epsilon$














1
0.8
0.6
0.4
0.2
0
10^{-3}
10^{-2}
10^{-1}
10^{0}
Probability of losing new interactions


1
0.8
0.6
0.4
0.2
0
10^{-3}
10^{-2}
10^{-1}
10^{0}
Probability of losing new interactions


1
0.8
0.6
0.4
0.2
0
10^{-3}
10^{-2}
10^{-1}
10^{0}
Probability of losing new interactions

$$f(\theta) = \mathbf{s}_{\text{sim.}} \quad \approx \quad \mathbf{s}_{\text{obs.}}$$


Tendency for specialization
10^{1}
10^{0}
10^{-1}
10^{-3}
10^{-2}
10^{-1}
10^{0}
Probability of losing new interactions


Tendency for specialization
10^{1}
10^{0}
10^{-1}
herbivory
pollination
dispersal
parasitism
phage
10^{-3}
10^{-2}
10^{-1}
10^{0}
Probability of losing new interactions


herbivory
pollination
dispersal
parasitism
phage
Tendency for specialization
10^{1}
$10^{0.8}$
$10^{0.6}$
$10^{0.4}$
$10^{0.2}$
10^{0}
10^{-2}
10^{-1}
10^{0}
Probability of losing new interactions


network structure is predicted by
simple evolutionary rules
types of networks differ on their
tendency to retain interactions