

Quantum Physics - Notes - v0.2.0

260236

October 2026

Preface

Every theory section in these notes has been taken from the sources:

- Quantum Mechanics. [1]
- Fondamenti di fisica atomica e quantistica. [2]
- Modern Quantum Mechanics. [3]

About:

 [GitHub repository](#)

These notes are an unofficial resource and shouldn't replace the course material or any other book on quantum physics. It is not made for commercial purposes. I've made the following notes to help me improve my knowledge and maybe it can be helpful for everyone.

As I have highlighted, a student should choose the teacher's material or a book on the topic. These notes can only be a helpful material.

Contents

1	From Wave-Particle Duality to Quantum States	4
1.1	Experimental Motivation: Wave-Particle Duality	4
1.2	Intrinsic Randomness of Quantum Measurements	22
	Index	31

1 From Wave-Particle Duality to Quantum States

1.1 Experimental Motivation: Wave-Particle Duality

The purpose of this section is not to give a history of the development of quantum mechanics, but rather to discover **what kind of mathematical theory we will eventually need** to describe the phenomena that we will discuss in these notes.

We begin from a very basic question: “*what is light?*”. Classically, there are two very different possibilities:

1. A **Particle** is something localized. At a given instant it **occupies a particular position, follows a trajectory, and arrives somewhere** as an individual object.
2. A **Wave**, on the other hand, is **spatially extended**. Different portion of the wave can overlap and interfere.

Historically, both descriptions were proposed for light. But for our purposes, the important question is not who proposed what. The important question is: **how can an experiment distinguish wave behaviour from particle behaviour?** In other words, definitions alone cannot choose between the two descriptions. So we **need an experimental criterion that allows us to distinguish between the two behaviors.**

What phenomenon is characteristic of waves? The wave vs. particle descriptions of light

Suppose we have light traveling from a source toward some obstacle and then onto a screen. **If light behaved only like classical particles**, the simplest picture would be:

source $\rightarrow$ particle trajectories $\rightarrow$ screen

Each particle follows some path, reaches a particular location, and contributes there.

But **waves behave differently** because different contributions can be added together. For two waves:

$$\psi_1, \quad \psi_2$$

The **combined disturbance**¹ is:

$$\psi_{\text{tot}} = \psi_1 + \psi_2$$

This simple operation has a crucial consequence: the **two waves may reinforce each other or cancel each other.** That phenomenon is called **Interference**.

¹The **total/resultant wave** produced when the two waves are present at the same place and time.

1 From Wave-Particle Duality to Quantum States

1.1 Experimental Motivation: Wave-Particle Duality

So already we have the first important distinction between waves and particles:

- **Particles** are localized objects.
- **Waves** are superposable² and extended disturbances.

At this point, “superposition” is still the familiar classical-wave idea.³ We are not yet stating the quantum superposition principle. Also, we have seen that the **interference phenomenon** is a characteristic of waves, but we have not yet seen any experimental evidence for it.

≡ Diffraction and interference as a signatures of waves

Interference is often observed experimentally through a **Diffraction** pattern. It is one of the **most convenient experimental manifestations of interference**. In general, interference is the general wave phenomenon, and diffraction is a **particular wave phenomenon in which a wave spreads after an aperture/obstacle, and the resulting diffraction pattern is produced by interference**. So we can have interference without diffraction, for example two independent water waves meeting each other.

Deepening: What does a “wave” actually have?

Before talking about light, think about something more intuitive: a **wave on a rope**. Imagine a rope that is fixed or held at the ends. If we move one point up and down, the disturbance propagates along the rope. The rope itself does **not** travel along with the wave. Each small piece of rope mainly moves up and down around its equilibrium position, while the **wave pattern** propagates horizontally.

Mathematically, at each point x along the rope, it has some displacement from equilibrium:

$$y(x, t)$$

Which is the description of the wave, where y is the vertical displacement of the rope at position x and time t ($y = 0$ is the equilibrium position). At one instant maybe $y = 2$ cm, and somewhere else $y = -1$ cm.

Thus, the quantity y is the **wave amplitude** at that point and instant. It indicates how far the rope is displaced from its equilibrium position.

For the case of **light**, instead of writing $y(x, t)$, we can think about:

$$E(x, t)$$

So the “amplitude” is the **value of the oscillating wave quantity**.

²They can be added together. More precisely, in a linear wave system, if ψ_1 and ψ_2 are allowed waves, then $\psi_1 + \psi_2$ is also an allowed wave.

³The sum of two waves is still a wave.

1 From Wave-Particle Duality to Quantum States

1.1 Experimental Motivation: Wave-Particle Duality

Deepening: What happens when two waves meet?

This is the defining ingredient behind interference. Suppose two waves arrive at the same place. Their amplitudes add:

$$E = E_1 + E_2$$

This is the superposition principle for classical waves. Now imagine at instant:

$$E_1 = +1, \quad E_2 = +1$$

Then:

$$E = +2$$

They reinforce each other. But perhaps instead:

$$E_1 = +1, \quad E_2 = -1$$

Then:

$$E = 0$$

They cancel each other. That is interference. So the fundamental thing that interference is **the amplitude**: $E_1 + E_2$.

Suppose light hits a screen. We don't see the electric field E directly:

$$E = +3, \quad E = -2, \quad E = +1, \quad \dots$$

Because the electromagnetic field oscillates extremely rapidly, and our eyes cannot follow it. Instead, the screen receives **energy** from the light.

- A region receiving more energy appears brighter.
- A region receiving less energy appears darker.

The quantity describing this is called the **Intensity** I , the **energy delivered per unit area per unit time**, roughly speaking.

For a light wave, intensity is proportional to $\propto$ the square of the wave amplitude:

$$I \propto |E|^2$$

 **Why square the amplitude?** One immediate reason the square makes sense is that amplitude can have a sign (e.g., $E = +1$ or $E = -1$), but **energy cannot be negative**. Squaring guarantees that the intensity is always positive. The sign is important for **interference**, but the final measured energy/intensity is always positive. Thus:

1. Add the amplitudes of the waves to get the total amplitude.
2. Square the total amplitude to get the intensity.

 Example 1: Interference of two waves

Imagine each wave alone would have amplitude magnitude 1. If only wave 1 existed:

$$E_1 = 1$$

So its intensity would be proportional to:

$$I_1 \propto 1^2 = 1$$

Similarly, if only wave 2 existed:

$$E_2 = 1$$

So its intensity would be proportional to:

$$I_2 \propto 1^2 = 1$$

We might naïvely think that two waves together should therefore give:

$$I = I_1 + I_2 = 1 + 1 = 2$$

But that is **not necessarily true**. We must first add their amplitudes.

- **Case A: the waves reinforce each other.** If:

$$E_1 = 1, \quad E_2 = 1$$

Then:

$$E = E_1 + E_2 = 1 + 1 = 2$$

So the intensity is:

$$I \propto |E|^2 = 2^2 = 4$$

We have **constructive interference**.

And if we had:

$$E_1 = -1, \quad E_2 = -1$$

Then:

$$E = E_1 + E_2 = -1 - 1 = -2$$

So the intensity is:

$$I \propto |E|^2 = |-2|^2 = 2^2 = 4$$

Again, we have **constructive interference**, even though the amplitudes are negative. The sign of the amplitude does not matter for the intensity.

- **Case B: the waves cancel each other.** If:

$$E_1 = 1, \quad E_2 = -1$$

Then:

$$E = E_1 + E_2 = 1 - 1 = 0$$

So the intensity is:

$$I \propto |E|^2 = 0^2 = 0$$

So the screen is dark there. This is **destructive interference**.

1 From Wave-Particle Duality to Quantum States

1.1 Experimental Motivation: Wave-Particle Duality

Let's generalize the previous example. Imagine one sinusoidal wave reaching some point P :

$$E_1(t) = E_0 \cos(\omega t)$$

Its average intensity is proportional to the square of the amplitude:

$$I_1 \propto E_0^2$$

So far, nothing strange. Now suppose **two waves** reach the same point P . Because ordinary wave equations are linear, the physical field is the sum:

$$E(t) = E_1(t) + E_2(t)$$

This is the classical **superposition principle for waves**. But intensity is not proportional to E , but rather to the **square** of the resulting amplitude:

$$I \propto |E|^2 \implies I \propto |E_1 + E_2|^2$$

Expanding:

$$|E_1 + E_2|^2 = |E_1|^2 + |E_2|^2 + 2\Re(E_1^* E_2)$$

The last term $2\Re(E_1^* E_2)$ is called the **Inference Term**. This term is the reason why:

$$|\psi_1 + \psi_2|^2 \neq |\psi_1|^2 + |\psi_2|^2$$

Or in words, the **intensity of two waves together is not equal to the sum of their individual intensities**. The inference term can be:

- **Constructive Interference** if the phases agree:

$$2\Re(E_1^* E_2) > 0$$

The two contributions reinforce each other. We have a **maximum** in the intensity.

- **Destructive Interference** if the phases disagree:

$$2\Re(E_1^* E_2) < 0$$

The contributions oppose each other and may cancel. We have a **minimum** in the intensity.

So an **alternating pattern of maxima and minima is a signature of interference**, and it is a **signature of wave-like behavior**. Therefore, observing such a diffraction/interference pattern provides an experimental criterion for identifying the wave-like nature of light.

Deepening: The interference term

Let's remove for a moment the absolute value from each term. Then the intensity is still proportional to the square of the amplitude:

$$I \propto (E_1 + E_2)^2$$

1 From Wave-Particle Duality to Quantum States

1.1 Experimental Motivation: Wave-Particle Duality

Expanding it exactly like ordinary algebra:

$$(E_1 + E_2)^2 = E_1^2 + E_2^2 + 2E_1E_2$$

The last term $2E_1E_2$ is the **Interference Term** in the simplest real-valued case. It is really **important** because without it, we would just have:

$$E_1^2 + E_2^2$$

Which is just “intensity from wave 1 plus intensity from wave 2”. The extra term tells us what happens when the two waves meet.

For example, if $E_1 = 1$ and $E_2 = 1$, then the intensity is:

$$I \propto (1 + 1)^2 = 4$$

But separately, the intensity from each wave is:

$$I_1 + I_2 = 1^2 + 1^2 = 2$$

The extra term $+2$ comes from the interference term:

$$2E_1E_2 = 2 \cdot 1 \cdot 1 = 2$$

So the waves reinforce each other. Now take $E_1 = 1$ and $E_2 = -1$. Then the intensity is:

$$I \propto (1 - 1)^2 = 0$$

But separately, the intensity from each wave is:

$$I_1 + I_2 = 1^2 + (-1)^2 = 1 + 1 = 2$$

The extra term -2 comes from the interference term:

$$2E_1E_2 = 2 \cdot 1 \cdot (-1) = -2$$

So the interference term is simply what can either increase the intensity or decrease it.

🔍 Now, why did we previously write the interference term as $2\Re(E_1^*E_2)$ instead of $2E_1E_2$? Because for light waves we often represent the amplitude using complex numbers. Then the correct version of the same algebra becomes:

$$\begin{aligned} |E|^2 &= |E_1 + E_2|^2 \\ \downarrow |z|^2 &= z \cdot z^* = (x + iy)(x - iy) = x^2 + y^2 \\ \downarrow \text{if we assume } E_1 + E_2 &= z \\ &= (E_1 + E_2)(E_1 + E_2)^* \\ &= (E_1 + E_2)(E_1^* + E_2^*) \\ &= E_1E_1^* + E_1E_2^* + E_2E_1^* + E_2E_2^* \\ \downarrow E_1E_1^* &= |E_1|^2, \quad E_2E_2^* = |E_2|^2 \\ &= |E_1|^2 + |E_2|^2 + E_1E_2^* + E_2E_1^* \end{aligned}$$

1 From Wave-Particle Duality to Quantum States

1.1 Experimental Motivation: Wave-Particle Duality

Now the last two terms are complex conjugates of each other. Let:

$$z = E_1 E_2^*$$

Thus its complex conjugate is:

$$z^* = (E_1 E_2^*)^* = E_1^* (E_2^*)^* = E_1^* E_2$$

Therefore:

$$E_1 E_2^* + E_2 E_1^* = z + z^*$$

And for any complex number:

$$z = a + ib$$

We have:

$$z^* = a - ib$$

So:

$$z + z^* = a + ib + a - ib = 2a$$

But a is the real part of z , represented as $\Re(z)$. So we have:

$$z + z^* = 2\Re(z)$$

Therefore:

$$E_1 E_2^* + E_2 E_1^* = 2\Re(E_1 E_2^*) \equiv 2\Re(E_1^* E_2)$$

And conceptually it is just the same cross term as $2E_1 E_2$.




Figure 1: **Diffraction and interference in a two-slit experiment.** An incident plane wave passes through two slits; the emerging waves spread, overlap, and interfere, producing an intensity pattern on the screen modulated by a diffraction envelope.

 Deepening

Some details about the Figure 1:

- ❓ **What is the “incident plane wave”?** On the left we have several **parallel blue vertical lines**, those lines are **wavefronts of the same incoming wave**.

A wavefront is a set of points where the wave is in the same state of oscillation. We do not need to know particular details; visually, think of each blue line as one crest/front of the wave. Because these wavefronts are straight and parallel, we call it a **plane wave**. So “*incident plane wave*” simply means **a wave arriving at the barrier with flat, parallel wavefronts**. “*Incident*” just means **incoming**.

Finally, the arrow indicates that the wave propagates from left to right.

- ❓ **What are the “two slits”?** The grey vertical object is a barrier with **two small openings**. Most of the wave is blocked by the barrier, only the portions reaching those openings pass through.

Here, the crucial observation is what happens **after** each slit. Instead of continuing only straight ahead, the wave **spreads out**. That’s why the figure draws curved blue wavefronts coming from each slit.

This spreading is the **diffraction** part! Each slit effectively gives us a new spreading wave. And now we have precisely what we need for **interference**: two waves reaching the same points on the screen (black rectangle on the right).

- ❓ **What happens between the slits and the screen?** Pick some point P on the screen. There are two wave contributions reaching P : ψ_1 from the upper slit and ψ_2 from the lower slit. The red straight lines in the figure illustrate examples of those two paths.

The two contributions add:

$$\psi(P) = \psi_1(P) + \psi_2(P)$$

But depending on the point P , the two paths have different lengths. So the waves may arrive “*in phase*” (reinforce, constructive interference), “*out of phase*” (cancel out, destructive interference), or somewhere in between (weaker reinforcement or weaker cancellation). And that’s why different points of the screen have different brightness.

- ❓ **Why is the center of the screen bright?** Call the two slit-to-screen distances r_1 and r_2 . At the exact center of the screen, by symmetry, the two distances are equal:

$$r_1 = r_2$$

Therefore the two waves have traveled the same distance, and they reinforce each other:

$$\psi_1 + \psi_2 \implies I \propto |\psi_1 + \psi_2|^2$$

So the intensity is large and the central bright band corresponds to constructive interference.

- ❓ **What are the dark bands?** Move upward or downward along the screen. Now one slit is slightly closer than the other:

$$r_1 \neq r_2$$

At certain position, the difference in traveled distance causes one wave to arrive exactly opposite in phase to the other.

For example:

$$\psi_1 = A, \quad \psi_2 = -A$$

Therefore:

$$\psi_1 + \psi_2 = A - A = 0$$

And hence the intensity is zero:

$$I \propto |\psi_1 + \psi_2|^2 = 0$$

So that location is dark, corresponding to destructive interference.

- ❓ **What is that red curve on the right?** The horizontal direction of the red graph represents the **intensity** I measured on the screen, while vertically we move along the screen coordinate y .

So where the solid red curve extends far to the right, the intensity is large and the screen is bright. Where it comes close to zero $I \approx 0$, the screen is dark.

So the oscillating solid red curve is essentially a graph of $I(y)$, the intensity as a function of vertical position on the screen.

- ❓ **What is the “envelope”?** Look at the **dashed red curve** labelled “envelope”. It is **not** “the level of interference”. Instead, it describes how large the interference peaks are allowed to become at different positions.

A more natural question is: *why are the bright fringes near the center stronger than the bright fringes farther away?* What is exactly what the dashed curve in figure is trying to show. The red solid curve $I(y)$ oscillates between bright and dark because of the **interference between the two slits**.

But notice also that the heights of the bright peaks are not all the same. Near the center, the peaks are large; farther away, they become smaller. The dashed curve simply joins the tops of those possible peaks. So the **envelope** is the smooth curve giving the

maximum possible intensity at each position on the screen (at each y). In other words, it is just a boundary for the interference pattern.

❓ **But why does the envelope exist?** Because the slits are not mathematical points; they have a finite width. Even **one slit by itself** produces a diffraction (i.e., spreading) pattern on the screen. So each slit does not send the same wave strength equally in every direction. It sends most strongly approximately forward, around the center of the screen, and more weakly at larger angles.

Therefore, before we even combine the two slits, each slit already has something like: strong near the center, weaker farther away. That broad distribution is the **single-slit diffraction pattern**. Now, if we combine the two slits, the two effects happen simultaneously and each slit diffracts; then the waves from the two slits interfere and the final pattern is the product of the two effects:

- **Single-slit diffraction** gives the broad envelope.
- **Two-slit interference** gives the oscillating pattern within that envelope.




Figure 2: **The interference term.** The resulting intensity depends on the relative phase $\phi_1 - \phi_2$: constructive interference produces maxima, while destructive interference produces minima.

1 From Wave-Particle Duality to Quantum States

1.1 Experimental Motivation: Wave-Particle Duality

⚙️ Electron diffraction

Now we take the criterion we just established for waves (i.e., the presence of interference)

$$\text{diffraction/interference} \implies \text{wave-like behaviour}$$

And apply it to something that, classically, we would expect to behave as a particle: **the electron.**

The surprising experimental observation is that a beam of electrons, when sent onto a crystalline sample, produces a **pattern that looks essentially like the diffraction pattern produced by light.**

💡 **The idea.** Before this experiment, the classical expectation was: electron behaves $\approx$ particle. *Why?* Because electrons have properties we normally associate with particles, they have mass, charge, momentum, and can form an electric current.

But now we perform an experiment designed to test for wave behaviour. We send electrons toward an ordered crystalline structure. And experimentally we observe, again, a **diffraction pattern.** And from the previous discussion, we know that it means:

$$\text{diffraction pattern} \implies \text{interference} \implies \text{wave-like behaviour}$$

So the immediate conclusion is that **electrons behave like waves.**

❓ **Why is this surprising?** Because the same physical object (the electron) was already well described as a particle in other experiments. So we have:

$$\text{electron} \quad \left\{ \begin{array}{l} \text{particle-like in some experiments} \\ \text{wave-like in diffraction experiments} \end{array} \right.$$

This is already the beginning of **wave-particle duality**. However, we shouldn't jump ahead just yet. We'll see it in more detail later. At this point, all we have established is that **an electron beam produces a wave-like diffraction pattern.**

❓ What exactly do we observe?

1. The electron source produces a collimated beam of electrons.
2. These electrons hit a crystal.
3. A crystal contains atoms arranged in a regular periodic structure.
4. The electrons interact with that periodic structure, and the detected electrons are not distributed uniformly everywhere. Instead, they **appear preferentially in specific directions**, giving a pattern of bright spots.

That is exactly the kind of spatial organization we expect from diffraction. The regular atomic arrangement plays, for electrons, a role analogous to a diffraction grating for light.

1 From Wave-Particle Duality to Quantum States

1.1 Experimental Motivation: Wave-Particle Duality




Figure 3: **Electron diffraction from a crystal.** A collimated electron beam produced by an electron gun is incident on a crystalline sample. The scattered electrons are detected on a fluorescent screen, where they form an ordered diffraction pattern with bright maxima separated by low-intensity regions.

1 From Wave-Particle Duality to Quantum States

1.1 Experimental Motivation: Wave-Particle Duality

In Figure 3, we show a schematic of the experimental set-up and the resulting diffraction pattern.

- In panel (a), the **electron gun** is the **source**. It emits a collimated beam of electrons **toward the crystal**. The crystal is not just a random obstacle: its atoms are arranged periodically, so it behaves like a diffraction structure for the electron beam.

The **arrows** leaving the crystal represent the **diffracted directions**. They are not meant to say that we know a classical trajectory for each electron; they indicate the **directions where the scattered electron signal is enhanced because of diffraction**.

Those **electrons then hit the fluorescent screen**. The screen converts the electron impacts into visible light: the electrons excite the fluorescent material, which then emits light.

- Panel (b) shows what we actually observe on that screen: a **set of bright spots at specific positions**. Those bright spots are the diffraction maxima. Between them, the signal is much weaker or absent. That ordered pattern is the important experimental result.

• Single-particle regime

From the previous discussion, we observed that a beam of electrons produces a diffraction pattern after interacting with the crystal, that suggests wave-like behaviour. But there is still **an alternative explanation** we have to **exclude**: **maybe the diffraction pattern is caused by interactions among many electrons**.

Electrons are charged particles, so when many of them are present simultaneously they **can interact with one another**. So we need to find a way to eliminate that possibility, and the solution is to **send only one electron at a time**.

❓ **How do we do that?** We **reduce the intensity** of the electron current emitted by the **electron gun** until successive electrons are separated by a sufficiently large time interval. So instead of having something like:

$$e^- e^- e^- e^- e^-$$

Inside the apparatus (i.e., in the region where the electrons are interacting with the crystal), we arrange approximately:

$$e^- \qquad e^- \qquad e^-$$

With a **large delay between one electron and the next**. So we reduce the current until one electron is sent at a time, with enough delay that individual arrivals can be resolved experimentally. This is what we call the **single-particle regime**.

In the next page (Figure 4), we show a schematic of the two regimes: single-particle and many-electron.

1 From Wave-Particle Duality to Quantum States

1.1 Experimental Motivation: Wave-Particle Duality




Figure 4: **Single-particle regime.** (a) At high current, many electrons are present simultaneously and can interact with each other. The screen shows a continuous diffraction pattern. (b) At low current, only one electron is present at a time, so electron-electron interaction is excluded. Each electron produces a localized flash on the screen

Why is this experiment important? Because now **two different electrons are not present in the apparatus at the same time.** Therefore, they cannot interact with each other during propagation. So if diffraction behaviour is still observed, we can no longer explain it by saying: “*electron 1 interacts with electron 2*” and this collective interaction generates the pattern. That **possibility has been removed.** Now the logic becomes:

many electrons simultaneously $\Rightarrow$ electron-electron interaction is possible,
 one electron at a time $\Rightarrow$ electron-electron interaction is excluded

So the experiment becomes much interesting.

1 From Wave-Particle Duality to Quantum States

1.1 Experimental Motivation: Wave-Particle Duality


☉ **What is actually detected?** Each electron reaches the fluorescent screen individually. When an electron hits the screen, it produces a localized luminous event: the electron excites the fluorescent material and a small flash of light is observed. So experimentally we do not see a continuous faint wave spread over the whole screen. We see:

one electron $\rightarrow$ one localized detection event

Then some time later:

next electron $\rightarrow$ another localized detection event

That is already very interesting, because the propagation had shown wave-like diffraction, while the detection is localized. In other words, we have demonstrated that **the electron is detected as a localized event** and, because we send electrons one at a time, we have removed the possibility that the pattern comes from electron-electron interactions.



one electron $\rightarrow$ one localized detection event

Figure 5: **Detection of individual electrons.** Each electron produces a localized flash on the fluorescent screen. The flashes are separated in time, and after many repetitions they accumulate to form the same diffraction pattern as in Figure 3.

Now, let's **repeat the experiment many times** to see what happens and if the position of the next spot can be predicted in advance.

1 From Wave-Particle Duality to Quantum States

1.1 Experimental Motivation: Wave-Particle Duality

Localized detection events and the emergence of a diffraction pattern

In the previous figures, electrons are sent **one at a time**. Each electron hits the fluorescent screen and produces one localized flash. So for a single electron, we do not see a little wave spread across the screen, we see one point.

The location of the next flash cannot be predicted from the previous ones. Because successive electrons arrive at different spots, then the next position cannot be determined beforehand. But the main point is that **random does not mean uniformly random**. The flashes do not populate every point of the screen equally. Instead, some regions receive many detections, while other regions receive very few or none.

When many single-electron detections are **accumulated**, those frequencies reproduce the same diffraction structure seen with the ordinary electron beam. In Figure 6, we show a schematic of the accumulation process. The first panel shows only a few detections, which appear scattered and do not reveal any structure. As more and more electrons are sent, the **flashes accumulate**.




Figure 6: **Accumulation of many single-electron detection events.** Each electron produces a localized flash on the fluorescent screen. After many repetitions, the flashes accumulate to form the same diffraction pattern as in Figure 3.

So after a small number of electrons, we might see something apparently messy ($N = 50$). After more electrons ($N = 500$), and finally after many electrons ($N = 3000$), the **diffraction pattern emerges statically**.

🔍 **Why this is surprising.** Each electron is detected as a particle:

one electron $\rightarrow$ one localized spot

But the ensemble of many electrons behaves as if some wave-like interference rule were controlling where detections are more or less likely:

many individual electrons $\rightarrow$ diffraction pattern

So we have **two apparently contradictory features in the same experiment**:

1 From Wave-Particle Duality to Quantum States

1.1 Experimental Motivation: Wave-Particle Duality

1. At **detection** (one electron at a time), the electron behaves like a **particle**. Because the electron appears at one precise location, not spread over the whole screen.
2. In the **accumulated distribution** (many electrons), the electron behaves like a **wave**. Because the spots organize according to a diffraction pattern.

So if we repeat the experiment a huge number of times, we can say something statistical: “*some positions will be hit much more often than others*”. So quantum mechanics is already forcing us toward two levels of description:

🔍 Individual event: **unpredictable** and **localized**.

📊 Many events: **highly structured statistics**.

📌 Key Takeaways 1

We began by asking how to identify wave behaviour:

interference/diffraction $\Rightarrow$ wave-like behaviour

Then we observed electron diffraction:

electrons $\rightarrow$ diffraction

Then we sent electrons one by one, eliminating electron-electron interactions:

one electron at a time

Each electron was detected locally:

one electron $\rightarrow$ one spot

But many such detections reconstructed the diffraction pattern:

many individual detections $\rightarrow$ wave-like diffraction pattern


Therefore, **quantum objects exhibit wave-like propagation and particle-like detection.** Or even more compactly:

interference during propagation + localization at detection

This is the experimental problem that quantum mechanics now has to explain. And the natural next question is no longer “*wave or particle?*” but: ***why can we predict the distribution but not the individual detection?***

1 From Wave-Particle Duality to Quantum States

1.1 Experimental Motivation: Wave-Particle Duality



1.2 Intrinsic Randomness of Quantum Measurements

🔍 Individual detection events are unpredictable

We have already established that electrons exhibit **wave-like diffraction** but are detected as **individual, localized particles**. Now we want to investigate a different question: *can we predict where the next electron will be detected?* This question introduces one of the most important features of quantum mechanics: **randomness in measurement outcomes**.

Let's consider the single-electron experiment again (page 14):

electron gun $\rightarrow$ crystal $\rightarrow$ fluorescent screen

We send electrons one at a time, keeping the experimental conditions unchanged. In particular, the electrons are prepared with the same nominal energy and momentum, and the crystal and screen remain in the same positions.

As we have seen, if we repeat the experiment only a few times (e.g., 3 times), we might observe:

$$e_1 \rightarrow x_1, \quad e_2 \rightarrow x_2, \quad e_3 \rightarrow x_3$$

Where e_i is the i -th electron and x_i is the position on the screen where it is detected. Even though we repeat the same preparation procedure, the detection positions x_1 , x_2 , and x_3 are **different** and **unpredictable**. So the important observation was that **the next detection position cannot be determined in advance from the information supplied by the experimental preparation or from the previous detections.**

🔍 **Why was this observation surprising?** Because a classical particle, like a bullet, would always hit the same spot on the screen if it is prepared in the same way.

In classical mechanics, if we know a particle's initial position and momentum, together with the forces acting on it, we can in principle calculate its trajectory:

$$(x(0), p(0)) + \text{equations of motion} \longrightarrow x(t)$$

Therefore, we can predict where the particle will arrive. If our classical prediction is inaccurate, we might suspect that we did not know the initial conditions precisely enough.

Instead, **quantum mechanics changes this situation.** Even when we prepare an electron in the same quantum state and repeat the same measurement, the standard quantum description generally does not provide a definite position for the next detection. Instead, it provides probabilities for different possible outcomes.

At this stage, we do not need to calculate those probabilities. What matters is the experimental observation:

$$\text{same preparation procedure} \not\Rightarrow \text{same detection position}$$

Notice that this does **not** mean quantum mechanics cannot predict anything. It means that **predicting the individual result** and **predicting the statistics of many results** are **different tasks**.

❓ **Does unpredictable mean equally likely everywhere?** No. Suppose the fluorescent screen has three possible detection regions: A , B , and C . The electron might arrive in any of them, but it does not follow that

$$P(A) = P(B) = P(C)$$

Some regions may be much more likely than others, and some may have a negligible probability of detection. That is perfectly compatible with the diffraction pattern we observed in the previous section.

unpredictable individual outcome $\neq$ uniformly random outcome

❓ **Have we already demonstrated intrinsic randomness?**⁴ We should distinguish the observation from its interpretation:

- What we have observed is that individual detection positions are unpredictable under our controlled experimental conditions.
- That **observation alone does not prove that randomness is fundamentally intrinsic rather than a consequence of unknown variables.**

One might still ask whether some inaccessible information determines each outcome, just as an apparently random coin toss can be determined by its initial conditions. We will not answer it yet, but we will investigate it in the next pages.

Now, we have identified the randomness of individual detections. The next question is: *if individual outcomes are unpredictable, how can the diffraction pattern be so reproducible?*

✓ Key Takeaways 2

One electron is sent toward the crystal.

↓

One localized detection occurs.

↓

Its position cannot generally be predicted in advance.

Quantum measurements can produce different individual outcomes even when the system is prepared in the same way.

⁴**Intrinsic** means inherent to the nature of something, rather than caused by external factors. In quantum physics, *intrinsic randomness* refers to randomness that is fundamental to the measurement outcome, not merely due to incomplete knowledge or imperfect measuring instruments.

📊 Statistical regularity from many repetitions

Thus **we cannot predict where the next electron will be detected**, but this raises an apparent contradiction: if individual electron detections are unpredictable, **how can the diffraction pattern be so regular and reproducible?** In other words, how can we reconcile the unpredictability of individual events with the predictability of the overall pattern?


⚡ **Random individual events do not imply a random overall distribution.** Let's return to our electron-diffraction experiment. We prepare the electron gun, crystal and fluorescent screen, and send electrons **one at a time**. For each electron, we observe one localized flash at a position that we cannot predict beforehand.

Now suppose we repeat the experiment N times, without changing the experimental conditions. We observe:

$$x_1, x_2, x_3, \dots, x_N$$

Where x_i is the detection position of electron i . At first, with only a few electrons, the spots appear scattered without an obvious pattern. However, as the number of detections increases, something remarkable happens: **the detections accumulate preferentially in certain regions, reproducing the characteristic diffraction pattern.**

This is the same experimental picture developed earlier on page 19, where individual localized flashes appear at unpredictable positions, but they occur with a spatial distribution associated with diffraction.



So we must distinguish between two different types of predictions:

- **Individual prediction:** *where will the next electron arrive?*
- **Statistical prediction:** *how frequently will electrons arrive in different regions after many repetitions?*

Quantum mechanics answers these questions differently.

1 From Wave-Particle Duality to Quantum States

1.2 Intrinsic Randomness of Quantum Measurements

From individual positions to relative frequencies. Imagine dividing the fluorescent screen into regions A , B , and C . After sending N electrons, we count how many were detected in each region: N_A , N_B , and N_C . If these are all the possible detection regions, then:

$$N_A + N_B + N_C = N$$

We can define the **relative frequency** of detections in region A :

$$f_A(N) = \frac{N_A}{N}$$

For example, imagine that after 1,000 detected electrons we have:




Figure 7: **Relative frequencies of electron detections.** A fluorescent screen is divided into three equal-sized regions A , B , and C . After $N = 1000$ detections, the hypothetical counts are $N_A = 700$, $N_B = 250$, and $N_C = 50$, corresponding to relative frequencies of 70%, 25%, and 5%. Each dot represents one localized electron detection. All values are illustrative.

The **next electron remains unpredictable**, it might arrive in A , B , or C . But if we repeat the same experiment a sufficiently large number of times, the relative frequencies tend to stabilize near characteristic values. For region A :

$$f_A(N) \longrightarrow P(A) \quad \text{as } N \rightarrow \infty$$

Under repeated, identically prepared independent trials, $P(A)$ is the probability of obtaining a detection in region A . This does **not** mean that every group of 100 electrons must contain exactly 70 detections in A . The counts still fluctuate (see Figure 8, page 26).

It means that, as the number of repetitions becomes large, the relative fluctuations become less significant and the estimated probabilities become increasingly reliable. This is what we mean by **Statistical Regularity**.



Figure 8: **Statistical convergence of relative frequencies.** The cumulative relative frequency $f_A(N) = N_A/N$ fluctuates as electrons are detected, but tends to stabilize near the hypothetical probability $P(A) = 0.70$ as N increases. The logarithmic horizontal axis emphasizes the fluctuations at small sample sizes. The blue curve connects selected cumulative frequencies, while the dashed red line represents the underlying probability. All values are illustrative.

🕒 **How does this explain the diffraction pattern?** A diffraction pattern contains bright and dark regions. On page 10 we saw that, for classical light, brightness is connected to intensity. So now we can give an **experimental interpretation to the electron-diffraction pattern**:

- At a position where the electron signal is strong, we observe many detections over time.

bright diffraction region $\implies$ **high** detection frequency

- At a position where the signal is weak, we observe fewer detections.

dark diffraction region $\implies$ **low** detection frequency

For an ideal detector, the number of electrons detected per unit area and time gives the local electron arrival flux. Thus, the **diffraction pattern tells us where electron detections are more or less likely**. The pattern is not visible because each electron deposits a fraction of itself in every bright region. It emerges because many complete, individually detected electrons accumulate with different frequencies in different regions.

One electron $\rightarrow$ one localized detection

Many repetitions $\rightarrow$ reproducible spatial distribution

1 From Wave-Particle Duality to Quantum States

1.2 Intrinsic Randomness of Quantum Measurements

 **What does quantum mechanics predict?** In classical mechanics, we often try to predict the trajectory of each particle, given sufficiently precise initial conditions.

In quantum mechanics, the standard theory generally predicts **probabilities for measurement outcomes**. For example, instead of determining the exact position x at which the next electron will be detected, the theory can predict a probability distribution over possible positions.

Those probabilities can then be tested experimentally by repeating the preparation and measurement many times. Later, we will learn the **mathematical rule that connects quantum states with measurement probabilities**: the **Born rule**. For now, we do not need that rule. The experimental observation already tells us that our model must explain both unpredictable individual detections and reproducible statistical distributions.

 **Randomness and regularity are compatible.** A random process can follow a well-defined probability distribution. So, **randomness describes the uncertainty of individual outcomes**, while **statistical regularity describes the reproducibility of their distribution**.

Individual measurement	Many repeated measurements
One localized flash	A collection of localized flashes
Exact position unpredictable	Relative frequencies become stable
Different outcomes are possible	The overall diffraction pattern is reproducible
Random individual result	Predictable statistical distribution

Key Takeaways 3

Individual electron detections are unpredictable



Repeat the experiment many times



Relative detection frequencies stabilize



A reproducible diffraction pattern emerges

Quantum Mechanics does not need to predict each individual detection to make precise experimental predictions. It must correctly predict the statistical distribution of possible outcomes.

❓ Why the diffraction pattern is not caused by interactions between particles

We have already encountered this topic on page 16, so this part is mainly about understanding *why the single-particle experiment is such an important experimental test*.


Our initial observation was that **an electron beam produces a diffraction pattern**. We interpreted this as evidence of wave-like behaviour. But there is a possible objection: *what if the diffraction pattern is caused by interactions between electrons rather than by the wave-like nature of electrons?*

This is not an unreasonable hypothesis. Electrons have negative electric charge. Consequently, they exert repulsive electric forces on one another. If many electrons travel through the apparatus⁵ simultaneously, they may influence each other's motion. One could therefore **hypothesize that their collective interactions produce the observed spatial distribution**. Thus, the question becomes: *how can we test this hypothesis experimentally?*

🗑️ **Eliminate electron-electron interactions.** We change a simple experimental parameter: the **electron emission rate**. We reduce the beam current until successive electrons are sufficiently separated in time that only one incident electron is inside the apparatus at any given moment.


Mathematically, if t_f is the **time needed for an electron to travel through the apparatus**, we want the interval between emissions to satisfy approximately: $\Delta t \gg t_f$.

(a) high rate: $\Delta t \lesssim t_f$



3 electrons inside at once: possible interaction

(b) low rate: $\Delta t \gg t_f$



only 1 electron inside:
no $e^- \leftrightarrow e^-$ interaction

⁵The apparatus is the complete experimental setup used to perform electron diffraction: the electron gun, the crystal, and the fluorescent screen. The electron gun emits electrons, the crystal scatters them, and the fluorescent screen detects their arrival (see page 17).

Now electron e_1 has completed its journey before electron e_2 enters the apparatus. Therefore, the two incident electrons **cannot interact** with one another during their passage.

❓ **What happens experimentally?** We already know the answer:

- First, each electron produces one localized flash on the fluorescent screen.
- Then, after repeating the experiment many times, the accumulated detections reproduce the diffraction pattern.

This result is decisive for our hypothesis, because if the diffraction pattern required interactions **between incident electrons**, it should disappear when those interactions are eliminated. But it does **not disappear**. Therefore, **electron-electron interactions are not necessary for diffraction**.

❓ **Does this mean the electron does not interact with anything?** No! We have eliminated interactions **between different incident electrons**, but the incident electron still interacts with the crystal. The **electron is scattered by the periodic structure of the crystal**. Without the scattering structure, this particular diffraction experiment would **not produce the same pattern**.

$\underbrace{\text{electron} \leftrightarrow \text{electron}}$	$\underbrace{\text{electron} \leftrightarrow \text{crystal}}$
excluded by the experiment	still present (and necessary?)

📌 Key Takeaways 4

Hypothesis: diffraction is caused by interactions among incident electrons

⇓

Send one electron at a time

⇓

Electron-electron interactions are eliminated

⇓

The diffraction pattern still emerges

⇓

The hypothesis is rejected

Wave-like diffraction is not a collective effect produced by interactions among the incident electrons. It is a feature that our description of individual electrons must accommodate (i.e., *must be capable of explaining the phenomenon*), or in other words, our mathematical model of a single electron must be able to explain why repeated single-electron experiments produce a diffraction pattern.

In summary, we **need a theory describing even a single electron must contain the ingredients necessary to predict the diffraction pattern**.

References

- [1] A. Berera and L.D. Debbio. *Quantum Mechanics*. Cambridge University Press, 2021.
- [2] F. Ciccacci. *Fondamenti di fisica atomica e quantistica*. Edises, 2020.
- [3] J.J. Sakurai and J. Napolitano. *Modern Quantum Mechanics*. Cambridge University Press, 2020.

Index

C

Constructive Interference 8

D

Destructive Interference 8

Diffraction 5

I

Inference Term 8

Intensity 6

Interference 4

Interference Term 9

P

Particle 4

S

Statistical Regularity 25

W

Wave 4