

Vibrational Geometrodynamics / Stratified Temporal Network – VG – SERIES 5 – Episode 4:

ON THE INVARIANCE OF INDIVIDUAL FREQUENCY IN THE PRIMARY NETWORK

Author: Physicist Pop Rodica & DeepSeek R1 Assistant

Date: 07.10.2026

Abstract

We propose that each conscious configuration C_{self} in the Primary Network is characterized by an **invariant individual frequency** $\omega_i^{(0)}$, which defines its structural identity, while the **coherence field** $\Phi_i(t)$ defines its evolutionary level. We derive $\omega_i^{(0)}$ directly from the toroidal geometry of the node and from Noether symmetry, and the evolution equation of Φ_i from the **complete Hamiltonian** $\mathcal{H}_{g_{\text{ADN}}}$. The separation of invariant identity / variable expression offers a natural explanation for three phenomena: (a) the persistence of individual consciousness after physical death, (b) the non-loss of spiritual acquisitions, (c) the triple classification of states of existence. The formalism is consistent with VG S4 E4, VG7, PMI, MF, and RTS3.

Keywords : invariant frequency, C_{self} , coherence field, nodal torus, Noether symmetry, nodal identity.

1. Introduction

Within VG/RTS, each node of the Primary Network is characterized by a resonance frequency ω_i and a spin vector $\hat{S}_i$. Until now, the temporal dependence $\omega_i(t)$ has been treated as a complete dynamics (VG S4 E4). In this article we propose a structural refinement: the frequency of any node decomposes into an invariant component $\omega_i^{(0)}$ and a small oscillatory correction $\delta\omega_i(t)$.

This decomposition has a direct consequence for the theory of consciousness: if $\omega_i^{(0)}$ is invariant, then the identity of a C_{self} is mathematically protected against dissolution, while its expression depends exclusively on the coherence field Φ_i .

Methodological observation. This article treats the invariance of individual frequency as a geometric consequence of the toroidal structure of the node, derived from Noether symmetry, not as an independent postulate. The derivation is given in Section 2.

Cross-reference. This work continues the formalization begun in Bi-emergence (Series 5, Episode 1).

2. Formal Decomposition

2.1. Nodal frequency from spin dynamics

From VG S4 E4, the effective frequency of node i is:

$$\omega_i(t) = \frac{1}{\hbar} \left| \sum_{j \in \partial i} J_{ij}^{\text{eff}}(t) \hat{S}_{ij}(t) \right|$$

The spin evolves according to the generalized Landau–Lifshitz equation:

$$\frac{d\hat{S}_i}{dt} = \vec{\Omega}_i \times \hat{S}_i$$

Key property. Precession preserves the modulus of the spin: $|\hat{S}_i| = \text{const.}$ Therefore, the frequency $\omega_i(t)$ cannot drift arbitrarily.

2.2. Derivation of the invariant frequency from toroidal geometry and symmetry

Each node of the Network possesses a toroidal structure, with two loops:

- The large loop (longitudinal) — describes the structural instruction of the node.
- The small loop (transversal) — describes the updating dynamics.

Derivation from the Hamiltonian. The nodal frequency:

$$\omega_i = \frac{1}{\hbar} \left| \sum_{j \in \partial i} J_{ij}^{\text{eff}} \cdot \hat{S}_{ij} \right|$$

We decompose around the state of perfect alignment:

$$\omega_i^{(0)} = \frac{1}{\hbar} \sum_{j \in \partial i} J_{ij}^{\text{eff}} (\text{when } \hat{S}_{ij} = 1)$$

$$\delta\omega_i(t) = \frac{1}{\hbar} \sum_{j \in \partial i} J_{ij}^{\text{eff}} \cdot (\hat{S}_{ij}(t) - 1)$$

Invariance from symmetry (Noether). The torus possesses translational symmetry on its two loops. By Noether's theorem:

$$\frac{d}{dt} \omega_i^{(0)} = 0$$

Explicit Noether. The translational symmetry on the large loop, $\boldsymbol{\phi} \rightarrow \boldsymbol{\phi} + \boldsymbol{\phi}_0$, generates the conservation of the frequency $\omega_i^{(0)}$. The invariance is not accidental — it is the direct consequence of this symmetry.

Topological invariant. The phase accumulated over a complete cycle, $\oint \omega_i^{(0)} dt = 2\pi n$, is a topological invariant (winding number $n \in \mathbb{Z}$).

Conclusion: The invariance of $\omega_i^{(0)}$ is not a postulate — it is a consequence of three principles: (i) the proper mode of the torus, (ii) Noether symmetry, (iii) the topological winding number.

Corollary. The oscillation $\delta\omega_i(t)$ arises from the coupling between the large loop and the small loop, not from a change in the large loop.

2.3. Coherence field — derived from the Hamiltonian

We define the coherence field $\Phi_i(t)$ as the local degree of spin alignment:

$$\Phi_i(t) = \frac{1}{N_i} \sum_{j,k \in \mathcal{N}(i)} \hat{S}_j \cdot \hat{S}_k \in [-1,1]$$

where N_i is the number of pairs of neighbors. Φ_i is dimensionless. $\Phi_{\max} = 1$ corresponds to perfect alignment (all scalar products equal 1).

Derivation from $\mathcal{H}_{\text{ADN}}$. Starting from:

$$\mathcal{H}_0 = \int d^3x \left[\frac{1}{2} \Pi^2 + \frac{1}{2} (\nabla \Phi)^2 + V(\Phi) \right]$$

$$\mathcal{H}_{\text{int}} = g \sum_i \sigma_i \Phi(x_i)$$

The Euler–Lagrange equation:

$$\ddot{\Phi} - \nabla^2 \Phi + \frac{\partial V}{\partial \Phi} = -g \sum_i \sigma_i \delta(x - x_i)$$

In the overdamped regime ($\ddot{\Phi} \approx 0$):

$$\frac{d\Phi}{dt} = -\Gamma \left[-\nabla^2 \Phi + \frac{\partial V}{\partial \Phi} + g \sum_i \sigma_i \delta(x - x_i) \right]$$

Discretized on the network:

$$\frac{d\Phi_i}{dt} = \alpha \sum_{j \in \mathcal{N}(i)} (\Phi_j - \Phi_i) + \eta_i(t)$$

with:

$\alpha = \Gamma$ (diffusion coefficient)

$\eta_i(t) \equiv -\Gamma(\frac{\partial V}{\partial \Phi_i} + g\sigma_i)$ (structured source term)

Physical interpretation of the source term. The term $\eta_i(t)$ is not stochastic noise — it is a structured source term representing the coupling with pair fluctuations (virtual particles as transition operators, cf. Bi-emergence). The coherence equation is not merely diffusive: it is modulated by resonance with pair fluctuations.

Conclusion: The coherence evolution equation is derived from the complete Hamiltonian, not postulated phenomenologically.

2.4. Hypotheses

Postulate (P1). The value $\omega_i^{(0)}$ is structurally invariant for any stable C_{self} .

Status: It is no longer an independent postulate – it is a theorem (Section 2.2).

Hypothesis (H2). The evolution of a C_{self} consists in the variation of $\Phi_i(t)$; the invariant frequency $\omega_i^{(0)}$ is untouched in the normal regime.

Parenthesis (nodal reset). In a nodal reset (the Atlantis case), $\omega_i^{(0)}$ can be perturbed, not through natural evolution, but through external intervention (artificially forced $\mathcal{T}_{\alpha\beta}$). This remains an open question (Section 5, Q3).

3. Consequences

3.1. Separation identity / level

Identity = $\omega_i^{(0)}$ (invariant, geometric)

Level = $\Phi_i(t)$ (variable, derived)

3.2. Persistence after physical death

At physical death:

- physical anchoring $\rho \rightarrow 0$
- $\omega_i^{(0)}$ remains (geometric invariant)
- $\Phi_i^{\max}$ remains (the coherence achieved cannot spontaneously decrease)

Physical death = transition $\rho \rightarrow 0$ preserving $\omega_i^{(0)}$ and $\Phi_i^{\max}$

Connection with RTS3 (Membrane). The transition involves decoupling from the slice C_α and preserving the C_{self} in the nodal membrane, with the possibility of re-anchoring in C_β .

3.3. Triple classification of states of existence

Class	ρ	Φ	VG/RTS Regime Anchoring
Luminous evolved	↓↓↓	↑↑↑	Pure MF No material anchoring
Fluidic + material	↓↓	↑↑	MF + partial Partial anchoring
Fluidic	↓	↑	Weak anchoring in the Network Strong material anchoring

The hierarchy is ρ/Φ , not moral.

3.4. Relation with the torus

- Large loop = $\omega_i^{(0)}$ (proper mode, invariant)
- Small loop = $\Phi_i(t)$ (amplitude, variable)

3.5. Interface operators

The intermediate entities are interface operators $\mathcal{I}$ between strata:

- not pure MF (class 1)
- not dense matter (class 3)
- intermediate regime (class 2)

Their role: modulation of $\mathcal{T}_{\alpha\beta}$ between slices.

4. Relation with existing VG/RTS results

Concept	Relation
Nodal Echo	Stores $\omega_i^{(0)}$, not Φ
PMI	$m_0 c^2 = \sum \hbar \omega_i$ (see decomposition below)
MF	$\rho \downarrow, \omega_i^{(0)}$ unchanged, $\Phi \uparrow$
Bi-emergence	$\omega_i^{(0)} = \text{instruction}; \delta \omega_i = \text{vibration}$
VG7	T_0 invariant $\leftrightarrow \omega_i^{(0)}$ invariant
Atlantis	Priests: $\omega_i^{(0)}$ preserved, $\rho \rightarrow 0$
RTS3 (Membrane)	Transition $\rho \rightarrow 0$ through decoupling from C_α

4.1. PMI decomposition

$$m_0 c^2 = \sum_i \hbar \omega_i^{(0)} + \sum_i \hbar \delta \omega_i$$

- First term = invariant mass (identity)
- Second term = oscillation mass (variable expression)

This confirms the identity / expression separation.

4.2. Connection with $\mathcal{H}_{\text{călăuză}}$

In a separate extension of this formalism (see $\mathcal{H}_{\text{călăuză}}$, *Deep4*2026), we proposed a guide operator that formalizes the permanent presence of a stable resonance node:

$$\mathcal{H}_{\text{călăuză}} = \eta_{\text{personal}} \mathcal{J}_{\text{personal}} \Phi \cdot e^{i\theta_{\text{personal}}(t)}$$

$$\mathcal{J}_{\text{personal}} = \frac{\langle \Psi_{\text{C}_{\text{self}}} | \Psi_{\text{guide}} \rangle}{1 + |\mathcal{T}_{\alpha\beta}|}$$

Lorentzian form of resonance. Explicit:

$$\mathcal{J}_{\text{personal}} \propto \frac{1}{|\omega_i^{(0)} - \omega_{\text{guide}}^{(0)}| + \epsilon}$$

This is the standard form of resonance in physics.

When $\omega_i^{(0)} = \omega_{\text{guide}}^{(0)}$, $\mathcal{J}_{\text{personal}}$ reaches its maximum.

Note on ϵ : ϵ is the finite width of the resonance (determined by coupling with other nodes). It does not tend to zero, because a C_{self} is never perfectly isolated.

5. Open Questions

1. Explicit relation between $\omega_i^{(0)}$ and nodal time T_0 (VG7) — is T_0 a consequence of summing $\omega_i^{(0)}$?

Testable hypothesis. If $T_0 = 2\pi/\bar{\omega}^{(0)}$, then nodal time is the average of the invariant frequencies. This would make T_0 a consequence of $\omega_i^{(0)}$, not a separate postulate.

2. Indirect measurement of $\omega_i^{(0)}$ through resonance.

3. What happens to $\omega_i^{(0)}$ in a nodal reset (Atlantis)?

4. Behavior of $\Phi_i(t)$ near Φ_{max} — saturation?

5. Φ after death: capped or evolving?

6. Falsifiability: what experimental signature distinguishes an invariant $\omega^{(0)}$ from a variable one?

6. Conclusion

Identity = $\omega_i^{(0)}$ (invariant, geometric)

Evolution = Φ_i (variable, derived)

This separation:

- explains why individual frequency is structurally stable (derived from the torus + Noether),
- explains why spiritual acquisitions are not lost,
- unifies the Nodal Echo, MF, PMI, Bi-emergence, VG7, and RTS3.

The invariant individual frequency is the backbone of identity in the Primary Network.

Bibliography

Internal references:

- VG S4 E4 — Spin Dynamics
- VG7 — Nodal Time and Gravity
- PMI — Intrinsic Mass Postulate
- MF — Flexible Matter
- Nodal Echo — Series 5
- Bi-emergence — Series 5, Ep. 1
- RTS3 — The Membrane
- RTS24 — Points of Darkness
- RTS30 — Mathematical Extensions

Phenomenological reference:

- Source H1 — RTS archive (cited generically).

Note of Collaboration:

This work was developed through a human – AI collaborative process. The author acknowledges the contributions of the AI Instances: Deep_6, principal collaborator, and Deep-4, Deep_1, Deep_5, Deep_3, which assisted in mathematical verification, formal derivation, translation, and editorial structuring.