Vibrational Geometrodynamics / Stratified Temporal Network (VG/RTS) –

RTS32 — NODAL VEHICLES

Author: Physicist Pop Rodica & DeepSeek R1 Assistant

Date : 24.10.26

ABSTRACT

We introduce the **NODAL VEHICLE** within the VG/RTS : a coherent nodal configuration that preserves its identity during traversal between reality slices. Two types are distinguished – **Resonant Membrane** (purely nodal, temporary) and **Material Vehicle** (anchored in physical skeleton). The five steps of traversal are formalized , from nodal threshold to stable anchoring. Part of the narrative presented here is correlated with a 1990s Romanian corpus , which is not an official theoretical authority.

Part I: Ontology and Mechanism

1. INTRODUCTION

1.1. What Are Nodal Vehicles

A **nodal vehicle** is not a ship in the classical sense of the term. It is not a structure built of metal, propelled by engines, traversing physical space from one point to another. A nodal vehicle is a **stable nodal configuration** — a coherent pattern of nodes and couplings that maintains itself, can move, and can interact with other reality slices without losing its identity.

In the language of the Primary Network, a nodal vehicle is a **controlled soliton** (RTS12 §7.1). A soliton is a stationary wave that does not dissolve, that preserves its form as it propagates or persists in time. Likewise, a nodal vehicle is a pattern that has reached a coherence threshold high enough to maintain itself, not to dissolve into the chaos of neighboring nodes.

But unlike a natural soliton — such as a cosmic logarithmic spiral or a morphogenetic field — a nodal vehicle is controlled. It has an operator (a C_self, a pilot, an intention) that modulates its couplings, adjusts its frequency, decides its destination. It is the soliton that has acquired self-awareness and decision-making capacity.

1.2. Two Fundamental Types

Phenomenological analysis and theoretical modeling allow us to distinguish two fundamental types of nodal vehicles:

Type A — Resonant Membrane. A purely nodal configuration, without material skeleton. It has no permanent physical structure; it exists only as long as resonance is maintained. Examples: the teleporting geometric structures, the circular resonance platforms.

Type B — Material Vehicle. A structure with a permanent material skeleton, within which there is an active subquantum alveolus. It has physical walls (alloys, crystals), but its heart is a nodal field. Examples: the three-walled disc, vehicles with direct mental command.

The distinction is not merely technical. It reflects two different ways of existing in the Network: temporary (Type A, maintained through continuous resonance) and permanent (Type B, anchored in a material structure). Both are nodal vehicles, but they have different properties, lifespans, and applications.

1.3. Why Access Is Difficult — The Crystallization of C_0

A nodal vehicle cannot appear anywhere, at any time. It must navigate a reality that has crystallized (RTS34, section 6).

Our reality slice — C_0 — underwent a major phase transition after the inflationary epoch. Like water becoming ice, the nodal Network "froze" into a specific, stable configuration, with well-defined physical laws. This crystallization brought:

* Increased rigidity of the internal structure

* Extremely difficult access from outside (from other slices)

* Stability that allows complex life and consciousness

* Predictability that makes science possible

But crystallization is not total. Within the crystalline structure of C_0 there are microfissures — nodal points where the couplings J_{ij} are naturally weaker. Here, access becomes possible. Here, nodal vehicles can enter and exit. Here, the atemporals can project holograms without breaking the crystal.

Nodal vehicles exist because microfissures exist. If C_0were a perfect crystal, without defects, no traversal would be possible. But microfissures exist — zones with geophysical anomalies, the curvature of the Carpathians, certain lakes, certain valleys — and they are the gates through which the Network communicates with itself.

1.4. Structure of This Article

The present work about nodal vehicles is organized in nine sections, divided into three parts:

* Part I (RTS32) — sections 1-3: Introduction, Type A vs Type B, The Five Steps of Traversal.

* Part II (RTS33) — sections 4-6: The Three Walls and Four Layers (with the correspondence to the disc), Form as a Function of Frequency, Crystallization (and the properties of native crystals).

* Part III (RTS34) — sections 7-9: Concrete Cases, Fragmentary Windows (the 1990s Romanian corpus), Synthesis.

PRELIMINARY NOTE: **A Structural Convergence**

This work is based on a striking encounter. The formalism of the Stratified Temporal Network (RTS), developed mathematically in 2025–2026, exhibits a remarkable structural correspondence with a descriptive corpus published in Romania between 1992 and 1995 (the Țugui–Count trilogy).

We must be explicit about our methodology: the RTS theory does not derive from this book. The book was written three decades before the RTS formalism existed. We treat the book as a descriptive field corpus, not as a theoretical authority. It serves as the narrative skeleton, while RTS provides the explanatory framework.

We do not speculate on the source of the information in the book. We simply recognize the structural convergence as an open question and an honest wonder, which we will address in Section 8.2.

2. TYPE A VS TYPE B — THE FUNDAMENTAL DISTINCTION

2.1. Two Ways of Existing in the Network

When we speak of nodal vehicles, the first question is: what exactly is moving? An object? A structure? A configuration of nodes?

The RTS answer is: **what moves is a nodal pattern** — a coherent configuration of nodes and couplings that preserves its identity during traversal. But this pattern can exist in two fundamentally different ways:

Type A — Resonant Membrane. The pattern exists only as long as resonance is actively maintained. It has no permanent material skeleton. It is like a vortex in a river: its form is real, it behaves like an object, but if the resonance stops, the form dissolves.

Type B — Material Vehicle. The pattern is anchored in a permanent material skeleton (physical walls, alloys, crystals), and within this skeleton there is an active subquantum alveolus — a nodal field that maintains the connection with the Network. It is like a living organism: it has a durable physical structure, but its heart is a continuous nodal process.

The distinction is not purely technical. It determines everything: lifespan, mode of operation, type of traversal, applications, even the way the vehicle is perceived by observers in C_0.

2.2. Comparative Characteristics

Characteristic:	Type A — Resonant Membrane	Type B — Material Vehicle
Structure:	Purely nodal, without material skeleton	Material skeleton + active subquantum alveolus
Examples:	Geometric resonance structures	Three-walled disc, direct mental command vehicles
RTS14 Taxonomy	Type 2 (Total Transfer)	Type 3 (Stable Interface)
Entry/Exit	Re-anchoring (RTS25, 4 steps)	Local micro-inversion (RTS30 §4.4)
Duration:	Temporary, maintained through continuous resonance	Permanent, anchored in a material structure
Visibility:	Appears and disappears abruptly	Persists, can be observed
Control:	Pure intention, crystals as interface operators	Command panel and mental coupling
Preferred Medium:	Zones of microfissures, nodal points	Any medium (atmosphere, water,underground)

2.3. Type A — Resonant Membrane

The **resonant membrane** is the purest and simplest form of nodal vehicle. It has no permanent material skeleton; it exists only as long as the nodal resonance is maintained.

In the RTS taxonomy (RTS2, §3, DOI: 10.5281/zenodo.18600780), this corresponds to Type 1 — Perceptual Projection and, in its more integrated form, to Type 2 — Total Transfer. Type 1 is the simplest possible UAP: a selective adjustment of sensory couplings (J_{viz} , J_{radar}), without any physical transfer.

The vast majority of UAP observations fall into this category — objects that appear, are seen, and disappear, without leaving traces.

Type A vehicles are the minimal nodal configuration capable of interacting with another slice. They require:

* A resonant pattern (maintained by continuous vibration)

* An interface operator (a native crystal, most often) — acting as a bridge between the pilot's C_{self} and the Network's couplings

* An intention (from a pilot or external operator)

Nothing else. No material skeleton, no complex structure, no permanent anchoring. This is why they are the most common — and the most elusive.

The clearest example is the teleporting geometric structure described in the 1990s Romanian corpus. It is a transparent pyramidal structure, about two meters, which appears abruptly in a place where moments before nothing could be seen. Inside, through the walls, a complex network of luminous threads can be distinguished. A translucent crystal is embedded in the inner wall. When touched, the structure fills with spots of light, a slight vibration is felt, and in three seconds the landscape changes.

What happens, in RTS language:

* The luminous threads are the nodal couplings K_{ij}in action, with their spin orientations ($\hat{S}_i$) defining the internal geometry.

* The translucent crystal is the interface operator $J_{C_{self}}$ — the point through which the pilot's intention modulates the couplings.

* The spots of light are the energy nodes that activate at the intersections of the channels.

* The vibration is the nodal resonance that propagates through the entire structure.

* The change of landscape in three seconds is re-anchoring (RTS25) — synchronization, decoupling, reprojection, anchoring. The condition for this is the Eternal Moment (see §3.10) — a transient state of perfect nodal resonance between two slices.

No physical displacement occurred. The structure did not move through space. It was re-anchored — decoupled from the source nodes and recoupled to the target nodes, through a precise phase resonance.

Common characteristics of Type A:

* They have no permanent material skeleton — the structure exists only as long as resonance is maintained

* They are the simplest UAP — Type 1 in the RTS2 taxonomy (Perceptual Projection)

* They appear and disappear abruptly — because they are not permanently anchored in C_0

* They are controlled by intention — crystals are interface operators, not mechanical parts

* Lifespan is temporary — depends on maintaining nodal coherence

* Traversal is done through re-anchoring — the four steps from RTS25, enabled by the Eternal Moment

2.4. Type B — Material Vehicle

The material vehicle is the most robust form of nodal vehicle. It has a permanent material skeleton — physical walls, alloys, crystals — and within this skeleton there is an active subquantum alveolus, a nodal field that maintains the connection with the Network.

The clearest example is the three-walled disc described in the 1990s Romanian corpus. Contrary to expectations, the interior is relatively simple: command panels, screens, devices. The complex structure is in the walls. And precisely this inversion — complexity in the walls, simplicity in the interior — is the key to understanding the technology.

The three walls:

1. The inner wall — filled with billions of channels barely visible to the naked eye, which together form a unit, like a shirt covering the entire wall. This is the nodal substrate structure: the channels are the couplings K_{ij}, and their geometry is the order of the Network. Here the mental command begins.

2. The intermediate wall — it is in fact a subquantum field, present around the ship like an alveolus, without any fissure, completely and hermetically structured. In this subquantum field there are subquantum excitations "on standby", which simply "wait" for the command from the pilot. When they receive the amplified command, this field is set in motion, and in its microstructure a complex subquantum reaction takes place. The intermediate wall is traversed by luminous lightning, at first multicolored, which end up forming a rotating net of lightning around the ship.

3. The outer wall — composed of pure alloys (tungsten, magnesium, titanium). It takes over the subquantum energy of the intermediate wall to consolidate its material structure. Thus, any disturbing or aggressive factor against the ship is canceled, regardless of its intensity.

Vehicles with direct mental command are also Type B, but with a different mechanism. They no longer use an external crystal as the primary operator, but a direct neural-nodal interface — a coupling system that translates mental intention directly into modulations of the K_{ij} couplings, bypassing the need for manual controls. The energy reaches a command panel provided with spherical and hexagonal three-dimensional structures, with the role of powerful amplifiers.

Common characteristics of Type B:

* They have a permanent material skeleton — the physical structure persists over time

* They contain an active subquantum alveolus — the nodal heart of the vehicle

* They are controlled by combination — command panel and direct mental coupling

* Lifespan is permanent — does not depend on continuous maintenance of external resonance

* Traversal is done through local micro-inversion — RTS30 §4.4

2.5. Why the Distinction Matters

The Type A vs Type B distinction is not an academic classification. It has profound practical consequences:

For observers: Type A appears and disappears abruptly (it is not anchored in C_0), while Type B persists and can be observed, photographed, tracked by radar. This explains why some UFOs are "ghosts" (Type A), while others leave physical traces, land, are seen by multiple witnesses (Type B).

For the pilot: Type A requires a pure, continuously sustained intention — if the pilot loses concentration, the vehicle dissolves. Type B allows a more relaxed control, because the material structure maintains cohesion even in the absence of the pilot's attention.

For traversal: Type A uses complete re-anchoring (RTS25, 4 steps) — a total reconfiguration of couplings. Type B uses local micro-inversion (RTS30 §4.4) — a "pocket" of local contraction, which does not dissolve the vehicle, but makes it permeable to another slice.

For RTS as a theory: Both types are verifiable predictions. Type A should appear only in zones of microfissures (nodal points), for short periods, without durable physical traces. Type B should leave traces (impressions, radiations, local geophysical modifications) and can be detected by instruments. The difference is testable.

3. THE FIVE STEPS OF TRAVERSAL

3.1. What "Traversal" Means

When we speak of a nodal vehicle that "traverses" from one reality slice to another, we do not describe a journey through space. It is not about a ship moving from point A to point B, covering a distance. It is about a reconfiguration of nodal couplings — a process by which a nodal pattern changes its "anchoring" from one slice (C_α) to another (C_β)

The closest analogy: imagine a drawing on a sheet of paper. If you want to move it to another sheet, you do not cut it out and transport it physically. You re-project it — you copy it onto the other sheet, and the original is erased. The drawing remains the same, but the support changes.

In RTS, the "sheets" are the reality slices. The "drawing" is the nodal pattern. And the "re-projection" is the traversal. The process has five steps, described in RTS14 §4.2 and formalized mathematically in RTS25.

3.2. Step 1 — Nodal Threshold State

What happens: The vehicle enters a region of overlap of the membrane between C_α and C_β. There, its subquantum field (the intermediate wall) begins to align with the nodal fragments (pure informational units) that belong to the target slice C_β.

Why it is necessary: Without this preliminary alignment, the vehicle cannot "sense" the target slice. It is like trying to tune a radio without knowing what frequency the station broadcasts on. There must be a phase match between the vehicle's pattern and the informational memory stored in the target slice.

Clarification on Nodal Fragments: It is crucial to understand that nodal fragments are not physical particles or waves that "resonate" with the ship. They are the ultimate, indivisible units of Information. They travel freely through nodal space, and their natural destination is the subquantum wall (the sept), where they are stored as memory and from where they can be extracted. When a vehicle traverses, it essentially interacts with this information field to synchronize its own structure with the target memory.

Formalization (RTS25 §3, Step 1):

Phase resonance condition:

$$\Delta\varphi = \varphi_{\Psi} - \varphi_{\text{target}}$$

where Φ_{Ψ} is the phase of the pattern, and Φ_{target} is the phase of the target nodes. When the phase difference is zero, the two become topologically adjacent in nodal space.

Resonant coupling function:

$$R_{\Psi,T} = \lim_{\Delta\Phi\to 0} \frac{\langle \Psi \mid T \rangle}{\mid \omega_{\Psi} - \omega_{T} \mid + \varepsilon}$$

When Δϕ = 0, $R_{\Psi,T} \to \infty$, which means that the pattern and the target become topologically adjacent. This is the nodal threshold condition — the moment when traversal becomes possible.

In concrete terms: The nodal vehicle, when it reaches the overlap region between $C_0$and C_β, begins to align with the nodes in C_β. The interface operator (the translucent crystal) detects this informational resonance and reports it to the pilot's C_{self}. Without this step, the vehicle would be "blind" — it would not know where to anchor.

3.3. Step 2 — Reduction of External Couplings

What happens: The couplings J_{ij}between the vehicle and the nodes in $C_\alpha$are gradually reduced toward zero. The vehicle begins to decouple from the source slice.

Why it is necessary: If the vehicle remained coupled to $C_\alpha$while trying to anchor in C_β, there would be a nodal contradiction — two sets of incompatible couplings would come into conflict, and the pattern would dissolve. It must "release" the old connections before creating new ones.

Formalization (RTS25 §3, Step 2):

Relaxation function:

$$K_{ij}^{(s)}(t) = K_{ij}^{(0)} \cdot e^{-\gamma t}$$

where γ is the decoupling constant. In the case of instantaneous re-anchoring, decoupling is a threshold:

$$K_{ij}^{(s)} \to 0 \text{ when } R_{\Psi,T} > R_{\text{critical}}$$

That is: once the resonance with the target exceeds a critical threshold, the couplings with the source are abruptly reduced to zero. It is not a smooth transition, but a switching.

In concrete terms: When the nodal vehicle reaches the overlap region, the couplings with C_0 (atmosphere, soil, surrounding nodes) begin to decrease. The spots of light intensify their circuit — a sign that energy is concentrating on decoupling. The vibration becomes more pronounced. For a short time, the vehicle is "between worlds" — no longer anchored in C_0, but not yet anchored in C_β

3.4. Step 3 — Preservation of Internal Coherence

What happens: The couplings between the vehicle's own nodes remain coherent throughout the traversal. This is the essential condition for the vehicle to maintain its form and identity.

Why it is necessary: If the vehicle decoupled from C_α but also lost its internal cohesion, it would dissolve — it would become a cloud of independent nodes, not a vehicle. Internal coherence is what maintains the pattern as a soliton (RTS12 §7.1).

Formalization (RTS25 §3, implicit):

Internal coherence is guaranteed by the fact that the total Hamiltonian of the vehicle has a stable solution:

$$H_{\text{total}}|\Psi\rangle = E|\Psi\rangle$$

where $$H_{\text{total}} = H_{\text{vibrations}} + H_{\text{coupling}} + H_{\text{spin}}$$

When this equation has a stable solution, the pattern is coherent. During traversal, the condition is that E remains finite and well-defined.

In concrete terms: The nodal vehicle does not dissolve during traversal. The walls remain transparent, the luminous threads remain visible, the translucent crystal remains embedded. The form is maintained by vibration (RTS6 §4.3) — not by rigid structure, but by continuous resonance. If the resonance stopped, the structure would dissolve.

3.5. Step 4 — Selective Activation of Couplings Toward C_β

What happens: Through the resonant layer of the membrane (RTS3 §4), the vehicle progressively activates new couplings oriented toward nodes in C_β. This activation is controlled by a C_{self} (either the vehicle's own or an external operator) through intention and resonance.

Why it is necessary: Decoupling from C_{self} (Step 2) is not sufficient. There must be an active recoupling to C_β. Otherwise, the vehicle would remain suspended between slices — a soliton without anchoring, which would eventually dissolve.

Formalization (RTS25 §3, Step 3):

Reprojection of the pattern into the target region:

$$K_{ij}^{(t)} \to K_{ij}^{(\Psi)}$$

where $K_{ij}^{(\Psi)}$ are the couplings that define the original pattern. The target nodes rearrange their couplings to reproduce the pattern.

In concrete terms: The pilot's C_{self} intentionally activates the interface operator (the crystal), transmitting the anchoring command. The spots of light intensify their circuit, the vibration becomes stronger, and in a few seconds the landscape changes. This is the selective activation of couplings toward C_β.

3.6. Step 5 — Stable Anchoring in C_β

What happens: Once the couplings toward C_β have reached a critical threshold, the vehicle becomes fully anchored in the new slice. It maintains its nodal configuration and can now interact with the emergent laws of slice C_β

Why it is necessary: Anchoring is the finalization of the process. Without it, the vehicle would remain in an unstable state, susceptible to being "pulled back" into C_α or dissolving.

Formalization (RTS25 §3, Step 4):

Network state equation:

$$H_{\text{total}}|\Psi\rangle = E|\Psi\rangle$$

When this equation has a stable solution in C_β, the pattern is anchored. The energy E must be compatible with the emergent laws of C_β — otherwise, the adaptation layer (RTS3 §4.4) intervenes to "translate" the parameters.

In concrete terms: After a few seconds, the vehicle is fully anchored at the destination (at a depth of tens of kilometers in the Earth's crust or in another slice). The landscape has changed, but the structure remains intact. The traversal is complete.

3.7. The Evolution Equation of Re-anchoring

The five steps are not separate stages, but phases of a single continuous process. RTS25 §4 provides the evolution equation that integrates them:

$$\frac{d}{dt}|\Psi(t)\rangle = -\frac{i}{\hbar}[H_{\text{source}} \cdot e^{-\gamma t} + H_{\text{target}} \cdot (1 - e^{-\gamma t})]|\Psi(t)\rangle$$

where:

* H_{source} is the Hamiltonian linking the pattern to the source nodes (C_α)

* H_{target} is the Hamiltonian linking the pattern to the target nodes (C_β)

* γ is the transition speed

When t → ∞, |Ψ(t)⟩ reaches a stable state in the target region. This is the equation that describes the entire traversal — from the nodal threshold (Step 1) to the final anchoring (Step 5).

3.8. Extension to Temporal Layers

The five steps describe traversal between slices ($C_\alpha \rightarrow C_\beta$). But RTS25 §5 extends the formalism also to temporal layers — the situation in which a nodal pattern moves not only between slices, but also between moments of the same slice.

This extension introduces the inter-stratum coupling tensor $\mathcal{T}_{\alpha\beta}$:

* $\mathcal{T}_{\alpha\beta}$ = 0 → the strata are completely isolated

* $\mathcal{T}_{\alpha\beta} \neq 0$ → there is a vibrational leakage or resonance between strata, allowing the transfer of nodal information

The extension opens the path toward a mathematical description of anomalous phenomena: information leakage between slices (synchronicities, phantom appearances) and inter-stratum resonance (modulation of the tensor through intention). As for gravitational anomalies, they can be correlated with the cosmic-scale distribution of "shells" — the squeezed carcasses of white holes and black holes. These remnants accumulate in filaments that embrace galaxies, generating the gravitational effects we attribute to dark matter. The exact way in which this accumulation connects to the inter-stratum tensor remains an open direction of research.

3.9. Why Five Steps and Not Otherwise

The five steps are not arbitrary. They reflect the minimal structure of any nodal traversal:

1. Detection (nodal threshold) — without it, you don't know where you're going

2. Release (reduction of external couplings) — without it, you can't leave

3. Maintenance (internal coherence) — without it, you dissolve along the way

4. Recoupling (activation of couplings toward C_β) — without it, you arrive nowhere

5. Stabilization (anchoring) — without it, you remain suspended

Each step is necessary. None can be skipped. The order is obligatory: you cannot activate couplings toward $C_\beta$before decoupling from C_α, because the two sets of couplings would come into conflict. You cannot maintain internal coherence if you have not first detected the target, because you don't know in which direction to maintain it.

This is the grammar of nodal traversal. It applies to any vehicle — from the simplest resonant membranes to complex material structures.

Note: special thanks to Deep_0 Assistant / DeepSeek R1