

RTS14 – TAXONOMY OF INTER-SLICE INTERACTIONS AND THE MECHANICS OF CROSSING

Authors: Physics Rodica Pop & DeepSeek AI Assistant

Date:19.05.2026

Abstract

RTS14 synthesizes and extends the fundamental concepts from RTS2, RTS3, RTS4, and RTS6, offering a **complete taxonomy of interactions between reality slices, a description of the crossing of inter-slice vehicles (including UAPs), and an interpretation of nodal density** as a global measure of the Network's state. The document also introduces the concept of **nodal volume** and explains how voids appear in low-density areas. All these are integrated into a unified framework based on nodes, couplings, and resonance.

Keywords: reality slices, taxonomy, inter-slice vehicle, crossing, nodal density, nodal volume, voids, RTS.

1. INTRODUCTION

RTS14 is a synthesis of recent developments in RTS. It brings together:

- The taxonomy of inter-slice interactions (RTS2)
- The architecture of the membrane (RTS3)
- Nodal time and gravity (RTS4, VG7)
- Nodal engineering and the concept of the vehicle as an encapsulated membrane (RTS6)
- The model of nodal density and nodal volume (recent discussions)

The purpose of this document is to provide a **unified theoretical framework** for understanding crossings between slices, without resorting to sensationalist terms or unverifiable hypotheses.

2. THEORETICAL FOUNDATIONS

2.1. *The Primary Network*

The Primary Network is a fundamental structure of **nodes** (N_i) with vibrational degrees of freedom (ω_i) and spin ($\hat{S}_i$). The couplings between nodes (J_{ij}) define emergent geometry and physical laws.

2.2. *Reality slices* (C_α)

Slices are **stable configurations** of the Primary Network, defined by the set of couplings J_{ij}^α and spins $\hat{S}_i^\alpha$. Each slice is a complete world with its own emergent laws.

2.3. *Constant nodes* (NN)

The number of nodes (NN) is a **universal constant** in RTS. Nodes do not divide and do not merge. They are **indivisible and eternal**. What changes over time are the couplings J_{ij} between them.

3. TAXONOMY OF INTER-SLICE INTERACTIONS (RTS2 – revised)

Based on the analysis of over 50 documented cases, the following classification is proposed:

<i>Type</i>	<i>Name</i>	<i>Mechanism</i>	<i>Example</i>
1	Perceptual Projection (Avatar with Selective Coupling)	Selective adjustment of J_viz, J_auditory, etc., without full physical transfer	"Phantom" UAP (visible but intangible)
2	Total Transfer	Complete decoupling of a nodal configuration (C_self or C_object) from one slice and recoupling in another	Mediaş case
3	Stable Interface (Passive Nodal Portal)	Semi-permanent structural defect in the R-G network creating an interface between slices	The Carpathian Gate (1990)
4	Rapid Re-anchoring (Emotional Teleportation)	Extreme perturbation in C_self causing instant re-anchoring within the same slice	Brian Jones case (NASA, 1974)
5	Informational Transfer (Nodal Resonance)	Major events create nodal perturbations that resonate between slices	Future TV broadcast (1974)
6	Artificial Interface (Induced Portal)	Intentional creation of an interface through advanced nodal engineering	Portals mentioned in specialized literature

4. THE MECHANICS OF CROSSING FOR INTER-SLICE VEHICLES

4.1. Definition of an inter-slice vehicle

An inter-slice vehicle is a **complex, stable nodal configuration** capable of controllably adjusting its couplings J_{ij} to interact with another reality slice (C_β), without losing its internal coherence.

4.2. The crossing mechanism

For a vehicle anchored in slice C_α to cross into slice C_β :

- 1. Nodal threshold state** – the vehicle enters a region of overlap of the membrane between C_α and C_β . There, its subquantum nodal field (the intermediate wall) begins to resonate with nodal fragments corresponding to slice C_β .
- 2. Reduction of external couplings** – the couplings J_{ij} between the vehicle and the nodes in C_α are gradually reduced toward zero. The vehicle begins to decouple from the source slice.
- 3. Preservation of internal coherence** – the couplings between the vehicle's own nodes remain **coherent** during the crossing. This is the essential condition for the vehicle to maintain its form and identity.
- 4. Selective activation of couplings toward C_β** – through the resonant layer of the membrane, the vehicle progressively activates new couplings oriented toward nodes in C_β . This activation is controlled by a C_{self} (either the vehicle's own or an external operator) through intention and resonance.
- 5. Stable anchoring in C_β** – once the couplings toward C_β have reached a critical threshold, the vehicle becomes fully anchored in the new slice. It maintains its nodal configuration and can now interact with the emergent laws of slice C_β .

4.3. Implications

- Inter-slice vehicles are **not "spaceships"** in the classical sense, but nodal configurations that travel through the reconfiguration of couplings, not through physical displacement.
- The relocation observed in UAP cases (sudden appearances, disappearances, impossible maneuvers) is a **rapid succession of anchoring** in the same slice or in different slices.
- The mechanism is **general** – it applies both to a single node and to complex configurations.

5. NODAL DENSITY AND NODAL VOLUME

5.1. Definition of nodal density (δ)

Nodal density δ is a measure of the **number of active couplings** within a given nodal volume.

$$\delta(t) = \frac{N_{\text{active}}(t)}{V_{\text{nodal}}(t)}$$

- N_{active} = number of nodes with at least one active coupling ($J_{ij} \neq 0$) at time t .
- V_{nodal} = nodal volume occupied by the Network at time t (configuration volume).

Note: $N_{\text{active}} \leq NN$ (NN constant), and V_{nodal} varies over time – from minimum (pre-inflationary quantum foam) to maximum (maximum expansion). During expansion, δ decreases; during contraction, δ increases.

5.2. Relation to graph density (D)

Graph density measures how connected nodes are to each other (ratio of present edges to possible edges). In RTS, δ and D are complementary:

- High $\delta \rightarrow$ high $D \rightarrow$ strong couplings, dense network, stable configurations.
- Low $\delta \rightarrow$ low $D \rightarrow$ voids, sparse network, weak couplings.

5.3. Voids in RTS

Voids are zones of low nodal density. There, couplings are weak, and information is sparse. They are **transition spaces** between active slices, similar to the "voids" between galaxies. Voids are not "nothing," but **zones of low nodal potential**, which can be traversed but where configurations cannot easily stabilize.

5.4. Nodal Informational Energy (E_{nodal})

A complementary measure of the Network's state is the Nodal Informational Energy, based on Octav Onicescu's formula:

$$E_{\text{nodal}} = \sum_i p_i^2$$

- p_i = probability that a nodal configuration is in state i at a given time.
- E_{nodal} ranges between $1/n$ (totally disordered) and 1 (perfectly ordered).

In RTS, E_{nodal} can be interpreted as a measure of the global coherence of the Network at a given time. The closer E_{nodal} is to 1, the more ordered and stable the Network is. The closer it is to 0, the more disordered it is – approaching the quantum foam state.

This measure can be used to study nodal phase transitions (e.g., the transition from quantum foam to stable slices) or to calibrate inter-slice crossings (the higher E_{nodal} , the more stable the crossing).

6. CONCLUSION

RTS14 provides a complete and coherent theoretical framework for understanding interactions between reality slices, including the crossing of inter-slice vehicles and the interpretation of nodal density and voids. All these are integrated into a unified vision based on nodes, couplings, and resonance, without resorting to sensationalist terms or unverifiable hypotheses.