

# Series of functions

G. Goron

EPITA

October 31, 2021



## Why are we interested in series?

We now look at the notion of convergence of a *series* of functions  $\sum f_n$ , where the  $f_n$  are defined over an interval  $I$  of  $\mathbb{R}$ .

## Why are we interested in series?

We now look at the notion of convergence of a *series* of functions  $\sum f_n$ , where the  $f_n$  are defined over an interval  $I$  of  $\mathbb{R}$ .

The approximation by a series of functions is a widely-used technique, you already encountered such things when dealing with Taylor expansions, power series, and Fourier series mainly.

## Why are we interested in series?

We now look at the notion of convergence of a *series* of functions  $\sum f_n$ , where the  $f_n$  are defined over an interval  $I$  of  $\mathbb{R}$ .

The approximation by a series of functions is a widely-used technique, you already encountered such things when dealing with Taylor expansions, power series, and Fourier series mainly.

By definition, a series of functions is at core a sequence of functions. Thus, the notions of pointwise and uniform convergence remain the same and present the same interests.

## Why are we interested in series?

We now look at the notion of convergence of a *series* of functions  $\sum f_n$ , where the  $f_n$  are defined over an interval  $I$  of  $\mathbb{R}$ .

The approximation by a series of functions is a widely-used technique, you already encountered such things when dealing with Taylor expansions, power series, and Fourier series mainly.

By definition, a series of functions is at core a sequence of functions. Thus, the notions of pointwise and uniform convergence remain the same and present the same interests.

However, the fact that we know series by their associated sequence opens a new path to explore the convergences, while it also makes it more difficult to use the methods we used to work with when dealing with plain sequences.

## Why are we interested in series?

We now look at the notion of convergence of a *series* of functions  $\sum f_n$ , where the  $f_n$  are defined over an interval  $I$  of  $\mathbb{R}$ .

The approximation by a series of functions is a widely-used technique, you already encountered such things when dealing with Taylor expansions, power series, and Fourier series mainly.

By definition, a series of functions is at core a sequence of functions. Thus, the notions of pointwise and uniform convergence remain the same and present the same interests.

However, the fact that we know series by their associated sequence opens a new path to explore the convergences, while it also makes it more difficult to use the methods we used to work with when dealing with plain sequences.

So, we have to devise new tools that are specific to series in order to prove in a swifter way that our requirements to use the wanted calculation properties are met.





## Pointwise convergence ( $\Sigma$ )

The notion of pointwise convergence is the same for series and sequences, if we consider that a series is nothing but a sequence of partial sums.

## Pointwise convergence ( $\Sigma$ )

The notion of pointwise convergence is the same for series and sequences, if we consider that a series is nothing but a sequence of partial sums.

However, here is quite a hindrance: we generally do not know how to calculate the limit of a numerical series.

## Pointwise convergence ( $\Sigma$ )

The notion of pointwise convergence is the same for series and sequences, if we consider that a series is nothing but a sequence of partial sums.

However, here is quite a hindrance: **we generally do not know how to calculate the limit of a numerical series.**

So far, you have only be taught how to assert or invalidate the convergence of a numerical series, yet without (almost) ever being able to determine towards which value it was converging.

## Pointwise convergence ( $\Sigma$ )

The notion of pointwise convergence is the same for series and sequences, if we consider that a series is nothing but a sequence of partial sums.

However, here is quite a hindrance: **we generally do not know how to calculate the limit of a numerical series.**

So far, you have only be taught how to assert or invalidate the convergence of a numerical series, yet without (almost) ever being able to determine towards which value it was converging.

Not knowing how to easily calculate the limit of a numerical series, and of course of a series of functions, makes it really hard to check if its convergence is uniform by evaluating the difference between the partial sum and the limit in infinite norm. We have to get around this problem.

## Uniform convergence ( $\sum$ )

Likewise, the notion of uniform convergence is similar to that of sequences of functions: we say a series of functions  $\sum f_n$  to be uniformly convergent over  $I$  towards its limit  $\mathcal{S}$  iff:

$$\sup_{x \in I} \left| \left( \sum_{i=0}^n f_n(x) \right) - \mathcal{S}(x) \right| \xrightarrow{n \rightarrow +\infty} 0, \quad \text{i.e. :} \quad \left\| \left( \sum_{i=0}^n f_n \right) - \mathcal{S} \right\|_{\infty} \xrightarrow{n \rightarrow +\infty} 0.$$

## Uniform convergence ( $\sum$ )

Likewise, the notion of uniform convergence is similar to that of sequences of functions: we say a series of functions  $\sum f_n$  to be uniformly convergent over  $I$  towards its limit  $\mathcal{S}$  iff:

$$\sup_{x \in I} \left| \left( \sum_{i=0}^n f_n(x) \right) - \mathcal{S}(x) \right| \xrightarrow{n \rightarrow +\infty} 0, \quad \text{i.e. :} \quad \left\| \left( \sum_{i=0}^n f_n \right) - \mathcal{S} \right\|_{\infty} \xrightarrow{n \rightarrow +\infty} 0.$$

Using for the remainder the notation  $R_n(x) = \sum_{i=n+1}^{\infty} f_n(x) = \mathcal{S}(x) - \sum_{i=0}^n f_n(x)$  we get:

$$\sum f_n \text{ converges uniformly over } I \text{ iff } \|R_n\|_{\infty} \xrightarrow{n \rightarrow +\infty} 0.$$

## Uniform convergence ( $\sum$ )

Likewise, the notion of uniform convergence is similar to that of sequences of functions: we say a series of functions  $\sum f_n$  to be uniformly convergent over  $I$  towards its limit  $\mathcal{S}$  iff:

$$\sup_{x \in I} \left| \left( \sum_{i=0}^n f_n(x) \right) - \mathcal{S}(x) \right| \xrightarrow{n \rightarrow +\infty} 0, \quad \text{i.e. :} \quad \left\| \left( \sum_{i=0}^n f_n \right) - \mathcal{S} \right\|_{\infty} \xrightarrow{n \rightarrow +\infty} 0.$$

Using for the remainder the notation  $R_n(x) = \sum_{i=n+1}^{\infty} f_n(x) = \mathcal{S}(x) - \sum_{i=0}^n f_n(x)$  we get:

$$\sum f_n \text{ converges uniformly over } I \text{ iff } \|R_n\|_{\infty} \xrightarrow{n \rightarrow +\infty} 0.$$

Thus, the series of functions converges uniformly if and only if its remainder  $R_n$  converges uniformly towards the zero function over  $I$ .

## Uniform convergence ( $\sum$ )

Likewise, the notion of uniform convergence is similar to that of sequences of functions: we say a series of functions  $\sum f_n$  to be uniformly convergent over  $I$  towards its limit  $\mathcal{S}$  iff:

$$\sup_{x \in I} \left| \left( \sum_{i=0}^n f_n(x) \right) - \mathcal{S}(x) \right| \xrightarrow{n \rightarrow +\infty} 0, \quad \text{i.e. :} \quad \left\| \left( \sum_{i=0}^n f_n \right) - \mathcal{S} \right\|_{\infty} \xrightarrow{n \rightarrow +\infty} 0.$$

Using for the remainder the notation  $R_n(x) = \sum_{i=n+1}^{\infty} f_n(x) = \mathcal{S}(x) - \sum_{i=0}^n f_n(x)$  we get:

$$\sum f_n \text{ converges uniformly over } I \text{ iff } \|R_n\|_{\infty} \xrightarrow{n \rightarrow +\infty} 0.$$

Thus, the series of functions converges uniformly if and only if its remainder  $R_n$  converges uniformly towards the zero function over  $I$ .

We can then study the uniform convergence of series without having identified their limit, only using their associated sequence.

However, finding upper bounds for the remainders is often quite difficult ...



## Uniform convergence ( $\sum$ ) - alternating series

Let  $(g_n)_{n \in \mathbb{N}}$  be a sequence of functions from  $I$  to  $\mathbb{R}$  such that for every  $x$  in  $I$ , the numerical sequence  $(g_n(x))_{n \in \mathbb{N}}$  is monotonic and tends towards 0.

## Uniform convergence ( $\sum$ ) - alternating series

Let  $(g_n)_{n \in \mathbb{N}}$  be a sequence of functions from  $I$  to  $\mathbb{R}$  such that for every  $x$  in  $I$ , the numerical sequence  $(g_n(x))_{n \in \mathbb{N}}$  is monotonic and tends towards 0.

In this case, Leibniz's test for alternating series enables us to prove that the series  $\sum (-1)^n g_n$  converges pointwise.

## Uniform convergence ( $\sum$ ) - alternating series

Let  $(g_n)_{n \in \mathbb{N}}$  be a sequence of functions from  $I$  to  $\mathbb{R}$  such that for every  $x$  in  $I$ , the numerical sequence  $(g_n(x))_{n \in \mathbb{N}}$  is monotonic and tends towards 0.

In this case, Leibniz's test for alternating series enables us to prove that the series  $\sum (-1)^n g_n$  converges pointwise.

Moreover, for every  $x$  we can find an upper bound for the remainder  $R_n(x)$ : we get indeed

$$\forall x \in I, \forall n \in \mathbb{N}, |R_n(x)| \leq |g_{n+1}(x)|.$$

## Uniform convergence ( $\sum$ ) - alternating series

Let  $(g_n)_{n \in \mathbb{N}}$  be a sequence of functions from  $I$  to  $\mathbb{R}$  such that for every  $x$  in  $I$ , the numerical sequence  $(g_n(x))_{n \in \mathbb{N}}$  is monotonic and tends towards 0.

In this case, Leibniz's test for alternating series enables us to prove that the series  $\sum (-1)^n g_n$  converges pointwise.

Moreover, for every  $x$  we can find an upper bound for the remainder  $R_n(x)$ : we get indeed

$$\forall x \in I, \forall n \in \mathbb{N}, |R_n(x)| \leq |g_{n+1}(x)|.$$

Such an upper bound can help determine whether the convergence is uniform: if  $\|g_n\|_{\infty} \xrightarrow{n \rightarrow +\infty} 0$  over  $I$ , then the convergence is uniform.

## Uniform convergence ( $\sum$ ) - necessary condition

**Proposition** If the series  $\sum f_n$  converges uniformly over  $I$ , then the sequence  $(f_n)$  converges uniformly over  $I$  towards the zero function.

## Uniform convergence ( $\sum$ ) - necessary condition

**Proposition** If the series  $\sum f_n$  converges uniformly over  $I$ , then the sequence  $(f_n)$  converges uniformly over  $I$  towards the zero function.

The reciprocal of this proposition is false: indeed, the uniform convergence of the sequence  $(f_n)$  does not even imply the pointwise convergence of the series.

## Uniform convergence ( $\sum$ ) - necessary condition

**Proposition** If the series  $\sum f_n$  converges uniformly over  $I$ , then the sequence  $(f_n)$  converges uniformly over  $I$  towards the zero function.

The reciprocal of this proposition is false: indeed, the uniform convergence of the sequence  $(f_n)$  does not even imply the pointwise convergence of the series.

For example, if we take  $f_n$  being the constant function equal to  $\frac{1}{n}$  over  $[0,1]$ , the sequence  $(f_n)$  converges uniformly towards the zero function  $\left( \|f_n\|_\infty = \frac{1}{n} \xrightarrow{n \rightarrow +\infty} 0 \right)$  yet for every  $x$ ,  $\sum f_n(x) = \sum \frac{1}{n}$  is a divergent Riemann series.

## Uniform convergence ( $\sum$ ) - necessary condition

**Proposition** If the series  $\sum f_n$  converges uniformly over  $I$ , then the sequence  $(f_n)$  converges uniformly over  $I$  towards the zero function.

The reciprocal of this proposition is false: indeed, the uniform convergence of the sequence  $(f_n)$  does not even imply the pointwise convergence of the series.

For example, if we take  $f_n$  being the constant function equal to  $\frac{1}{n}$  over  $[0,1]$ , the sequence  $(f_n)$  converges uniformly towards the zero function  $\left( \|f_n\|_\infty = \frac{1}{n} \xrightarrow{n \rightarrow +\infty} 0 \right)$  yet for every  $x$ ,  $\sum f_n(x) = \sum \frac{1}{n}$  is a divergent Riemann series.

This result thus gives a necessary but not sufficient condition. It can be used by contraposition to prove that a series does not converge uniformly.



# Uniform convergence ( $\sum$ ) - properties

## Continuity

As for sequences, the interest of uniform convergence is to give sufficient hypotheses to enable the use of results on the limits of series. We will reformulate here the three results we got previously to adapt them to series.

# Uniform convergence ( $\sum$ ) - properties

## Continuity

As for sequences, the interest of uniform convergence is to give sufficient hypotheses to enable the use of results on the limits of series. We will reformulate here the three results we got previously to adapt them to series.

### **Theorem** [Uniform convergence and continuity]

Let  $(f_n)$  be a sequence of **continuous** functions over  $I$ .

We suppose that the series  $\sum f_n$  converges **uniformly** over  $I$ .

Then its limit  $\mathcal{S} = \sum_{n=0}^{+\infty} f_n$  is **continuous** over  $I$ .

# Uniform convergence ( $\sum$ ) - properties

## Swapping sums and integrals

**Theorem** [Swapping sums and integrals]

In this case,  $I$  is a **segment**  $[a, b]$ .

We suppose that  $(f_n)$  is a sequence of continuous functions over  $[a, b]$ .

If the series  $\sum f_n$  converges **uniformly** over  $[a, b]$  then the series  $\sum \int_a^b f_n(t) dt$  converges and:

$$\sum_{n=0}^{+\infty} \int_a^b f_n(t) dt = \int_a^b \left( \sum_{n=0}^{+\infty} f_n(t) \right) dt$$

# Uniform convergence ( $\sum$ ) - properties

## Swapping sums and derivatives

### **Theorem** [Swapping sums and derivatives]

We suppose that:

- ▶ the  $f_n$  functions are of class  $C^1$  over  $I$  ;
- ▶  $\sum f_n$  converges (pointwise) over  $I$  ;
- ▶  $\sum f'_n$  converges uniformly over  $I$ .

Then  $\sum_{n=0}^{+\infty} f_n$  is differentiable over  $I$  and:

$$\left( \sum_{n=0}^{+\infty} f_n(x) \right)' = \sum_{n=0}^{+\infty} f'_n(x)$$

# Summary

Pointwise and uniform convergences ( $\sum$ )

Absolute convergence

Normal convergence

Reminders about power series

Applications: Weierstrass approximation theorems

## Absolute convergence

The absolute convergence for series of functions extends the notion of absolute convergence for numerical series. Thus, like the pointwise convergence, it is a convergence that looks at each point separately, and that will not keep the interesting properties of the functions.

# Absolute convergence

The absolute convergence for series of functions extends the notion of absolute convergence for numerical series. Thus, like the pointwise convergence, it is a convergence that looks at each point separately, and that will not keep the interesting properties of the functions.

**Definition** [Absolute convergence]

We say that the series  $\sum f_n$  converges **absolutely** if the series  $\sum |f_n|$  converges pointwise.

# Absolute convergence

The absolute convergence for series of functions extends the notion of absolute convergence for numerical series. Thus, like the pointwise convergence, it is a convergence that looks at each point separately, and that will not keep the interesting properties of the functions.

**Definition** [Absolute convergence]

We say that the series  $\sum f_n$  converges **absolutely** if the series  $\sum |f_n|$  converges pointwise.

If we work with positives functions, obviously the pointwise and absolute convergence are the same thing.



# Absolute convergence

The absolute convergence for series of functions extends the notion of absolute convergence for numerical series. Thus, like the pointwise convergence, it is a convergence that looks at each point separately, and that will not keep the interesting properties of the functions.

**Definition** [Absolute convergence]

We say that the series  $\sum f_n$  converges **absolutely** if the series  $\sum |f_n|$  converges pointwise.

If we work with positives functions, obviously the pointwise and absolute convergence are the same thing.

As for numerical series:

**Proposition**

The absolute convergence implies the pointwise convergence.

## Absolute and uniform convergences

The absolute convergence will be of little interest to us since it has no link with the uniform convergence.

## Absolute and uniform convergences

The absolute convergence will be of little interest to us since it has no link with the uniform convergence.

Indeed, there exist series for which the convergence is uniform yet not absolute:

*example : the series  $\sum g_n$  where the  $g_n$  functions are constant, equal to  $\frac{(-1)^n}{n}$  converges uniformly (Leibniz's test for alternating series with*

*$\|R_n\|_\infty \leq \|g_{n+1}\|_\infty = \frac{1}{n+1} \xrightarrow{n \rightarrow +\infty} 0$ ), yet not absolutely since  $\sum |g_n| = \sum \frac{1}{n}$  is a divergent Riemann series;*

# Absolute and uniform convergences

The absolute convergence will be of little interest to us since it has no link with the uniform convergence.

Indeed, there exist series for which the convergence is uniform yet not absolute:

*example : the series  $\sum g_n$  where the  $g_n$  functions are constant, equal to  $\frac{(-1)^n}{n}$  converges uniformly (Leibniz's test for alternating series with*

*$\|R_n\|_\infty \leq \|g_{n+1}\|_\infty = \frac{1}{n+1} \xrightarrow{n \rightarrow +\infty} 0$ ), yet not absolutely since  $\sum |g_n| = \sum \frac{1}{n}$  is a divergent Riemann series;*

and others for which the convergence is absolute yet not uniform (for instance, any series of positive functions that converges pointwise yet not uniformly).

# Summary

Pointwise and uniform convergences ( $\sum$ )

Absolute convergence

Normal convergence

Reminders about power series

Applications: Weierstrass approximation theorems

## Normal convergence

The normal convergence of a series of functions is the strongest of all the convergences we will study, which means that it implies all the others.

# Normal convergence

The normal convergence of a series of functions is the strongest of all the convergences we will study, which means that it implies all the others.

**Definition** [Normal convergence of a series of functions]

Let  $(f_n)$  be a sequence of bounded<sup>a</sup> functions over  $I$ .

We say that the series  $\sum f_n$  converges **normally** over  $I$  if the numerical series  $\sum \|f_n\|_\infty$  is convergent.

---

<sup>a</sup>at least after a certain rank.

# Normal convergence

The normal convergence of a series of functions is the strongest of all the convergences we will study, which means that it implies all the others.

**Definition** [Normal convergence of a series of functions]

Let  $(f_n)$  be a sequence of bounded<sup>a</sup> functions over  $I$ .

We say that the series  $\sum f_n$  converges **normally** over  $I$  if the numerical series  $\sum \|f_n\|_\infty$  is convergent.

---

<sup>a</sup>at least after a certain rank.

This characterisation is thus simpler, yet way less accurate.



# Properties of the normal convergence

## **Proposition**

The normal convergence implies the pointwise and absolute convergences.

## Properties of the normal convergence

### **Proposition**

The normal convergence implies the pointwise and absolute convergences.

One just needs to remark that by definition,  $|f_n(x)| \leq \|f_n\|_\infty$ .

## Properties of the normal convergence

### **Proposition**

The normal convergence implies the pointwise and absolute convergences.

One just needs to remark that by definition,  $|f_n(x)| \leq \|f_n\|_\infty$ .

### **Proposition**

The normal convergence implies the uniform convergence.

# Properties of the normal convergence

## **Proposition**

The normal convergence implies the pointwise and absolute convergences.


One just needs to remark that by definition,  $|f_n(x)| \leq \|f_n\|_\infty$ .

## **Proposition**

The normal convergence implies the uniform convergence.

The study of the normal convergence is a quick way, when it gives a favourable outcome, to determine that the convergence of a series of functions is uniform. However, this is not a necessary condition: when the convergence is not normal, we have to study the series more in detail.

# Links between the different convergences



# Summary

Pointwise and uniform convergences ( $\sum$ )

Absolute convergence

Normal convergence

Reminders about power series

Applications: Weierstrass approximation theorems

A power series  $\sum a_n z^n$  admits a radius of convergence  $R \in \mathbb{R}_+ \cup \{+\infty\}$ : it is the supremum of the moduli of complex numbers for which the series converges pointwise (or is bounded).

A power series  $\sum a_n z^n$  admits a radius of convergence  $R \in \mathbb{R}_+ \cup \{+\infty\}$ : it is the supremum of the moduli of complex numbers for which the series converges pointwise (or is bounded).

Then, the series is absolutely convergent over the open ball of radius  $R$  centred at 0 (over  $\mathbb{R}$ , it is the interval  $] -R; R[$ ) and on any segment included in this interval, the convergence is normal thus uniform ( $[r_1, r_2]$  with  $-R < r_1 < r_2 < R$ ).



A power series  $\sum a_n z^n$  admits a radius of convergence  $R \in \mathbb{R}_+ \cup \{+\infty\}$ : it is the supremum of the moduli of complex numbers for which the series converges pointwise (or is bounded).

Then, the series is absolutely convergent over the open ball of radius  $R$  centred at 0 (over  $\mathbb{R}$ , it is the interval  $] -R; R[$ ) and on any segment included in this interval, the convergence is normal thus uniform ( $[r_1, r_2]$  with  $-R < r_1 < r_2 < R$ ). If the series is still convergent at  $R$  (respectively  $-R$ ), the convergence is even uniform on every segment  $[r, R]$  (respectively  $[-R, r]$ ) with  $r \in ] -R, R[$ .

A power series  $\sum a_n z^n$  admits a radius of convergence  $R \in \mathbb{R}_+ \cup \{+\infty\}$ : it is the supremum of the moduli of complex numbers for which the series converges pointwise (or is bounded).

Then, the series is absolutely convergent over the open ball of radius  $R$  centred at 0 (over  $\mathbb{R}$ , it is the interval  $] -R; R[$ ) and on any segment included in this interval, the convergence is normal thus uniform ( $[r_1, r_2]$  with  $-R < r_1 < r_2 < R$ ). If the series is still convergent at  $R$  (respectively  $-R$ ), the convergence is even uniform on every segment  $[r, R]$  (respectively  $[-R, r]$ ) with  $r \in ] -R, R[$ .

Thus, the convergence being uniform, power series are an excellent way to approximate functions while keeping their interesting properties.

# Summary

Pointwise and uniform convergences ( $\sum$ )

Absolute convergence

Normal convergence

Reminders about power series

Applications: Weierstrass approximation theorems

## Weierstrass approximation theorem (polynomial)

**Theorem** [Weierstrass approximation theorem (polynomial)]

Every continuous function over a segment  $[a, b]$  is the limit of a sequence of polynomial functions that converges uniformly.

# Weierstrass approximation theorem (polynomial)

**Theorem** [Weierstrass approximation theorem (polynomial)]

Every continuous function over a segment  $[a, b]$  is the limit of a sequence of polynomial functions that converges uniformly.

The Russian mathematician Sergei Bernstein gave a constructive proof of this theorem: if  $f$  is continuous over  $[0, 1]$ , it is the uniform limit of the sequence of the polynomials

$$P_n : x \longmapsto \sum_{k=0}^n f\left(\frac{k}{n}\right) \binom{n}{k} x^k (1-x)^{n-k}.$$

# Weierstrass approximation theorem (polynomial)

**Theorem** [Weierstrass approximation theorem (polynomial)]

Every continuous function over a segment  $[a, b]$  is the limit of a sequence of polynomial functions that converges uniformly.

The Russian mathematician Sergei Bernstein gave a constructive proof of this theorem: if  $f$  is continuous over  $[0, 1]$ , it is the uniform limit of the sequence of the polynomials

$$P_n : x \longmapsto \sum_{k=0}^n f\left(\frac{k}{n}\right) \binom{n}{k} x^k (1-x)^{n-k}.$$

The right part of the expression is inspired from the binomial law of probabilities. The proof uses many results from the theory of probabilities, among which the weak law of large numbers; it enables to prove the pointwise and uniform convergences.

## Weierstrass approximation theorem (trigonometric)

**Theorem** [Weierstrass approximation theorem (trigonometric)]

Every continuous periodic function is the limit of a sequence of trigonometric polynomials that converges uniformly.

## Weierstrass approximation theorem (trigonometric)

### **Theorem** [Weierstrass approximation theorem (trigonometric)]

Every continuous periodic function is the limit of a sequence of trigonometric polynomials that converges uniformly.

We call trigonometric polynomial any function with complex values under the form

$P : x \mapsto \sum_{k=-n}^n c_k e^{ikx}$ . Such a function, when taking only real values, can also be

written  $P : x \mapsto a_0 + \sum_{k=1}^n (a_k \cos(kx) + b_k \sin(kx))$ .



## Weierstrass approximation theorem (trigonometric)

### **Theorem** [Weierstrass approximation theorem (trigonometric)]

Every continuous periodic function is the limit of a sequence of trigonometric polynomials that converges uniformly.

We call trigonometric polynomial any function with complex values under the form

$P : x \mapsto \sum_{k=-n}^n c_k e^{ikx}$ . Such a function, when taking only real values, can also be

written  $P : x \mapsto a_0 + \sum_{k=1}^n (a_k \cos(kx) + b_k \sin(kx))$ .

This result derives from the theory of Fourier series; the Hungarian mathematician Lipót Fejér proved that for every  $2\pi$ -periodic continuous function  $f$ , the sequence of

Cesàro means  $\frac{1}{p+1} \sum_{n=0}^p S_n(f)$  of the sequence of partial sums of its Fourier series

$S_n(f) : x \mapsto \sum_{k=-n}^n c_k(f) e^{ikx}$  with  $c_k(f) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(t) e^{-ikt} dt$  converges uniformly

towards  $f$ .