

Sequences of functions

Pointwise and uniform convergence

G. Goron

EPITA

October 18, 2021

Interest of the convergence

We take a look here at the notion of convergence of a sequence of functions $(f_n) \in (\mathbb{R}^I)^{\mathbb{N}}$, where I is an interval of \mathbb{R} .

Interest of the convergence

We take a look here at the notion of convergence of a sequence of functions

$(f_n) \in (\mathbb{R}^I)^{\mathbb{N}}$, where I is an interval of \mathbb{R} .

When we approximate a function by a sequence of functions, it is necessary to know if this approximation is correct and if it is a good one. That is what we are going to discuss here.

Interest of the convergence

We take a look here at the notion of convergence of a sequence of functions

$(f_n) \in (\mathbb{R}^I)^{\mathbb{N}}$, where I is an interval of \mathbb{R} .

When we approximate a function by a sequence of functions, it is necessary to know if this approximation is correct and if it is a good one. That is what we are going to discuss here.

The notion of convergence as we can guess it intuitively unfortunately does not allow to keep a lot of interesting properties of functions. Thus, we will try to refine it a little.

Interest of the convergence

We take a look here at the notion of convergence of a sequence of functions

$(f_n) \in (\mathbb{R}^I)^{\mathbb{N}}$, where I is an interval of \mathbb{R} .

When we approximate a function by a sequence of functions, it is necessary to know if this approximation is correct and if it is a good one. That is what we are going to discuss here.

The notion of convergence as we can guess it intuitively unfortunately does not allow to keep a lot of interesting properties of functions. Thus, we will try to refine it a little. The existence of several ways to define a distance over a space of functions can also offer several ways to look at the problem, it is good to know how to determine which one is the most interesting.

Pointwise convergence

The pointwise convergence is the most intuitive notion of convergence, because it extends to functions the notion of convergence for numerical sequences.

Pointwise convergence

The pointwise convergence is the most intuitive notion of convergence, because it extends to functions the notion of convergence for numerical sequences. We consider the interval I where the function is defined as a collection of distinct points, and we examine **each of them separately**. Thus, for every $x \in I$, we determine towards which value the sequence $(f_n(x))$ converges (if it does converge). It is a way to consider the convergence of a sequence of functions as a gathering of convergences of numerical sequences.

Pointwise convergence

The pointwise convergence is the most intuitive notion of convergence, because it extends to functions the notion of convergence for numerical sequences. We consider the interval I where the function is defined as a collection of distinct points, and we examine **each of them separately**. Thus, for every $x \in I$, we determine towards which value the sequence $(f_n(x))$ converges (if it does converge). It is a way to consider the convergence of a sequence of functions as a gathering of convergences of numerical sequences.

However, dealing with each point separately lets us lose the global properties of the function: we do not use the topology of I which links the values the function takes at its different arguments.

Pointwise convergence

Definition [Pointwise convergence of a sequence of functions]

We say that (f_n) *converges pointwise* towards the function $f \in \mathbb{R}^I$ iff:

$$\forall x \in I, \quad f_n(x) \longrightarrow f(x).$$

Pointwise convergence

Definition [Pointwise convergence of a sequence of functions]

We say that (f_n) *converges pointwise* towards the function $f \in \mathbb{R}^I$ iff:

$$\forall x \in I, \quad f_n(x) \longrightarrow f(x).$$

Which can be written as:

$$\forall x \in I, \forall \varepsilon \in \mathbb{R}_+^*, \exists N_\varepsilon \in \mathbb{N}, \forall n \in \mathbb{N}, \quad n \geq N_\varepsilon \implies |f_n(x) - f(x)| < \varepsilon.$$

Here is a simple example: let us look at the sequence of functions $f_n : x \mapsto x^n$ over the segment $[0, 1]$.

Here is a simple example: let us look at the sequence of functions $f_n : x \mapsto x^n$ over the segment $[0, 1]$.

► for every $n > 0$, $f_n(0) = 0$ thus $f_n(0) \xrightarrow{n \rightarrow +\infty} 0$;

Here is a simple example: let us look at the sequence of functions $f_n : x \mapsto x^n$ over the segment $[0, 1]$.

- ▶ for every $n > 0$, $f_n(0) = 0$ thus $f_n(0) \xrightarrow{n \rightarrow +\infty} 0$;
- ▶ for every $n \geqslant 0$, $f_n(1) = 1$ thus $f_n(1) \xrightarrow{n \rightarrow +\infty} 1$;

Here is a simple example: let us look at the sequence of functions $f_n : x \mapsto x^n$ over the segment $[0, 1]$.

- ▶ for every $n > 0$, $f_n(0) = 0$ thus $f_n(0) \xrightarrow{n \rightarrow +\infty} 0$;
- ▶ for every $n \geq 0$, $f_n(1) = 1$ thus $f_n(1) \xrightarrow{n \rightarrow +\infty} 1$;
- ▶ for every $x \in]0; 1[$, $x^n \xrightarrow{n \rightarrow +\infty} 0$.

Here is a simple example: let us look at the sequence of functions $f_n : x \mapsto x^n$ over the segment $[0, 1]$.


- ▶ for every $n > 0$, $f_n(0) = 0$ thus $f_n(0) \xrightarrow{n \rightarrow +\infty} 0$;
- ▶ for every $n \geq 0$, $f_n(1) = 1$ thus $f_n(1) \xrightarrow{n \rightarrow +\infty} 1$;
- ▶ for every $x \in]0; 1[$, $x^n \xrightarrow{n \rightarrow +\infty} 0$.

The sequence of function thus converges pointwise towards the function


$$f : x \mapsto \begin{cases} 0 & \text{if } 0 \leq x < 1; \\ 1 & \text{if } x = 1. \end{cases}$$

We can graphically represent the evolution of the f_n functions towards their limit:

We can graphically represent the evolution of the f_n functions towards their limit:




We can graphically represent the evolution of the f_n functions towards their limit:



Slight disappointment: all the f_n functions are smooth, yet the limit function f isn't even continuous.


We can graphically represent the evolution of the f_n functions towards their limit:



Slight disappointment: all the f_n functions are smooth, yet the limit function f isn't even continuous.

The pointwise convergence does not guarantee the conservation of smoothness properties.

We can graphically represent the evolution of the f_n functions towards their limit:



Slight disappointment: all the f_n functions are smooth, yet the limit function f isn't even continuous.

The pointwise convergence does not guarantee the conservation of smoothness properties.

And that is far from being the only problem ...

Pointwise convergence

Among the interesting properties that the pointwise convergence does not keep at the limit, we have:

Pointwise convergence

Among the interesting properties that the pointwise convergence does not keep at the limit, we have:

- ▶ continuity: even if (f_n) converges pointwise towards f , f is not always continuous;

Pointwise convergence

Among the interesting properties that the pointwise convergence does not keep at the limit, we have:

- ▶ continuity: even if (f_n) converges pointwise towards f , f is not always continuous;
- ▶ inversion between integral and limits: if (f_n) converges pointwise towards f , we do not necessarily get

$$\lim_{n \rightarrow +\infty} \int_I f_n(t) dt \stackrel{?}{=} \int_I f(t) dt \left(= \int_I \lim_{n \rightarrow +\infty} f_n(t) dt \right)$$

Pointwise convergence

Among the interesting properties that the pointwise convergence does not keep at the limit, we have:

- ▶ continuity: even if (f_n) converges pointwise towards f , f is not always continuous;
- ▶ inversion between integral and limits: if (f_n) converges pointwise towards f , we do not necessarily get

$$\lim_{n \rightarrow +\infty} \int_I f_n(t) dt \stackrel{?}{=} \int_I f(t) dt \left(= \int_I \lim_{n \rightarrow +\infty} f_n(t) dt \right)$$

- ▶ inversion between differentiation and limits: if (f_n) converges pointwise towards f , we do not necessarily get

$$\lim_{n \rightarrow +\infty} f'_n(t) \stackrel{?}{=} f'(t) = \left(\lim_{n \rightarrow +\infty} f_n \right)'$$

So when these properties do matter in the problem we are looking at (if for instance we want to calculate an integral by successive approximations), the pointwise convergence is too inaccurate a tool to be used safely.

So when these properties do matter in the problem we are looking at (if for instance we want to calculate an integral by successive approximations), the pointwise convergence is too inaccurate a tool to be used safely.

Thus, we will need to devise more powerful ways to evaluate the convergence of sequences of functions, particularly in regards to the properties at the core of our problems.

Uniform convergence

The uniform convergence is a stronger notion than the pointwise convergence (meaning that the first implies the latter). It will allow to check that some interesting properties are kept through our approximations.

Uniform convergence

The uniform convergence is a stronger notion than the pointwise convergence (meaning that the first implies the latter). It will allow to check that some interesting properties are kept through our approximations.

Definition [Uniform convergence of a sequence of functions] We say that a sequence (f_n) *converges uniformly* towards a function f over I iff, as long as the following quantities are defined:

$$\sup_{x \in I} |f_n(x) - f(x)| \xrightarrow{n \rightarrow +\infty} 0.$$

that is to say

$$\|f_n - f\|_{\infty} \xrightarrow{n \rightarrow +\infty} 0.$$

Which can be written as

$$\forall \varepsilon \in \mathbb{R}_+^*, \exists N_{\varepsilon} \in \mathbb{N}, \forall n \in \mathbb{N}, (n \geq N_{\varepsilon} \implies \forall x \in I, |f_n(x) - f(x)| < \varepsilon).$$

Uniform convergence

The uniform convergence is a stronger notion than the pointwise convergence (meaning that the first implies the latter). It will allow to check that some interesting properties are kept through our approximations.

Definition [Uniform convergence of a sequence of functions] We say that a sequence (f_n) *converges uniformly* towards a function f over I iff, as long as the following quantities are defined:

$$\sup_{x \in I} |f_n(x) - f(x)| \xrightarrow{n \rightarrow +\infty} 0.$$

that is to say

$$\|f_n - f\|_{\infty} \xrightarrow{n \rightarrow +\infty} 0.$$

Which can be written as

$$\forall \varepsilon \in \mathbb{R}_+^*, \exists N_{\varepsilon} \in \mathbb{N}, \forall n \in \mathbb{N}, (n \geq N_{\varepsilon} \implies \forall x \in I, |f_n(x) - f(x)| < \varepsilon).$$

Remark: the uniform convergence is the convergence with respect to the infinite norm (or distance).

Uniform convergence: in practice

In order to determine if a convergence is uniform, we can use several methods:

Uniform convergence: in practice

In order to determine if a convergence is uniform, we can use several methods:

- ▶ We can use simple upper or lower bounds, or any kind of approximation using analytic methods;

Uniform convergence: in practice

In order to determine if a convergence is uniform, we can use several methods:

- ▶ We can use simple upper or lower bounds, or any kind of approximation using analytic methods;
- ▶ Some properties are useful to work with (smoothness, monotonicity etc.)

Uniform convergence: in practice

In order to determine if a convergence is uniform, we can use several methods:

- ▶ We can use simple upper or lower bounds, or any kind of approximation using analytic methods;
- ▶ Some properties are useful to work with (smoothness, monotonicity etc.)
- ▶ If the functions are differentiable, we can look for their maxima / suprema using a study of their variations.

Uniform convergence: in practice

In order to determine if a convergence is uniform, we can use several methods:

- ▶ We can use simple upper or lower bounds, or any kind of approximation using analytic methods;
- ▶ Some properties are useful to work with (smoothness, monotonicity etc.)
- ▶ If the functions are differentiable, we can look for their maxima / suprema using a study of their variations.

List not exhaustive. One can also imagine using sequential criteria, particularly when one wants to come up with a simple counter-example.

Properties of the uniform convergence

Uniform \implies Pointwise

Theorem

Uniform convergence implies pointwise convergence (towards the same limit).

Properties of the uniform convergence

Uniform \implies Pointwise

Theorem

Uniform convergence implies pointwise convergence (towards the same limit).

In other words, if (f_n) converges uniformly towards f over I then it converges pointwise towards f .

Obviously, the reciprocal is false (use the previous example of $f_n : x \mapsto x^n$ over $[0, 1]$, where $\|f_n - f\|_\infty = 1$ for every n).

The pointwise convergence remains insufficient, yet it is still a necessary step to find the limit towards which we want to determine whether the convergence is uniform or not.

Properties of the uniform convergence

Continuity

Theorem

If a sequence of continuous functions converges uniformly, then its limit is a continuous function.

Properties of the uniform convergence

Continuity

Theorem

If a sequence of continuous functions converges uniformly, then its limit is a continuous function.

The uniform convergence thus preserves continuity.

Properties of the uniform convergence

Continuity

Theorem

If a sequence of continuous functions converges uniformly, then its limit is a continuous function.

The uniform convergence thus preserves continuity.

Remark: the contraposition of this property can be used to show that a convergence is not uniform.

Properties of the uniform convergence

Swapping integrals and limits

Theorem

In this case, I is a **segment** $[a, b]$.

If (f_n) converges uniformly towards f over $[a, b]$, then:

$$\lim_{n \rightarrow +\infty} \int_a^b f_n(t) dt = \int_a^b \lim_{n \rightarrow +\infty} f_n(t) dt = \int_a^b f(t) dt.$$

Properties of the uniform convergence

Swapping integrals and limits

Theorem

In this case, I is a segment $[a, b]$.

If (f_n) converges uniformly towards f over $[a, b]$, then:

$$\lim_{n \rightarrow +\infty} \int_a^b f_n(t) dt = \int_a^b \lim_{n \rightarrow +\infty} f_n(t) dt = \int_a^b f(t) dt.$$

Thus, if the convergence is uniform, we can «swap» the signs of integration and limit.

Properties of the uniform convergence

Swapping integrals and limits

Theorem

In this case, I is a **segment** $[a, b]$.

If (f_n) converges uniformly towards f over $[a, b]$, then:

$$\lim_{n \rightarrow +\infty} \int_a^b f_n(t) dt = \int_a^b \lim_{n \rightarrow +\infty} f_n(t) dt = \int_a^b f(t) dt.$$

Thus, if the convergence is uniform, we can «swap» the signs of integration and limit.

Remark: this result becomes false if I is not bounded. Particularly, let us consider the sequence (g_n) with $\forall n \in \mathbb{N}^*, \forall x \in \mathbb{R}, g_n(x) = \begin{cases} \frac{1}{n} & \text{if } x \in [-n; n] \\ 0 & \text{else} \end{cases}$ then (g_n) converges

uniformly towards the zero function over \mathbb{R} , yet for every n , $\int_{\mathbb{R}} g_n(t) dt = 2 \not\xrightarrow[n \rightarrow +\infty]{} 0$.

Properties of the uniform convergence

Swapping derivatives and limits

Beware: here the hypotheses are way more restrictive.

Theorem

Let us suppose that:

- ▶ the f_n functions are of class C^1 over I ;
- ▶ (f_n) converges (pointwise) over I towards f ;
- ▶ (f'_n) converges uniformly over I towards a function g .

Then f is differentiable over I and $f' = g$, that is to say:

$$\frac{d}{dx} \left(\lim_{n \rightarrow +\infty} f_n(x) \right) = \lim_{n \rightarrow +\infty} \frac{d}{dx} (f_n(x))$$

Properties of the uniform convergence

Swapping derivatives and limits

Beware: here the hypotheses are way more restrictive.

Theorem

Let us suppose that:

- ▶ the f_n functions are of class C^1 over I ;
- ▶ (f_n) converges (pointwise) over I towards f ;
- ▶ (f'_n) converges uniformly over I towards a function g .

Then f is differentiable over I and $f' = g$, that is to say:

$$\frac{d}{dx} \left(\lim_{n \rightarrow +\infty} f_n(x) \right) = \lim_{n \rightarrow +\infty} \frac{d}{dx} (f_n(x))$$

So, here again we can «swap» the signs of limit and differentiation.

Remark: for all the properties we talked about here, the uniform convergence is a sufficient yet not necessary condition. There exists indeed sequences of continuous functions that converge pointwise towards continuous functions, and so on. The uniform convergence is just a relatively simple way to make sure that these important results can be applied.