

Logical Formalism

Exercise Sheet 1

Adrien Pommellet, EPITA

August 14, 2024

Main Exercises

The Dumbest Proofs

Find the errors in the following proofs. Note that the result may be true, even if the method isn't.

Exercise 1 - *Proving that $\sin(x + \pi) = -\sin(x)$.*

$$\sin(x + \pi) = \sin(x) + \sin(\pi) \quad (1)$$

$$= \sin(x) - 1 \quad (2)$$

$$= (-1)\sin(x) \quad (3)$$

$$= -\sin(x) \quad (4)$$

Exercise 2 - *Proving that $2 = 4$.*

$$2 = 4 \quad (5)$$

$$\implies 2 - 3 = 4 - 3 \quad (6)$$

$$\implies -1 = 1 \quad (7)$$

$$\implies (-1)^2 = 1^2 \quad (8)$$

$$\implies 1 = 1 \quad (9)$$

Indeed, $1 = 1$. Hence, $2 = 4$.

Exercise 3 - *The exponential function has an upper bound.* Given a real number x , since the exponential function is strictly increasing, $e^x < e^{x+1}$. Consider $M = e^{x+1}$. Hence for all real numbers x , $e^x < M$. The exponential function therefore has an upper bound.

Sets and Formulas

Exercise 4 - *A strange equation.* Let E be a set and $A, B \in \mathcal{P}(E)$ be two subsets of E . Prove that $\forall X \in \mathcal{P}(E), (A \cap X = B) \iff (B \subseteq X) \wedge (X \subseteq B \cup A^c) \wedge (B \subseteq A)$.

Exercise 5 - \forall and \vee . Consider $A = \forall x \in E, P(x) \vee Q(x)$ and $B = (\forall x \in E, P(x)) \vee (\forall x \in E, Q(x))$. Does A imply B , or B imply A ? Find a counter-example if one of these implications does not hold.

Exercise 6 - \exists and \wedge . Consider $A = \exists x \in E, P(x) \wedge Q(x)$ and $B = (\exists x \in E, P(x)) \wedge (\exists x \in E, Q(x))$. Does A imply B , or B imply A ? Find a counter-example if one of these implications does not hold.

Extra Exercises

Classical Logics

Exercise 7 - *Negating \vee .* Let ABC be a triangle. What is the negation of the proposition *ABC is isosceles*?

Exercise 8 - *An equivalence.* Prove that $\forall n \in \mathbb{N}$, n is a multiple of 3 if and only if the sum of its digits is a multiple of 3 as well.

Set Theory

Exercise 9 - *Properties of \emptyset .* Prove that for any set A , $A \cap \emptyset = \emptyset$ and $A \cup \emptyset = A$.

Exercise 10 - *A triple equivalence.* Prove that the three following propositions are equivalent, given two sets A and B :

1. $A \subseteq B$.
2. $B^c \subseteq A^c$.
3. $B^c \cap A = \emptyset$.

Exercise 11 - *Properties of the complement.* Prove that for any set $A \in \mathcal{P}(E)$, $A \cap A^c = \emptyset$ and $A \cup A^c = E$.

Quantifiers

Exercise 12 - *Negating logical formulas.* Write the negation of the following formulas (using as simple an equivalent formula as possible):

1. $A \implies (B \wedge \neg(C \vee D))$.

2. $A \vee B \iff C$.

3. $\forall x \in X, \forall y \in Y, (P(x) \wedge (\exists z \in Z, Q(y, z)))$.