

Metric spaces

Distances and norms

G. Goron

EPITA

October 16, 2021

We choose to work on a vector space E over \mathbb{R} or \mathbb{C} .

We choose to work on a vector space E over \mathbb{R} or \mathbb{C} .

Definition [Norm]

Let E be a vector space over a field \mathbb{K} — here \mathbb{R} or \mathbb{C} .

An application $N : E \longrightarrow \mathbb{R}_+$ is called a *norm* over E if it checks the following

$$\begin{array}{lll} \text{properties:} & \left\| \begin{array}{ll} \forall x \in E, & \mathbf{N}(x) = 0 \iff x = 0 & \textit{it is «definite»} \\ \forall x \in E, \forall \lambda \in \mathbb{K}, & \mathbf{N}(\lambda x) = |\lambda| \mathbf{N}(x) & \textit{(homogeneity)} \\ \forall (x, y) \in E^2, & \mathbf{N}(x + y) \leq \mathbf{N}(x) + \mathbf{N}(y) & \textit{(triangle inequality)} \end{array} \right. \end{array}$$

We choose to work on a vector space E over \mathbb{R} or \mathbb{C} .

Definition [Norm]

Let E be a vector space over a field \mathbb{K} — here \mathbb{R} or \mathbb{C} .

An application $N : E \longrightarrow \mathbb{R}_+$ is called a *norm* over E if it checks the following

properties:		$\forall x \in E,$	$\mathbf{N}(x) = 0 \iff x = 0$	<i>it is «definite»</i>
		$\forall x \in E, \forall \lambda \in \mathbb{K},$	$\mathbf{N}(\lambda x) = \lambda \mathbf{N}(x)$	<i>(homogeneity)</i>
		$\forall (x, y) \in E^2,$	$\mathbf{N}(x + y) \leq \mathbf{N}(x) + \mathbf{N}(y)$	<i>(triangle inequality)</i>

The homogeneity property goes together well with the structure of vector space.

A norm N over E induces a distance d_N over E through the relation:

$$\forall (x, y) \in E^2, \quad d_N(x, y) = N(x - y).$$

A norm N over E induces a distance d_N over E through the relation:

$$\forall (x, y) \in E^2, \quad d_N(x, y) = N(x - y).$$

Hence, the norm of a vector can be viewed as its distance to 0_E :

$$N(x) = d_N(x, 0_E).$$

A norm N over E induces a distance d_N over E through the relation:

$$\forall (x, y) \in E^2, \quad d_N(x, y) = N(x - y).$$

Hence, the norm of a vector can be viewed as its distance to 0_E :

$$N(x) = d_N(x, 0_E).$$

Remark: there are also distances over E that do not come from any norm.

A bit of topology:

Definition [Open / Closed ball]

We call *open ball* centred at a of radius r with respect to d the set

$$B_d(a, r) = \{x \in E \mid d(a, x) < r\}.$$

You get a *closed ball* if you substitute a large inequality for the strict one.

A bit of topology:

Definition [Open / Closed ball]

We call *open ball* centred at a of radius r with respect to d the set

$$B_d(a, r) = \{x \in E \mid d(a, x) < r\}.$$

You get a *closed ball* if you substitute a large inequality for the strict one.

The concept of ball extends the notion of neighbourhood you already used in \mathbb{R} to multidimensional spaces.

Definition [p -norms over \mathbb{R}^n]

Let $p \in [1, +\infty[$. We call p -norm over \mathbb{R}^n the norm denoted by $\|\cdot\|_p$ defined for every $x = (x_1, \dots, x_n) \in \mathbb{R}^n$ by

$$\|x\|_p = \left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}}.$$

This definition is extended to $p = +\infty$ by looking at the limit:

$$\|x\|_\infty = \sup_{1 \leq i \leq n} |x_i| = \max_{1 \leq i \leq n} |x_i|$$

We denote by d_p the distance induced by $\|\cdot\|_p$.

Definition [p -norms over \mathbb{R}^n]

Let $p \in [1, +\infty[$. We call p -norm over \mathbb{R}^n the norm denoted by $\|\cdot\|_p$ defined for every $x = (x_1, \dots, x_n) \in \mathbb{R}^n$ by

$$\|x\|_p = \left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}}.$$

This definition is extended to $p = +\infty$ by looking at the limit:


$$\|x\|_\infty = \sup_{1 \leq i \leq n} |x_i| = \max_{1 \leq i \leq n} |x_i|$$

We denote by d_p the distance induced by $\|\cdot\|_p$.

d_1 is called Manhattan distance (or taxicab distance), while d_2 is the Euclidean distance.

Here is a graph showing the three distances d_1, d_2, d_∞ between two points of \mathbb{R}^2 :

Here is a graph showing the three distances d_1, d_2, d_∞ between two points of \mathbb{R}^2 :



Intuitively, one can remark the inequality:

$$\|x - y\|_{\infty} \leq \|x - y\|_2 \leq \|x - y\|_1$$

Intuitively, one can remark the inequality:

$$\|x - y\|_{\infty} \leq \|x - y\|_2 \leq \|x - y\|_1$$

And exactly as it seems, for every $x \in \mathbb{R}^n$ we get:

$$\|x\|_{\infty} \leq \|x\|_2 \leq \|x\|_1 \leq n\|x\|_{\infty}$$

Intuitively, one can remark the inequality:

$$\|x - y\|_{\infty} \leq \|x - y\|_2 \leq \|x - y\|_1$$

And exactly as it seems, for every $x \in \mathbb{R}^n$ we get:

$$\|x\|_{\infty} \leq \|x\|_2 \leq \|x\|_1 \leq n\|x\|_{\infty}$$

And more generally:

$$p < q \implies \|x\|_p \geq \|x\|_q.$$

Definition [Equivalent norms]

Two norms N and N' over E are said to be equivalent if and only if there exist two strictly positive real numbers k_1 and k_2 such that:

$$\forall x \in E, \quad k_1 N(x) \leq N'(x) \leq k_2 N(x).$$

Definition [Equivalent norms]

Two norms N and N' over E are said to be equivalent if and only if there exist two strictly positive real numbers k_1 and k_2 such that:

$$\forall x \in E, \quad k_1 N(x) \leq N'(x) \leq k_2 N(x).$$

Please remark that k_1 and k_2 do not depend on the chosen x .

Definition [Equivalent norms]

Two norms N and N' over E are said to be equivalent if and only if there exist two strictly positive real numbers k_1 and k_2 such that:


$$\forall x \in E, \quad k_1 N(x) \leq N'(x) \leq k_2 N(x).$$

Please remark that k_1 and k_2 do not depend on the chosen x .


Theorem [Equivalence of norms in finite dimension]

If E is a vector space of finite dimension, then all norms over E are equivalent.

Balls of radius 1 for norms 1, 2, ∞ :




Balls of radius 1 for norms 1, 2, ∞ :




- ▶ If we choose another radius, we get the same graph up to the scale (homogeneity).

Balls of radius 1 for norms 1, 2, ∞ :



- ▶ If we choose another radius, we get the same graph up to the scale (homogeneity).
- ▶ The ball inclusions translate the inequalities between norms; these can be determined by comparing the radii in the optimal framing of the ball of a given norm by two balls of another one norm.

Balls of radius 1 for norms 1, 2, ∞ :



- ▶ If we choose another radius, we get the same graph up to the scale (homogeneity).
- ▶ The ball inclusions translate the inequalities between norms; these can be determined by comparing the radii in the optimal framing of the ball of a given norm by two balls of another one norm.
- ▶ This is also topologically interesting: all these norms depict the same structure.

Definition [p -norms over sets of functions]

Let I be a segment of \mathbb{R} not reduced to a single element, let us look at the space $\mathcal{C} = C^0(I, \mathbb{R})$ of continuous functions over I with values in \mathbb{R} .

Let $p \in [1, +\infty[$. We call p -norm over \mathcal{C} the norm denoted by $\|\cdot\|_p$ defined for every $f \in \mathcal{C}$ by

$$\|f\|_p = \left(\int_I |f(t)|^p dt \right)^{\frac{1}{p}}.$$

For $p = +\infty$, we set:

$$\|f\|_\infty = \sup_{x \in I} |f(x)|.$$

We denote by d_p the distance induced by $\|\cdot\|_p$.

It is possible to change the hypotheses on regularity to define distances and norms over different kinds of function spaces, yet this forces to diminish the strains given by the axioms.

It is possible to change the hypotheses on regularity to define distances and norms over different kinds of function spaces, yet this forces to diminish the strains given by the axioms.

Moreover, the theorem of equivalence of norms in finite dimension unfortunately does not hold anymore. We even get the result:

If $p \neq q$ then $\|\cdot\|_p$ and $\|\cdot\|_q$ are not equivalent over \mathcal{C} .