

# Fourier Series

## 1 - Inner product spaces

G. Goron

EPITA

November 3, 2021



In order to approach the topic of Fourier series, we need to talk about the adapted algebraic structure: the one we call inner product space.

In order to approach the topic of Fourier series, we need to talk about the adapted algebraic structure: the one we call inner product space.

The calculation of the coefficients of the Fourier series, the theorems of convergence, all the important results ... use properties that can be derived from our knowledge of that kind of structure.

In order to approach the topic of Fourier series, we need to talk about the adapted algebraic structure: the one we call inner product space.

The calculation of the coefficients of the Fourier series, the theorems of convergence, all the important results ... use properties that can be derived from our knowledge of that kind of structure.

It is an extension of the notion of vector space. On a vector space  $E$  over a field  $\mathbb{K}$ , is added an inner operation  $E \times E \rightarrow \mathbb{K}$  that, to two elements of  $E$ , maps a scalar from  $\mathbb{K}$ .

In order to approach the topic of Fourier series, we need to talk about the adapted algebraic structure: the one we call inner product space.

The calculation of the coefficients of the Fourier series, the theorems of convergence, all the important results ... use properties that can be derived from our knowledge of that kind of structure.

It is an extension of the notion of vector space. On a vector space  $E$  over a field  $\mathbb{K}$ , is added an inner operation  $E \times E \rightarrow \mathbb{K}$  that, to two elements of  $E$ , maps a scalar from  $\mathbb{K}$ . This operation is called **inner product**.



# Inner product

In order to be an inner product, an application  $\varphi : E \times E \rightarrow \mathbb{K}$  has to check several properties.



## Inner product

In order to be an inner product, an application  $\varphi : E \times E \rightarrow \mathbb{K}$  has to check several properties.

These will ensure that this operation is an interesting one to add to a structure of vector space, allowing us particularly to induce a norm.

## Inner product

In order to be an inner product, an application  $\varphi : E \times E \rightarrow \mathbb{K}$  has to check several properties.

These will ensure that this operation is an interesting one to add to a structure of vector space, allowing us particularly to induce a norm.

These properties vary slightly depending on the chosen field  $\mathbb{K} = \mathbb{R}$  or  $\mathbb{C}$ .

## Bilinearity (and sesquilinearity)

The first property facilitates the manipulations with respect to the vector space structure: it is the **bilinearity** / **sesquilinearity**.

## Bilinearity (and sesquilinearity)

The first property facilitates the manipulations with respect to the vector space structure: it is the **bilinearity** / **sesquilinearity**.

Reminder: a function  $f$  over a  $\mathbb{K}$ -vector space  $E$  is said to be **linear** if

$$\forall (x, y) \in E^2, \forall \lambda \in \mathbb{K}, \quad f(x + y) = f(x) + f(y) \quad \text{and} \quad f(\lambda x) = \lambda f(x)$$

## Bilinearity (and sesquilinearity)

The first property facilitates the manipulations with respect to the vector space structure: it is the **bilinearity / sesquilinearity**.

Reminder: a function  $f$  over a  $\mathbb{K}$ -vector space  $E$  is said to be **linear** if

$$\forall (x, y) \in E^2, \forall \lambda \in \mathbb{K}, \quad f(x + y) = f(x) + f(y) \quad \text{and} \quad f(\lambda x) = \lambda f(x)$$

If  $\mathbb{K} = \mathbb{C}$ , we say that  $g$  is **semi-linear** if

$$\forall (x, y) \in E^2, \forall \lambda \in \mathbb{K}, \quad g(x + y) = g(x) + g(y) \quad \text{and} \quad g(\lambda x) = \bar{\lambda} f(x)$$

## Bilinearity (and sesquilinearity)

We define the bilinearity / sesquilinearity thus:

$$\mathbb{K} = \mathbb{R}$$

$\varphi : E \times E \rightarrow \mathbb{K}$  is **bilinear** if:

# Bilinearity (and sesquilinearity)

We define the bilinearity / sesquilinearity thus:

$$\mathbb{K} = \mathbb{R}$$

$\varphi : E \times E \rightarrow \mathbb{K}$  is **bilinear** if:

- ▶ it is **linear with respect to the left variable** :

$\forall y \in E, \varphi_y : x \mapsto \varphi(x, y)$  is linear ;

# Bilinearity (and sesquilinearity)

We define the bilinearity / sesquilinearity thus:

$$\mathbb{K} = \mathbb{R}$$

$\varphi : E \times E \rightarrow \mathbb{K}$  is **bilinear** if:

- ▶ it is linear with respect to the left variable :

$$\forall y \in E, \varphi_y : x \mapsto \varphi(x, y) \text{ is linear ;}$$

- ▶ it is linear with respect to the left variable :

$$\forall x \in E, \varphi_x : y \mapsto \varphi(x, y) \text{ is linear.}$$



# Bilinearity (and sesquilinearity)

We define the bilinearity / sesquilinearity thus:

$$\mathbb{K} = \mathbb{R}$$

$$\mathbb{K} = \mathbb{C}$$

$\varphi : E \times E \rightarrow \mathbb{K}$  is **bilinear** if:

- ▶ it is **linear with respect to the left variable** :

$$\forall y \in E, \varphi_y : x \mapsto \varphi(x, y) \text{ is linear ;}$$

- ▶ it is **linear with respect to the left variable** :

$$\forall x \in E, \varphi_x : y \mapsto \varphi(x, y) \text{ is linear.}$$

$\varphi : E \times E \rightarrow \mathbb{K}$  est **left-side sesquilinear** if:

# Bilinearity (and sesquilinearity)

We define the bilinearity / sesquilinearity thus:

$$\mathbb{K} = \mathbb{R}$$

$\varphi : E \times E \rightarrow \mathbb{K}$  is **bilinear** if:

- ▶ it is **linear with respect to the left variable** :  
 $\forall y \in E, \varphi_y : x \mapsto \varphi(x, y)$  is linear ;
- ▶ it is **linear with respect to the left variable** :  
 $\forall x \in E, \varphi_x : y \mapsto \varphi(x, y)$  is linear.

$$\mathbb{K} = \mathbb{C}$$

$\varphi : E \times E \rightarrow \mathbb{K}$  est **left-side sesquilinear** if:

- ▶ it is **semi-linear with respect to the left variable** :  
 $\forall y \in E, \varphi_y : x \mapsto \varphi(x, y)$  is semi-linear ;

# Bilinearity (and sesquilinearity)

We define the bilinearity / sesquilinearity thus:

$$\mathbb{K} = \mathbb{R}$$

$\varphi : E \times E \rightarrow \mathbb{K}$  is **bilinear** if:

- ▶ it is **linear with respect to the left variable** :  
 $\forall y \in E, \varphi_y : x \mapsto \varphi(x, y)$  is linear ;
- ▶ it is **linear with respect to the right variable** :  
 $\forall x \in E, \varphi_x : y \mapsto \varphi(x, y)$  is linear.

$$\mathbb{K} = \mathbb{C}$$

$\varphi : E \times E \rightarrow \mathbb{K}$  est **left-side sesquilinear** if:

- ▶ it is **semi-linear with respect to the left variable** :  
 $\forall y \in E, \varphi_y : x \mapsto \varphi(x, y)$  is semi-linear ;
- ▶ it is **linear with respect to the right variable** :  
 $\forall x \in E, \varphi_x : y \mapsto \varphi(x, y)$  is linear.

# Bilinearity (and sesquilinearity)

We define the bilinearity / sesquilinearity thus:

$$\mathbb{K} = \mathbb{R}$$

$\varphi : E \times E \rightarrow \mathbb{K}$  is **bilinear** if:

- ▶ it is **linear with respect to the left variable** :  
 $\forall y \in E, \varphi_y : x \mapsto \varphi(x, y)$  is linear ;
- ▶ it is **linear with respect to the right variable** :  
 $\forall x \in E, \varphi_x : y \mapsto \varphi(x, y)$  is linear.

$$\mathbb{K} = \mathbb{C}$$

$\varphi : E \times E \rightarrow \mathbb{K}$  est **left-side sesquilinear** if:

- ▶ it is **semi-linear with respect to the left variable** :  
 $\forall y \in E, \varphi_y : x \mapsto \varphi(x, y)$  is semi-linear ;
- ▶ it is **linear with respect to the right variable** :  
 $\forall x \in E, \varphi_x : y \mapsto \varphi(x, y)$  is linear.

Choosing the left or right side for the sesquilinearity is arbitrary.

# Symmetry

The second property, symmetry, echoes the symmetry we observe when working with norms and distances.

# Symmetry

The second property, symmetry, echoes the symmetry we observe when working with norms and distances.

Like for the previous property, the symmetry is slightly different depending on whether the field is  $\mathbb{R}$  or  $\mathbb{C}$ .

# Symmetry

The second property, symmetry, echoes the symmetry we observe when working with norms and distances.

Like for the previous property, the symmetry is slightly different depending on whether the field is  $\mathbb{R}$  or  $\mathbb{C}$ .

$$\mathbb{K} = \mathbb{R}$$

We say that  $\varphi$  is **symmetric** if:

$$\forall (x, y) \in E, \quad \varphi(y, x) = \varphi(x, y).$$

# Symmetry

The second property, symmetry, echoes the symmetry we observe when working with norms and distances.

Like for the previous property, the symmetry is slightly different depending on whether the field is  $\mathbb{R}$  or  $\mathbb{C}$ .

$$\mathbb{K} = \mathbb{R}$$

We say that  $\varphi$  is **symmetric** if:

$$\forall (x, y) \in E, \quad \varphi(y, x) = \varphi(x, y).$$

$$\mathbb{K} = \mathbb{C}$$

We say that  $\varphi$  is **hermitian** if:

$$\forall (x, y) \in E, \quad \varphi(y, x) = \overline{\varphi(x, y)}.$$



# Symmetry

The second property, symmetry, echoes the symmetry we observe when working with norms and distances.

Like for the previous property, the symmetry is slightly different depending on whether the field is  $\mathbb{R}$  or  $\mathbb{C}$ .

$$\mathbb{K} = \mathbb{R}$$

We say that  $\varphi$  is **symmetric** if:

$$\forall (x, y) \in E, \quad \varphi(y, x) = \varphi(x, y).$$

$$\mathbb{K} = \mathbb{C}$$

We say that  $\varphi$  is **hermitian** if:

$$\forall (x, y) \in E, \quad \varphi(y, x) = \overline{\varphi(x, y)}.$$

We will see later why it is necessary to use conjugates when working with complex numbers.

## Positivity and definiteness

The last two properties will help establish the link between inner product and norms.

## Positivity and definiteness

The last two properties will help establish the link between inner product and norms.

We say that  $\varphi$  is **positive** if:

$$\forall x \in E, \quad \varphi(x, x) \in \mathbb{R}_+.$$

## Positivity and definiteness

The last two properties will help establish the link between inner product and norms. We say that  $\varphi$  is **positive** if:

$$\forall x \in E, \quad \varphi(x, x) \in \mathbb{R}_+.$$

Take care, it does not mean that  $\varphi$  only takes positive values. Indeed, for  $x \neq y$ , we can have  $\varphi(x, y) < 0$  over  $\mathbb{R}$ , and if  $\mathbb{K} = \mathbb{C}$  we can get any complex number. But, if we take the same argument both on the left and the right, the result will be a positive real number.

## Positivity and definiteness

The last two properties will help establish the link between inner product and norms. We say that  $\varphi$  is **positive** if:

$$\forall x \in E, \quad \varphi(x, x) \in \mathbb{R}_+.$$

Take care, it does not mean that  $\varphi$  only takes positive values. Indeed, for  $x \neq y$ , we can have  $\varphi(x, y) < 0$  over  $\mathbb{R}$ , and if  $\mathbb{K} = \mathbb{C}$  we can get any complex number. But, if we take the same argument both on the left and the right, the result will be a positive real number.

Moreover, we say that  $\varphi$  is **definite** if

$$\varphi(x, x) = 0 \iff x = 0_E.$$

## Positivity and definiteness

The last two properties will help establish the link between inner product and norms. We say that  $\varphi$  is **positive** if:

$$\forall x \in E, \quad \varphi(x, x) \in \mathbb{R}_+.$$

Take care, it does not mean that  $\varphi$  only takes positive values. Indeed, for  $x \neq y$ , we can have  $\varphi(x, y) < 0$  over  $\mathbb{R}$ , and if  $\mathbb{K} = \mathbb{C}$  we can get any complex number. But, if we take the same argument both on the left and the right, the result will be a positive real number.

Moreover, we say that  $\varphi$  is **definite** if

$$\varphi(x, x) = 0 \iff x = 0_E.$$

This property is to be paralleled with the separation axiom of distances and norms.

## Inner product: definition

We say that  $\varphi$  is an **inner product** over  $\mathbb{K}$  if  $\varphi$  is:

$$\mathbb{K} = \mathbb{R}$$

- ▶ bilinear
- ▶ symmetric
- ▶ positive
- ▶ definite

$$\mathbb{K} = \mathbb{C}$$

- ▶ sesquilinear (left or right)
- ▶ hermitian
- ▶ positive
- ▶ definite

## Inner product: definition

We say that  $\varphi$  is an **inner product** over  $\mathbb{K}$  if  $\varphi$  is:

$$\mathbb{K} = \mathbb{R}$$

- ▶ bilinear
- ▶ symmetric
- ▶ positive
- ▶ definite

$$\mathbb{K} = \mathbb{C}$$

- ▶ sesquilinear (left or right)
- ▶ hermitian
- ▶ positive
- ▶ definite

If so, we say that  $(E, +, \cdot, \varphi)$  is a structure of **inner product space**.



## Inner product: definition

We say that  $\varphi$  is an **inner product** over  $\mathbb{K}$  if  $\varphi$  is:

$$\mathbb{K} = \mathbb{R}$$

- ▶ bilinear
- ▶ symmetric
- ▶ positive
- ▶ definite

$$\mathbb{K} = \mathbb{C}$$

- ▶ sesquilinear (left or right)
- ▶ hermitian
- ▶ positive
- ▶ definite

If so, we say that  $(E, +, \cdot, \varphi)$  is a structure of **inner product space**.

If  $E$  is of finite dimension, we call it an **Euclidean space** if  $\mathbb{K} = \mathbb{R}$  and a **Hermitian space** if  $\mathbb{K} = \mathbb{C}$ .

## Standard inner products - $\mathbb{R}^n$ and $\mathbb{C}^n$

On  $\mathbb{R}^n$ , the standard inner product  $\langle \cdot, \cdot \rangle$  maps to the vectors  $x = (x_1, \dots, x_n)$  and  $y = (y_1, \dots, y_n)$  the scalar

$$\langle x, y \rangle = \sum_{i=1}^n x_i y_i.$$

## Standard inner products - $\mathbb{R}^n$ and $\mathbb{C}^n$

On  $\mathbb{R}^n$ , the standard inner product  $\langle \cdot, \cdot \rangle$  maps to the vectors  $x = (x_1, \dots, x_n)$  and  $y = (y_1, \dots, y_n)$  the scalar

$$\langle x, y \rangle = \sum_{i=1}^n x_i y_i.$$

On  $\mathbb{C}^n$ , the definition is almost similar:  $\langle \cdot, \cdot \rangle$  maps to the vectors  $x = (x_1, \dots, x_n)$  and  $y = (y_1, \dots, y_n)$  the scalar

$$\langle x, y \rangle = \sum_{i=1}^n \overline{x_i} y_i.$$

## Standard inner products - $\mathbb{R}^n$ and $\mathbb{C}^n$

On  $\mathbb{R}^n$ , the standard inner product  $\langle \cdot, \cdot \rangle$  maps to the vectors  $x = (x_1, \dots, x_n)$  and  $y = (y_1, \dots, y_n)$  the scalar

$$\langle x, y \rangle = \sum_{i=1}^n x_i y_i.$$

On  $\mathbb{C}^n$ , the definition is almost similar:  $\langle \cdot, \cdot \rangle$  maps to the vectors  $x = (x_1, \dots, x_n)$  and  $y = (y_1, \dots, y_n)$  the scalar

$$\langle x, y \rangle = \sum_{i=1}^n \overline{x_i} y_i.$$

Remark: the formula over  $\mathbb{C}$  still stands over  $\mathbb{R}$ , since the conjugate of a real number is itself.

## Standard inner products - $\mathbb{R}^n$ and $\mathbb{C}^n$

On  $\mathbb{R}^n$ , the standard inner product  $\langle \cdot, \cdot \rangle$  maps to the vectors  $x = (x_1, \dots, x_n)$  and  $y = (y_1, \dots, y_n)$  the scalar

$$\langle x, y \rangle = \sum_{i=1}^n x_i y_i.$$

On  $\mathbb{C}^n$ , the definition is almost similar:  $\langle \cdot, \cdot \rangle$  maps to the vectors  $x = (x_1, \dots, x_n)$  and  $y = (y_1, \dots, y_n)$  the scalar

$$\langle x, y \rangle = \sum_{i=1}^n \bar{x}_i y_i.$$

Remark: the formula over  $\mathbb{C}$  still stands over  $\mathbb{R}$ , since the conjugate of a real number is itself.

We can see that these functions check all the needed properties if we remember that for every  $z \in \mathbb{C}$ ,  $z \cdot \bar{z} = |z|^2 \in \mathbb{R}_+$ .

## Standard inner products - spaces of functions

Over spaces of smooth enough functions, we generally define the standard inner product as being, for functions with real values:

$$\langle f, g \rangle = \int_I f(t)g(t)dt$$

and for functions with complex values:

$$\langle f, g \rangle = \int_I \overline{f(t)}g(t)dt$$

## Standard inner products - spaces of functions

Over spaces of smooth enough functions, we generally define the standard inner product as being, for functions with real values:

$$\langle f, g \rangle = \int_I f(t)g(t)dt$$

and for functions with complex values:

$$\langle f, g \rangle = \int_I \overline{f(t)}g(t)dt$$

Of course, in order to be allowed to do so, the integrals must be defined, which happens particularly if we choose continuous or piecewise continuous functions defined over a segment.

## Induced norm

The way we defined inner products help us get the following interesting result:

**Proposition** [Norm induced by an inner product]

Let  $E$  be a vector space, together with an inner product  $\varphi$ .

Then, the map  $N_\varphi$  defined for every  $x \in E$  by  $N_\varphi(x) = \sqrt{\varphi(x, x)}$  is a norm over  $E$ .

It is called **norm induced by  $\varphi$**  over  $E$ .



## Induced norm

The way we defined inner products help us get the following interesting result:

**Proposition** [Norm induced by an inner product]

Let  $E$  be a vector space, together with an inner product  $\varphi$ .

Then, the map  $N_\varphi$  defined for every  $x \in E$  by  $N_\varphi(x) = \sqrt{\varphi(x, x)}$  is a norm over  $E$ .

It is called **norm induced by  $\varphi$**  over  $E$ .

The structure of inner product space thus comes together with built-in notions of distance and norm.

## Euclidean norms

- ▶ Over  $\mathbb{R}^n$  or  $\mathbb{C}^n$ , the norm induced by the standard inner product is the Euclidean norm  $\|\cdot\|_2$  :

$$\|x\|_2 = \left( \sum |x_i|^2 \right)^{\frac{1}{2}}$$

## Euclidean norms

- ▶ Over  $\mathbb{R}^n$  or  $\mathbb{C}^n$ , the norm induced by the standard inner product is the Euclidean norm  $\|\cdot\|_2$  :

$$\|x\|_2 = \left( \sum |x_i|^2 \right)^{\frac{1}{2}}$$

- ▶ Likewise, over  $C^0([a,b], \mathbb{R})$  or another such space, the norm induced by the standard inner product is the Euclidean norm  $\|\cdot\|_2$  :

$$\|x\|_2 = \left( \int_a^b |f(t)|^2 dt \right)^{\frac{1}{2}}$$



The two following results enable us to prove that the application  $N_\varphi$  induced by  $\varphi$  is a norm over  $E$ .

The two following results enable us to prove that the application  $N_\varphi$  induced by  $\varphi$  is a norm over  $E$ .

They had already been talked about during the tutorial about distances. Now you will know whence they come :-)

## Cauchy-Schwarz inequality

**Theorem** [Cauchy-Schwarz inequality]

Let  $E$  be an inner product space (over  $\mathbb{R}$  or  $\mathbb{C}$ ); we denote by  $\langle \cdot, \cdot \rangle$  its inner product and by  $\| \cdot \|$  the induced norm. Then:

$$| \langle x, y \rangle | \leq \|x\| \cdot \|y\|$$

## Cauchy-Schwarz inequality

**Theorem** [Cauchy-Schwarz inequality]

Let  $E$  be an inner product space (over  $\mathbb{R}$  or  $\mathbb{C}$ ); we denote by  $\langle \cdot, \cdot \rangle$  its inner product and by  $\| \cdot \|$  the induced norm. Then:

$$| \langle x, y \rangle | \leq \|x\| \cdot \|y\|$$

This result can be rewritten thus:

$$| \langle x, y \rangle | \leq \sqrt{\langle x, x \rangle} \sqrt{\langle y, y \rangle}.$$



## Cauchy-Schwarz inequality

### **Theorem** [Cauchy-Schwarz inequality]

Let  $E$  be an inner product space (over  $\mathbb{R}$  or  $\mathbb{C}$ ); we denote by  $\langle \cdot, \cdot \rangle$  its inner product and by  $\| \cdot \|$  the induced norm. Then:

$$| \langle x, y \rangle | \leq \|x\| \cdot \|y\|$$

This result can be rewritten thus:

$$| \langle x, y \rangle | \leq \sqrt{\langle x, x \rangle} \sqrt{\langle y, y \rangle}.$$

The proof of this result is interesting: if we evaluate the expression  $\langle x + ty, x + ty \rangle$ , we get an degree 2 equation of the variable  $t$ , that cannot take strictly negative values (positivity), meaning that its discriminant is negative (possibly 0 iff  $x + ty = 0$ ). This discriminant gives the statement of the Cauchy-Schwarz inequality.

## Minkowski inequality

The Minkowski inequality can be derived from the Cauchy-Schwarz inequality:

**Theorem** [Minkowski inequality]

With the notations used for the previous theorem:

$$\sqrt{\langle x+y, x+y \rangle} \leq \sqrt{\langle x, x \rangle} + \sqrt{\langle y, y \rangle}$$

That is to say:

$$\|x+y\| = \|x\| + \|y\|.$$

## Minkowski inequality

The Minkowski inequality can be derived from the Cauchy-Schwarz inequality:

**Theorem** [Minkowski inequality]

With the notations used for the previous theorem:

$$\sqrt{\langle x+y, x+y \rangle} \leq \sqrt{\langle x, x \rangle} + \sqrt{\langle y, y \rangle}$$

That is to say:

$$\|x+y\| = \|x\| + \|y\|.$$

The Minkowski inequality establishes that the norm induced by an inner product checks the property of triangle inequality.

## Minkowski inequality

The Minkowski inequality can be derived from the Cauchy-Schwarz inequality:

**Theorem** [Minkowski inequality]

With the notations used for the previous theorem:

$$\sqrt{\langle x+y, x+y \rangle} \leq \sqrt{\langle x, x \rangle} + \sqrt{\langle y, y \rangle}$$

That is to say:

$$\|x+y\| = \|x\| + \|y\|.$$

The Minkowski inequality establishes that the norm induced by an inner product checks the property of triangle inequality.

You already proved it for particular cases during the first tutorial, when proving the triangle inequality for the Euclidean norms.



# Orthogonality

This will not be an exhaustive lesson into the concept of orthogonality, just an introduction to the interesting concept to explain the theory of Fourier series.

# Orthogonality

This will not be an exhaustive lesson into the concept of orthogonality, just an introduction to the interesting concept to explain the theory of Fourier series. We for instance will not dwell on the notions of orthogonality to a space and their problems, but instead will focus on the orthogonality between vectors, only to introduce the orthogonal projection.

# Orthogonality

This will not be an exhaustive lesson into the concept of orthogonality, just an introduction to the interesting concept to explain the theory of Fourier series.

We for instance will not dwell on the notions of orthogonality to a space and their problems, but instead will focus on the orthogonality between vectors, only to introduce the orthogonal projection.

Similarly, we will not talk about the Gram-Schmidt process, it is to be studied later during the lessons about matrix decompositions.



# Orthogonality

## **Definition** [Orthogonality]

In an inner product space  $E$  together with an inner product  $\langle \cdot, \cdot \rangle$ , we say that two vectors  $x$  and  $y$  are **orthogonal** if  $\langle x, y \rangle = 0$ .

We say that a set of vectors  $(e_1, \dots, e_n)$  is (pairwise) orthogonal if, for every  $(i, j)$  such that  $i \neq j$ ,  $\langle e_i, e_j \rangle = 0$ .

If, moreover, for every  $i$  then  $\langle e_i, e_i \rangle = 1$  then the set is said to be **orthonormal**.

# Orthogonality

## **Definition** [Orthogonality]

In an inner product space  $E$  together with an inner product  $\langle \cdot, \cdot \rangle$ , we say that two vectors  $x$  and  $y$  are **orthogonal** if  $\langle x, y \rangle = 0$ .

We say that a set of vectors  $(e_1, \dots, e_n)$  is (pairwise) orthogonal if, for every  $(i, j)$  such that  $i \neq j$ ,  $\langle e_i, e_j \rangle = 0$ .

If, moreover, for every  $i$  then  $\langle e_i, e_i \rangle = 1$  then the set is said to be **orthonormal**.

For instance, in  $\mathbb{R}^2$  together with its standard inner product, the vectors  $(0, 1)$  and  $(1, 0)$  are orthogonal. They even form an orthonormal basis of  $\mathbb{R}^2$ .

# Orthogonality

## **Definition** [Orthogonality]

In an inner product space  $E$  together with an inner product  $\langle \cdot, \cdot \rangle$ , we say that two vectors  $x$  and  $y$  are **orthogonal** if  $\langle x, y \rangle = 0$ .

We say that a set of vectors  $(e_1, \dots, e_n)$  is (pairwise) orthogonal if, for every  $(i, j)$  such that  $i \neq j$ ,  $\langle e_i, e_j \rangle = 0$ .

If, moreover, for every  $i$  then  $\langle e_i, e_i \rangle = 1$  then the set is said to be **orthonormal**.

For instance, in  $\mathbb{R}^2$  together with its standard inner product, the vectors  $(0, 1)$  and  $(1, 0)$  are orthogonal. They even form an orthonormal basis of  $\mathbb{R}^2$ .

In  $\mathbb{R}^2$ , you already know orthogonality as perpendicularity.

# Orthogonality

## **Definition** [Orthogonality]

In an inner product space  $E$  together with an inner product  $\langle \cdot, \cdot \rangle$ , we say that two vectors  $x$  and  $y$  are **orthogonal** if  $\langle x, y \rangle = 0$ .

We say that a set of vectors  $(e_1, \dots, e_n)$  is (pairwise) orthogonal if, for every  $(i, j)$  such that  $i \neq j$ ,  $\langle e_i, e_j \rangle = 0$ .

If, moreover, for every  $i$  then  $\langle e_i, e_i \rangle = 1$  then the set is said to be **orthonormal**.

For instance, in  $\mathbb{R}^2$  together with its standard inner product, the vectors  $(0, 1)$  and  $(1, 0)$  are orthogonal. They even form an orthonormal basis of  $\mathbb{R}^2$ .

In  $\mathbb{R}^2$ , you already know orthogonality as perpendicularity.

Orthogonality is an important property when it comes to get decompositions (in a basis for instance): it ensures that no mixed products will pollute the calculations involving the inner product.

## Orthogonal supplementary subspace

**Proposition** [Orthogonal supplementary subspace]

Let  $F$  be a vector subspace of finite dimension of  $E$ . Then the set of vectors of  $E$  that are orthogonal to every vector of  $F$ , called **orthogonal of  $F$**  and denoted by  $F^\perp$ , is a supplementary subspace of  $F$  in  $E$ .

## Orthogonal supplementary subspace

### **Proposition** [Orthogonal supplementary subspace]

Let  $F$  be a vector subspace of finite dimension of  $E$ . Then the set of vectors of  $E$  that are orthogonal to every vector of  $F$ , called **orthogonal of  $F$**  and denoted by  $F^\perp$ , is a supplementary subspace of  $F$  in  $E$ .

Careful, if we choose  $F$  of infinite dimension, this result is not always true. However, this will be the case for the well-chosen spaces in which we will process to the Fourier decompositions.

## Orthogonal supplementary subspace

### **Proposition** [Orthogonal supplementary subspace]

Let  $F$  be a vector subspace of finite dimension of  $E$ . Then the set of vectors of  $E$  that are orthogonal to every vector of  $F$ , called **orthogonal of  $F$**  and denoted by  $F^\perp$ , is a supplementary subspace of  $F$  in  $E$ .

Careful, if we choose  $F$  of infinite dimension, this result is not always true. However, this will be the case for the well-chosen spaces in which we will process to the Fourier decompositions.

Remark: the Gram-Schmidt process is a constructive method to build an orthonormal basis of a vector subspace of finite dimension, ensuring that such bases exist.

## Orthogonal projection

**Definition** [Orthogonal projection]

Let  $F$  be a vector subspace of  $E$  with an orthogonal supplementary subspace  $F^\perp$ .

We call orthogonal projection onto  $F$  the function  $p_F$  whose restriction to  $F$  is the identity function of  $F$  and whose kernel is  $F^\perp$ .



## Orthogonal projection

**Definition** [Orthogonal projection]

Let  $F$  be a vector subspace of  $E$  with an orthogonal supplementary subspace  $F^\perp$ .

We call orthogonal projection onto  $F$  the function  $p_F$  whose restriction to  $F$  is the identity function of  $F$  and whose kernel is  $F^\perp$ .

Remark: we then have  $x = p_F(x) + (x - p_F(x))$  and we can check that  $p_F(x)$  and  $x - p_F(x)$  are orthogonal.

## Orthogonal projection

### **Definition** [Orthogonal projection]

Let  $F$  be a vector subspace of  $E$  with an orthogonal supplementary subspace  $F^\perp$ .

We call orthogonal projection onto  $F$  the function  $p_F$  whose restriction to  $F$  is the identity function of  $F$  and whose kernel is  $F^\perp$ .

Remark: we then have  $x = p_F(x) + (x - p_F(x))$  and we can check that  $p_F(x)$  and  $x - p_F(x)$  are orthogonal.

If  $F$  is of finite dimension, we can even write:

if  $(f_1, \dots, f_p)$  is an orthogonal basis of  $F$ , then

$$p_F(x) = \sum_{i=1}^p \frac{\langle x, f_i \rangle}{\langle f_i, f_i \rangle} f_i.$$

If the basis  $(f_1, \dots, f_p)$  is orthonormal, we get

$$p_F(x) = \sum_{i=1}^p \langle x, f_i \rangle f_i.$$

## Pythagoras's theorem

**Theorem** [Pythagoras's theorem]

If  $x$  and  $y$  are orthogonal, then  $\langle x+y, x+y \rangle = \langle x, x \rangle + \langle y, y \rangle$ , that is to say:

$$\|x+y\|^2 = \|x\|^2 + \|y\|^2.$$

## Pythagoras's theorem

### **Theorem** [Pythagoras's theorem]

If  $x$  and  $y$  are orthogonal, then  $\langle x+y, x+y \rangle = \langle x, x \rangle + \langle y, y \rangle$ , that is to say:

$$\|x+y\|^2 = \|x\|^2 + \|y\|^2.$$

This result can be generalised to any orthogonal set: if the set  $(e_1, \dots, e_n)$  is orthogonal then

$$\left\| \sum_{i=1}^n e_i \right\|^2 = \sum_{i=1}^n \|e_i\|^2.$$

## Pythagoras's theorem

### **Theorem** [Pythagoras's theorem]

If  $x$  and  $y$  are orthogonal, then  $\langle x+y, x+y \rangle = \langle x, x \rangle + \langle y, y \rangle$ , that is to say:

$$\|x+y\|^2 = \|x\|^2 + \|y\|^2.$$

This result can be generalised to any orthogonal set: if the set  $(e_1, \dots, e_n)$  is orthogonal then

$$\left\| \sum_{i=1}^n e_i \right\|^2 = \sum_{i=1}^n \|e_i\|^2.$$

It is a rewriting of the Pythagoras's theorem you already know for right-angled triangles, but in a general inner product space and more than two dimensions.