The Economics of Inward Migration: Relocation versus Densification for Computation-Maximizing Civilizations

Tim Swanson
The Omega Centauri Society / Post Oak Labs
tim@postoaklabs.com

October 2026

Draft v2.7 (last revised 2026-10-24) — fourth paper of the set; companion to *The Macro Transcension Hypothesis* (Paper A), the inward-migration review (Paper B),
the Omega Centauri campaign (Paper C), the engineered-IMBH systems paper (Paper E), the joint accretion bound (Paper F),
the residual X-ray census (Paper G), and the mass-tension paper (Paper H);
prepared for omegacentauri.me

Abstract

The inward-migration resolutions of the Fermi paradox hold that computation-optimizing civilizations relocate to thermodynamically privileged environments, with rapidly spinning intermediate-mass black holes (IMBHs) in dense old clusters as the strongest candidate destination. Bostrom (2003) priced the opportunity cost of delayed expansion and Bennett et al. (2019) showed that dormancy confers no thermodynamic advantage, but relocation itself, the abandonment of accumulated local infrastructure for a transit of 10^5–10^6 years toward a large deferred payoff, is one for which we are not aware of a prior pricing. This paper treats that trade as a decision problem over a common utility (discounted integrated computation) with three strategies: stay-and-densify (Matrioshka-style local engineering), migrate (beamed-sail relocation to the nearest suitable IMBH cluster), and seed-and-stay (a self-replicating seed dispatched while densification continues at home). Using payoff kernels assembled from established physics, we derive closed-form crossover conditions.

The central result is a threshold on the effective discount-plus-hazard rate: migration dominates densification whenever $\rho+\lambda < \ln(Gp_s)/\tau$, provided $Gp_s > 1$, where G is the destination computation-rate multiplier (10^3–10^9 on power alone, from an ADIOS-suppressed floor to full Eddington capacity, depending on fuel imports), p_s the transit survival probability, and τ the transit-plus-construction time. At fiducial parameters the threshold is $\rho + \lambda \lesssim 2 \times 10^{-4}$ yr^{-1}: any lineage whose combined discount-plus-hazard half-life exceeds roughly 3,500 years should migrate. Under additive utility, symmetric valuation, and a seed fidelity within the tolerance $(1 - f)(Gp_s)^{x-1}$ of the migrant's, with $x = (\rho + \lambda)/\rho^*$ (about 4.5×10^{-5} at $x = 0.5$, rising to unity at the threshold and falling to 2×10^{-9} as $\rho + \lambda$ goes to zero), the seed-and-stay hybrid dominates both pure strategies throughout the migration-favourable region and in a band extending modestly beyond the threshold (to $\rho + \lambda \approx 1.7\,\rho^*$ at a physical seed cost $f = 10^{-6}$), because seed mass is a negligible fraction of local output; beyond that band the exponential discount on the deferred payoff extinguishes the seed's value. Embedding the decision rule in a mixed population quantifies how much inward migration thins the expected loud population, and yields a new population-level residue: the predicted sky ratio of Matrioshka-type infrared sources to quiet-cluster systems, for which archival infrared nulls supply the joint-likelihood frame for a limit conditional on an assumed residue-to-formation ratio and maintenance lifetimes. All

results are conditional on the optimization premise shared by the hypothesis family, on symmetric valuation of drifted successors, and on power-only destination accounting; the contribution is the pricing structure, not the premise.

Keywords: Fermi paradox; SETI; technosignatures; transcension; migration; decision theory; discounting; intermediate-mass black holes; Dyson spheres

Contents

1 Introduction **4**
- 1.1 The pricing gap . . . 4
- 1.2 Epistemic register . . . 4
- 1.3 Structure . . . 5

2 The decision problem **5**
- 2.1 Agent, utility, and strategies . . . 5
- 2.2 Parameters . . . 6

3 Payoff kernels **6**
- 3.1 Strategy S: the Matrioshka ceiling . . . 6
- 3.2 Strategy M: the IMBH platform . . . 8
- 3.3 Discounted values . . . 9

4 Crossover conditions **11**
- 4.1 Migrate versus stay . . . 11
- 4.2 Meaning of the threshold . . . 13

5 The hybrid result **15**
- 5.1 Dominance . . . 15
- 5.2 Consequences . . . 15

6 Population consequences **17**

7 Observational corollaries: the residue ratio **18**

8 Limits and objections **20**

9 Conclusion **22**

1 Introduction

1.1 The pricing gap

Three papers precede this one. The Macro Transcension Hypothesis (Swanson, 2026c, hereafter Paper A) argues that rapidly spinning massive black holes in dense old stellar systems are the global optimum among present-day environments for long-term computation, and identifies Omega Centauri as the most accessible test system. A critical review of the inward-migration family (Swanson, 2026b, hereafter Paper B) places that hypothesis among six decades of related proposals and grades their falsifiability. An observational companion (Swanson, 2026d, hereafter Paper C) specifies the instrument-matched campaign. Paper B also catalogued the family's open problems, and its third problem motivates this paper directly: the economics of the migration itself has never been worked out.

The gap is specific. Bostrom (2003) priced delay: every century of postponed expansion forgoes an irrecoverable harvest of free energy. Bennett et al. (2019) showed that the erasure cost aestivation defers can be paid at any epoch by transferring entropy into existing non-maximal-entropy reservoirs, and that this is not improved by waiting for a colder background, so the aestivation strategy of Sandberg et al. (2016) gains nothing from dormancy. The expansion decision itself has economic precedent: Hosek (2007) proposed a resolution of the paradox from common economic assumptions, in which rational resource allocation leads planet-bound civilizations to forgo interstellar exploration and colonization altogether. We are not aware of a prior pricing of relocation. A lineage that migrates abandons its accumulated local infrastructure, spends 10^5–10^6 years in transit, accepts a risk of total loss en route, and only then begins to collect a payoff that is, by the physics of Paper A, orders of magnitude larger per unit of everything. Whether that trade clears depends on quantities the inward-family literature has so far left unmodelled: discount rates, hazard rates, transit risk, and the option of sending a seed instead of moving.

This matters beyond bookkeeping. If migration clears the trade only for implausibly patient agents, the inward family loses its force as a Fermi solution regardless of the destination physics. If it clears easily, the family's incomplete-compliance problem (Paper B, problem P4) sharpens into a quantitative question: what fraction of lineages must migrate for the observed silence to follow? Both outcomes constrain the family.

1.2 Epistemic register

Epistemic status: This paper is decision theory over established physics. It makes no claim that extraterrestrial intelligence exists. Every result is conditional on the optimization premise (P1 of Paper A): that some long-lived technological lineages maximize long-term computation. The contribution is the pricing structure and its observational corollaries.

The paper makes three claims:

1. **(Decision analysis.)** Under the stated utility and payoff kernels, the choice among stay, migrate, and seed-and-stay reduces to closed-form threshold conditions on the discount rate, hazard rate, transit time, and destination multiplier (Section 4).

2. **(Hybrid boundary.)** Under additive utility and the seed-fidelity assumption made explicit in Section 5 ($p_d \geq p_d^{\mathrm{M}}$, the seed drifts no more than a deliberating migrant), the seed-and-stay hybrid weakly dominates both pure strategies wherever migration clears, a near-immediate consequence of a nearly free seed; the substantive result is the narrowness of the band beyond the threshold in which the hybrid still pays, $\rho + \lambda \approx 1.7\,\rho^*$ at seed cost $f = 10^{-6}$, because the exponential discount on the deferred payoff closes the window within a factor of two of ρ^* (Section 5).

3. **(Population corollary.)** Folding the decision rule into a mixed population yields the expected thinning of the loud population and a new residue, the Matrioshka-to-quiet-cluster sky ratio, that archival infrared surveys already constrain (Sections 6–7).

We do not claim the parameter priors are measurable; the contribution is the structure of the crossover the analysis exposes, with point estimates serving only as orientation. We flag model dependence throughout (Section 8).

1.3 Structure

Section 2 states the decision problem. Section 3 assembles the payoff kernels from the physics of Paper A. Section 4 derives the crossover conditions and Section 5 the hybrid-dominance result. Section 6 treats the population level, Section 7 the observational corollaries, Section 8 the limits, and Section 9 concludes.

2 The decision problem

2.1 Agent, utility, and strategies

Consider a technological lineage established at a planetary system a distance d from the nearest system satisfying the Paper A selection criteria (an IMBH of 10^3–$10^5\,M_\odot$ in a dense, old, quiescent cluster; criterion 3, the dense low-mass stellar reservoir, is what excludes nearer isolated stellar-mass holes, which lack a fuel supply at the Eddington scale this analysis assumes). Following the family's shared premise, the lineage values integrated computation. Its utility over a policy π is

$$U(\pi) \;=\; \int_0^\infty C_\pi(t)\, e^{-\rho t}\, dt, \tag{1}$$

where $C_\pi(t)$ is the computation rate delivered at time t (bit-erasures per second, the binding currency by the Landauer accounting of Paper A, Section 4.2), t is measured in years throughout (ρ, λ, and τ below are all yr and yr^{-1}; the erasures-per-second rates of Section 3 are annualized by the implicit factor of $3.156 \times 10^7\ \mathrm{s\,yr}^{-1}$ before entering U, which cancels in every dimensionless ratio used below and is stated once here rather than at each occurrence), and $\rho \geq 0$ an exponential discount rate. We use exponential discounting for tractability and revisit hyperbolic alternatives (Laibson, 1997) in Section 8; the patient limit $\rho \to 0$ recovers the Ramsey (1928) zero-discount benchmark, an argument Bostrom (2003) makes for agents with unbounded horizons.

For readers who have not worked with exponential discounting: the parameter ρ measures impatience. A payoff delivered τ years from now is worth $e^{-\rho\tau}$ times the same payoff today. At the fiducial $\tau = 10^5$ yr, even a modest $\rho = 10^{-4}\ \mathrm{yr}^{-1}$ discounts the destination payoff by $e^{-10} \approx 5 \times 10^{-5}$ — a factor of 20,000. The migration question is whether the destination advantage G overcomes this factor, and the answer depends logarithmically on G but linearly on ρ and τ.

Three strategies span the space of single-destination policies:

S (stay and densify). Remain at the home system and build toward the local optimum: Dyson-type collection feeding Matrioshka-style nested computing shells (Dyson, 1960; Bradbury, 1999). Payoff begins almost immediately and is capped by stellar output and radiator physics.

M (migrate). Relocate the lineage by beamed-sail transport (Lubin, 2016) to the IMBH system, build the Paper A staged architecture, and compute there. Payoff is deferred by the transit-plus-construction time τ and discounted by a survival probability p_s, but multiplied by the destination gain G.

H (seed and stay). Dispatch a self-replicating seed payload (Freitas, 1980; Armstrong and Sandberg, 2013) of negligible mass while executing S at home; the seed bootstraps the destination in parallel. The lineage pays a small fraction f of local output for the launch infrastructure and collects both payoffs, with the deferred term weighted by a goal-drift valuation p_d introduced in Section 3; the seed's transit hazards (radiation damage, replication error) also differ in kind from those facing a deliberating lineage in transit.

Strategy H is simply the architecture Paper A already assumes (gram-to-kilogram scouts, then seed factories), read as an economic option rather than a mission profile. What has not been stated before is its dominance structure (Section 5). Multi-destination seeding, one lineage dispatching $k > 1$ seeds to distinct clusters, generalizes H directly: the per-lineage cost scales to kf and the hybrid condition of Section 5 applies seed by seed. This paper does not model that generalization separately.

2.2 Parameters

Table 1 collects the parameters and fiducial values. Transit time uses Paper A's beamed-sail speeds (0.1–0.2 c; Hoang et al. 2017, conservative against the $> c/4$ of gram-scale probes in Lubin 2016) and the distance to ω Cen (5.49 kpc; Häberle et al. 2025) as the fiducial destination, giving $\tau_{\text{transit}} \approx 0.9$–$1.8\times10^5$ yr; we fold construction ($\lesssim 10^3$ yr to first ISCO-swarm operation; Paper A, staged-construction appendix) into a single $\tau = 10^5$ yr fiducial. The bare cruise time at the tabulated $v = 0.15c$ is $d/v = 1.19 \times 10^5$ yr, so the round fiducial corresponds to an effective $\approx 0.18c$; Table 1 states the bare number so the rows can be checked independently. We keep $\tau_{\text{fid}} = 10^5$ yr as the printed fiducial rather than the bare 1.19×10^5 yr, so the threshold understates the true transit time by at most 19 per cent before construction time is added back in; the residence-survival debit is insensitive to this choice, differing by less than 2×10^{-6} between the two readings of τ. Beamed-sail launch provides no deceleration at an unprepared destination, and braking by magsail (Zubrin and Andrews, 1991; Perakis and Hein, 2016) or by photon-assisted maneuvers (Heller and Hippke, 2017) can multiply the cruise time by a factor of a few; because the threshold of Section 4 scales linearly in $1/\tau$, we propagate this allowance explicitly there (Table 2) rather than absorbing it into the range. The hazard rate λ prices the goal-stability problem (Paper B, problem P2): the probability per year that the lineage ceases to pursue the plan, through value drift, fragmentation, or extinction. We do not claim to know λ; it enters the thresholds additively with ρ, which is the analytically convenient way to carry an unknown.

See Paper F App. A for the shared-parameter table across the set; this paper's distance fiducial $d = 5.49$ kpc matches the set consensus there.

3 Payoff kernels

Epistemic status: This section assembles rates from established physics, detailed in Paper A Section 4; nothing here is new physics. The kernels deliberately understate the migration payoff, for reasons given below.

3.1 Strategy S: the Matrioshka ceiling

A complete Dyson-type collector intercepts the stellar output, $L_* = 3.8 \times 10^{26}$ W for a solar analogue (the modern solar constant; Dyson 1960 motivates the collector concept but prints the rounder 4×10^{26} W). Computation against the ambient sink is bounded by the Landauer cost at the effective radiator temperature T_{rad} (Landauer, 1961; Bradbury, 1999): the innermost shells run hot and fast, the outermost approach the

Table 1: Decision parameters and fiducial values. Two entries carry cross-paper dependencies: the residence-survival figure imports Paper E's flyby accounting, whose operative channel is now the bound-cusp diffusion floor rather than the (non-operative) adiabatically corrected unbound-field curve, and is bimodal rather than a single number (Section 4), and the $G = 10^6$ variant imports Paper E's ambient-supply estimate, which falls to $G \approx 10^3$ under Paper E's ADIOS-suppressed ceiling of $2.6 \times 10^3\ L_\odot$ (Swanson, 2026a) (Section 3).

Symbol	Meaning	Fiducial	Source / rationale
d	distance to destination	5.49 kpc	ω Cen (Häberle et al., 2025)
v	transit speed	$0.15\,c$	beamed sail (Lubin, 2016; Hoang et al., 2017)
τ	transit + construction	10^5 yr	bare $d/v = 1.19\times10^5$ yr at $0.15c$; fiducial $\approx$ effective $0.18c$
p_s	transit survival probability (deliberating lineage)	0.5	placeholder; Hoang et al. (2017) gives erosion depths only
p_s^{seed}	transit survival probability (seed payload)	0.5 adopted pending dedicated modelling	seed hazards (radiation, replication error) differ in kind from lineage transit hazards (Section 3); no separate estimate exists, so $p_s^{\mathrm{seed}} = p_s$ is a placeholder, not a derived equality
G	destination rate multiplier	10^9	power ratio; 10^6 without fuel imports, 10^3 under ADIOS suppression (Section 3)
ρ	discount rate	free	(Ramsey, 1928; Bostrom, 2003)
λ	goal-stability hazard rate	free	Paper B, problem P2
f	seed cost, fraction of the discounted S payoff U_{S} diverted (a permanent fractional tax; a one-time launch cost of f years of output is equivalent to $f(\rho+\lambda)$)	10^{-6} adopted (physical bound $\ll 10^{-3}$)	gram–kg payloads (Lubin, 2016; Freitas, 1980)
p_d	goal-drift valuation of seed output	free, $\lesssim 1$ by engineering	Section 3
p_d^{M}	goal-drift valuation of a deliberating migrant's own output	free	rescues $U_{\mathrm{H}} \geq U_{\mathrm{M}}$ only if $p_d \geq p_d^{\mathrm{M}}$ (Section 5)

CMB floor $T_\gamma = 2.7$ K,[1] and the whole stack is radiator-limited, the constraint Bradbury (1999) identified as binding. An upper bound that ignores every engineering loss is

$$C_S \leq \frac{L_*}{k_B T_\gamma \ln 2} \simeq 1.5 \times 10^{49} \text{ erasures s}^{-1}, \tag{2}$$

where the denominator, $k_B T_\gamma \ln 2 \simeq 2.6 \times 10^{-23}$ J, is the minimum energy per erasure against the CMB sink at $T_\gamma = 2.7$ K. This bound is available after a build time we absorb into the timeline as effectively immediate on the scales of interest (10^2–10^3 yr; Armstrong and Sandberg 2013). Strategy S also faces a slow ceiling decay (stellar evolution) that we neglect: neglecting it favours S, which is the conservative direction for our conclusion.

Equation (2) is not attainable even in principle, and it matters why not. Rejecting the full L_* at $T_{rad} \to T_\gamma$ requires zero net flux through infinite radiator area; the standard trade is that radiated power scales as T_{rad}^4 while the erasure cost scales linearly in T_{rad}, so finite architectures optimize at T_{rad} a few times T_γ and forfeit one to two orders of magnitude in C_S (Sandberg et al., 2016; Bennett et al., 2019; Badescu and Cathcart, 2000). The nesting itself has been priced: applying the Landsberg formalism to Dyson spheres as work extractors and computational engines, Wright (2023) finds little to no advantage in nested shells and concludes that the best use of mass is generally a very small, hot sphere, undercutting the idealization from which Strategy S takes its name. We nonetheless carry the unattainable bound, because any downward correction to C_S raises the destination multiplier G and moves every threshold below in migration's favour: Eq. (2) is the conservative choice for the stay option.

3.2 Strategy M: the IMBH platform

At the destination, the sustainable power is the Eddington luminosity of the host, $L_{Edd} \simeq 2.5 \times 10^{35}$ W at $2 \times 10^4\, M_\odot$ (Paper A, Eq. 2), a factor $\sim 10^9$ (6.6×10^8, rounded) over the solar Dyson ceiling. The erasure cost against a horizon sink falls by a further factor $\sim 10^{12}$ (Paper A, Section 4.2), but that factor is a floor-to-floor comparison of theoretical erasure minima, not a realized rate; Paper A (§4.2) is explicit that the realized gain, a per-erasure cost ratio against the 2.7 K sky, is about 10^3 in a fed flow and about 10^7 in the ambient medium if the carrier modes are kept in the vacuum state ($N_{eff} \ll 1$) from emitter to horizon, and a factor 20 deficit (~ 0.05 on the conservative budget) if they are thermally occupied by the 2.7 K sky, and states that the $\sim 10^{12}$ figure should not be quoted "wherever the realized gain is meant." $G \equiv C_M/C_S$ is a ratio of delivered computation rates, so the erasure-floor factor does not belong in it unless realized: the admissible multiplier on Eq. (2) is the power ratio alone, times the realized (not floor-to-floor) erasure gain of $\sim 10^3$–10^7 under the cold-mode condition, giving a Paper-A-compliant realized band of 6.6×10^{11}–6.6×10^{15} (rounded, $\sim 10^{12}$–10^{16}), and the power ratio alone, 6.6×10^8, if the modes are thermally occupied. We adopt

$$G \equiv \frac{C_M}{C_S} = 10^9 \tag{3}$$

as the fiducial, crediting the migration payoff with the power ratio only (rounded up from the recomputed 6.6×10^8, a +2.1% anti-conservative rounding on ρ^* that is dwarfed by the order-of-magnitude uncertainties carried elsewhere in G). The Eddington figure is a capacity, not a fuel supply: the infrastructure companion (Swanson, 2026a, hereafter Paper E) budgets the ambient Bondi-fed accretion available in a gas-poor

[1] The floor can be approached without leaving home: Garrett (2026) argue that computation-dominated civilizations should dissipate at the lowest ambient temperature, approaching the CMB floor with vast cold radiators whose area grows as T^{-4} and whose grey, line-free waste heat emerges in the submillimetre, constrained by archival surveys at the $\sim 10^{20}$ W level. Equation (2) credits S with that floor throughout, so G is measured against the best non-relocating cold computer.

cluster at $\sim 8 \times 10^5\, L_\odot$, so the power-alone multiplier is $G \sim 10^6$ unless the civilization imports fuel, which raises G at the cost of visibility (enhanced accretion luminosity is one of Paper E's residues). The recurring logistics of that import programme, chiefly angular-momentum removal from infalling mass (priced in Paper A), are not carried in the kernel; they are absorbed into the conservatism of the power-only G, while the far smaller seed cost f is carried explicitly because it gates the hybrid boundary of Section 5. At $G = 10^6$ the threshold of Section 4 becomes $\rho^* \simeq \ln(5 \times 10^5)/10^5$ yr $\approx 1.3 \times 10^{-4}$ yr^{-1}, against 2.0×10^{-4} at $G = 10^9$: three orders of magnitude in the multiplier move the threshold by a factor of 1.5, the logarithmic robustness on which the analysis leans. The $G = 10^6$ figure imports Paper E's ambient-supply estimate and moves with it: under the ADIOS outflow-suppression standard for hot low-Eddington flows (Swanson, 2026a), the delivered rate falls to a ceiling of $2.6 \times 10^3\, L_\odot$, giving $G \approx 2.6 \times 10^3$ and moving the threshold to $\rho^* = \ln(1.3 \times 10^3)/10^5$ yr $\approx 0.7 \times 10^{-4}$ yr^{-1} (0.6×10^{-4} at the rounded $G = 10^3$ that serves as the lower end of the adopted bracket), the same order. Two further reasons for the understatement. First, horizon-sink disposal shifts the binding constraint to waste-heat transport (Paper A, Section 4.2), whose engineering efficiency is unmodelled; using the power ratio alone keeps the kernel on demonstrated physics. Second, every conclusion below strengthens monotonically in G, so a conservative G makes the thresholds conservative. Readers preferring the full realized-erasure-gain stack may substitute $G = 6.6 \times 10^{11}$–6.6×10^{15} (the Paper-A-compliant band, valid under the cold-mode condition of Paper A); the thresholds move by the logarithm, i.e. by a factor of ≤ 1.8 (Section 4). The floor-to-floor $\sim 10^{21}$ figure is a thermodynamic ceiling on erasure cost alone, not a realized payoff, and is not substituted here. One accounting caveat: if computation at either site is performed reversibly (Bennett, 1973), erasures are not proportional to useful computation and the two sites' conversion factors could in principle differ. Because G is a ratio of erasure budgets, architecture-dependent reversibility factors cancel to first order when both sites use the same logic family, which is the natural assumption for a single lineage. Three inputs are imported from Paper E and its companions: the ambient Bondi ceiling, the range of G, and the residence-survival fractions of Section 4. Those companions are preprints and are not yet refereed, so the figures that depend on them are provisional.

3.3 Discounted values

With C_S available from $t = 0$ and C_M from $t = \tau$ with probability p_s, and with the hazard rate entering as an additional exponential attrition on plan continuation, the strategy values under Eq. (1) are

$$U_S = \frac{C_S}{\rho + \lambda}, \tag{4}$$

$$U_M = p_s\, e^{-(\rho+\lambda)\tau}\, \frac{G\, C_S}{\rho + \lambda}, \tag{5}$$

$$U_H = (1 - f)\, U_S \;+\; U_M. \tag{6}$$

Dividing Eq. (5) by Eq. (4) gives the migrate-to-stay value ratio in closed form,

$$\frac{U_M}{U_S} \;=\; G\, p_s\, e^{-(\rho+\lambda)\tau}, \tag{7}$$

in which the $1/(\rho+\lambda)$ prefactors cancel and λ survives only inside the transit exponential. That cancellation depends on what λ prices and on how drifted successors are valued. If λ is *lineage death* (extinction or fragmentation that terminates the payoff stream wherever the lineage sits), it attrites both strategies identically over the exploitation phase and the prefactors cancel as written. If instead λ is *plan abandonment* (value drift away from computation maximization), the question is whether the deciding lineage credits

computation performed by drifted successors. We adopt the symmetric valuation: a drifted lineage that has arrived still computes, and its output is credited on the same terms as a drifted homebody's, consistent with the redistribution reading of Section 4. Under that valuation, drift voids the migration payoff only when it strikes during transit (a plan abandoned mid-cruise delivers nothing anywhere), which is the same structure as lineage death: the ratio is again $G p_s\, e^{-(\rho+\lambda)\tau}$ and the two interpretations yield the same threshold. An asymmetric valuation, in which only undrifted migrant output counts while any homebody output counts, would multiply Eq. (7) by $\rho/(\rho+\lambda)$; we do not adopt it, because it values the same drift event differently at the two sites for no stated reason.

Equations (4)–(5) also let both payoff streams run forever, and the physical horizons differ. Strategy S is bounded by the host star's remaining main-sequence life, $T_S \approx 5 \times 10^9$ yr for a solar analogue. Star lifting extends this horizon substantially only for fully convective low-mass hosts, to $T_S \sim 10^{11}$–10^{12} yr; a solar analogue, the fiducial host here, gains only $\sim 3 \times 10^9$ yr from the same mechanism (Scoggins and Kipping, 2023). Strategy M at the Eddington rate consumes roughly one solar mass per 2,300 yr at $\eta = 0.1$ and exhausts an accessible reservoir of 10^5–$10^6\ M_\odot$ in $T_M \approx 2.3 \times 10^8$–$2.3 \times 10^9$ yr; this reservoir range is adopted for round numbers rather than derived, and sits 1.9–19× below Paper A's working cluster-mass reservoir ($1.9 \times 10^6\ M_\odot$ within 10 pc), a conservative (shorter-T_M) choice whose only downstream effect is the finite-horizon cap below. With finite horizons the ratio becomes

$$\frac{U_M}{U_S} \;=\; G\, p_s\, e^{-(\rho+\lambda)\tau}\, \frac{1 - e^{-(\rho+\lambda)T_M}}{1 - e^{-(\rho+\lambda)T_S}}. \tag{8}$$

Three consequences follow. At and near the threshold of Section 4, $(\rho+\lambda)T_M \gtrsim 4 \times 10^4$, so both brackets equal 1 to exponential accuracy and ρ^* is unchanged; the cancellation argument holds for all $\rho + \lambda \gtrsim$ a few $\times 10^{-8}$ yr^{-1} (a few $\times 10^{-9}$ yr^{-1} only at the long-T_M end of the range: at the short end, $T_M \approx 2.3 \times 10^8$ yr, $\rho + \lambda = 3 \times 10^{-9}$ yr^{-1} gives $(\rho+\lambda)T_M = 0.69$ and the bracket $1 - e^{-0.69} = 0.50$, not 1). In the patient limit $\rho + \lambda \to 0$ the bracket's value depends on which T_S horizon is operative. Against the unlifted horizon ($T_S \approx 5 \times 10^9$ yr, the solar-analogue fiducial), the bracket is $T_M/T_S \in [0.046, 0.46]$ (from $T_M \approx 2.3 \times 10^8$–$2.3 \times 10^9$ yr, l'Hôpital's rule on Eq. 8 at $\rho + \lambda \to 0$), capping the migration advantage at $\sim p_s G\, T_M/T_S \approx 2.3 \times 10^7$–$2.3 \times 10^8$. Against the star-lifted horizon for a fully convective low-mass host ($T_S \sim 10^{11}$–10^{12} yr, an illustrative alternative to the fiducial rather than a bound the solar-analogue case reaches; Section 3), the bracket shrinks and the cap instead falls to $\sim 1.15 \times 10^5$–1.15×10^7, rather than letting the advantage grow without bound in either branch. And the finite-horizon kernel is what makes the $\rho + \lambda \to 0$ Ramsey benchmark well-defined at all, since the infinite-horizon utilities (4)–(5) individually diverge there. One coupling worth stating: G and T_M anticorrelate at fixed reservoir, because the multiplier is set by the burn rate; the Bondi-fed $G = 10^6$ variant consumes $\sim 10^3$ times more slowly, giving $T_M \sim 10^{12}$ yr $> T_S$, so the patient-limit cap does not bind that branch.

A related asymmetry concerns the seed. Equation (6) treats seed arrival as all-or-nothing through p_s, but a seed can arrive functional with drifted goals, yielding computation the home lineage only partially values, and it carries its own transit-survival probability p_s^{seed} rather than the deliberating lineage's p_s (Table 1). A continuous treatment multiplies the hybrid's deferred term by a drift-weighted valuation $p_d \in [0, 1]$, the expected fraction of the destination payoff the deciding lineage credits after goal drift in the seed line, and substitutes p_s^{seed} for p_s in the deferred exponential:

$$U_H \;=\; (1 - f)\, U_S \;+\; p_d\, p_s^{\text{seed}}\, e^{-(\rho+\lambda)\tau}\, \frac{G\, C_S}{\rho + \lambda}. \tag{9}$$

Because a seed does not deliberate en route, its drift channels (radiation damage to stored values, replication error during bootstrap) differ from the institutional-drift channels of a deliberating lineage in transit, and




Figure 1: Undiscounted computation-rate histories for the three strategies at fiducial parameters (Table 1). Strategy S reaches its radiator-limited ceiling (Eq. 2) within centuries and holds it. Strategy M delivers nothing for $\tau = 10^5$ yr (its curve is drawn only for $t \geq \tau$; during transit there is no output), then a rate $G = 10^9$ times higher, conditional on transit survival. The hybrid follows S until the seed's destination comes online, then collects both; strictly its early segment is $(1 - f)\, C_{\mathrm{S}}$ plus the seed line, an offset invisible at this scale for $f \leq 10^{-6}$. The decision problem is whether the area under the M step, discounted and risk-weighted, exceeds the head start of S.

plausibly p_d is close to 1 by engineering; but nothing below requires that, since p_d enters the hybrid condition (Eq. 13) inside the same logarithm as $G p_s$. A deliberating lineage that migrates is exposed to the same institutional-drift channel over its own exploitation horizon; we denote the corresponding valuation p_d^{M}, so that $U_{\mathrm{M}}^{\mathrm{drift}} = p_d^{\mathrm{M}}\, U_{\mathrm{M}}$ (Table 1). Comparing Eq. (9) with $U_{\mathrm{M}}^{\mathrm{drift}}$ gives the general H-versus-M breakeven,

$$(1 - f)\, U_{\mathrm{S}} \;>\; \left(p_d^{\mathrm{M}}\, p_s \;-\; p_d\, p_s^{\mathrm{seed}}\right) \frac{G\, C_{\mathrm{S}}}{\rho + \lambda}\, e^{-(\rho+\lambda)\tau}, \tag{10}$$

which reduces to the one-line comparison of Section 5 when $p_s^{\mathrm{seed}} = p_s$ and collapses further to a sufficient condition $p_d \geq p_d^{\mathrm{M}}$ when the right-hand side is bounded above by zero. Equation (6) treats the two payoff streams as additive, which is exact if the lineage's utility is linear in computation and the seed's draw on local resources is the fraction f; Section 8 discusses when additivity fails (identity-centred utilities that do not value a forked descendant's computation).

Figure 1 shows the undiscounted rate histories: S delivers early and low, M late and high, H both.

4 Crossover conditions

4.1 Migrate versus stay

From Eqs. (4)–(5), $U_{\mathrm{M}} > U_{\mathrm{S}}$ reduces to

$$\boxed{\rho + \lambda \;<\; \frac{\ln(G\, p_s)}{\tau} \;\equiv\; \rho^*, \quad \text{valid for } G\, p_s > 1.} \tag{11}$$

Table 2: Sensitivity of the patience threshold to the transit-plus-construction time τ, at fixed $Gp_s = 5 \times 10^8$. The $\tau = 10^5$ yr row is the rounded fiducial (the bare cruise at $0.15c$ is 1.19×10^5 yr); the larger values represent deceleration by magsail or drag multiplying the cruise time by 2–3 (Zubrin and Andrews, 1991; Perakis and Hein, 2016; Heller and Hippke, 2017).

τ [yr]	$\rho^* = \ln(Gp_s)/\tau$ [yr^{-1}]	$t_{1/2} = \ln 2/\rho^*$ [yr]
1×10^5	2.0×10^{-4}	3,500
2×10^5	1.0×10^{-4}	6,900
3×10^5	6.7×10^{-5}	10,400

At $Gp_s \leq 1$ the logarithm is non-positive and Eq. (11) has no solution with $\rho + \lambda \geq 0$: migration never clears regardless of patience. The threshold depends on the enormous multiplier only through its logarithm: the structural reason the result is robust to the order-of-magnitude uncertainties in G. At the fiducials ($G = 10^9$, $p_s = 0.5$, $\tau = 10^5$ yr),

$$\rho^* = \frac{\ln(5 \times 10^8)}{10^5 \text{ yr}} \simeq 2.0 \times 10^{-4} \text{ yr}^{-1}, \tag{12}$$

corresponding to a combined discount-plus-hazard half-life $t_{1/2} = \ln 2/\rho^* \approx 3{,}500$ yr; this is a planning half-life in the usual sense only under $\lambda = 0$, since the threshold binds the sum $\rho + \lambda$, not the discount rate alone. Any lineage whose effective concern horizon exceeds a few millennia (a modest demand against the 10^8-yr coherence the family's architectures assume elsewhere) prefers migration to densification. Under the Paper-A-compliant realized-gain ceiling ($G = 6.6 \times 10^{15}$) the threshold rises to $\rho^* \approx 3.6 \times 10^{-4}$ yr^{-1} ($t_{1/2} \approx 1{,}900$ yr); under a pessimistic $p_s = 0.01$ it falls only to 1.6×10^{-4} yr^{-1}, and even $p_s = 10^{-4}$ moves it only to 1.15×10^{-4} yr^{-1}, a factor of 1.7; transit risk is nearly irrelevant inside the logarithm and the operative lever is the transit time τ. Both threshold and half-life additionally require $Gp_s > 1$: at $Gp_s \leq 1$ the logarithm is non-positive and no discount-plus-hazard rate clears the trade (Section 4 below returns to this floor).

Because $\rho^* \propto 1/\tau$ linearly, τ is the one parameter where a factor of a few moves the answer, and the deceleration allowance of Section 2.2 acts on τ directly. Table 2 propagates it: if braking multiplies the cruise time by 2–3, the threshold falls to $0.7\text{–}1.0 \times 10^{-4}$ yr^{-1} and the headline half-life lengthens to ∼7–10 kyr. The qualitative conclusion (patience horizons of millennia, against the 10^8-yr coherence assumed elsewhere in the family) is unchanged; the quoted 3,500-yr figure is the no-braking end of the band. These single-factor sensitivities are not independent: the joint pessimistic corner, ADIOS-suppressed $G = 10^3$ combined with braked $\tau = 3 \times 10^5$ yr, gives $\rho^* = \ln(500)/(3 \times 10^5) \approx 2.07 \times 10^{-5}$ yr^{-1} ($t_{1/2} \approx 33{,}500$ yr), a factor of ∼10 below the headline threshold and ∼3 below either single-factor pessimization alone. Worse, at $G = 10^3$ the tabulated low end $p_s = 10^{-4}$ (Section 4 above) gives $Gp_s = 0.1 < 1$: Eq. (11)'s validity condition fails, the logarithm is negative, and no discount-plus-hazard rate makes migration pay in that corner.

The $p_s = 0.5$ fiducial is anchored, loosely, in the only quantitative literature on transit attrition: Hoang et al. (2017) model gas and dust damage to relativistic spacecraft and find survivable but non-negligible erosion at $0.2c$ over the parsec-scale column to α Centauri. Over kiloparsec columns the damage integral is $\sim 4 \times 10^3$ times larger and mitigation (shielding, redundancy, en-route repair) is unmodelled, so 0.5 should be read as an order-unity placeholder rather than a conservative bound; the logarithmic dependence just exhibited is what makes this imprecision tolerable. Note that p_s prices transit alone; residence survival does not enter ρ^* (Eq. 11) and is reported here only as a check that the destination itself holds up once occupied. Paper E's flyby accounting with mass-segregated remnants carries two channels for the

operating envelope. The unbound stellar field is adiabatically decoupled from every orbit an installation could occupy (the adiabatic parameter runs from 2.2 at the stripping radius to 4×10^3 at the ISCO); the correction that would apply to it, an envelope minimum of 7.6×10^8 yr reaching the cluster-age cap in the interior, is retained by Paper E only as the measure of that decoupling and does not operate physically. The channel that does operate is bound-cusp diffusion: a bound 31 $M_\odot$ remnant is present at the envelope edge in 24.8 per cent of 2×10^4 Monte Carlo histories (Paper E's bound-remnant channel), giving a diffusion floor with median 4.3×10^7 yr (16th percentile 4.3×10^7 yr) in that branch, while 72.3 per cent of histories reach the 12-Gyr cluster-age cap because no bound remnant, or only a distant one, is drawn. Paper E's earlier fixed-number convention gave 53 per cent presence, and that figure is superseded. Residence survival to the 10^8-yr horizon is therefore bimodal, with 75 per cent of histories surviving to that horizon at the edge: in the minority branch the diffusion floor sits below the horizon by a factor of a few, so survival there is not assured by the flyby channel alone and would need the episodic-hazard and engineered-response accounting below rather than a single multiplicative p_s; the majority, cap-limited branch is survival for draws with an absent or distant bound remnant. Because p_s prices transit alone by definition (Eq. 11), this figure cannot enter ρ^* by construction, not because residence turns out not to matter: the insensitivity is a scoping fact about what the threshold measures, not a finding about residence durability. Section 4's threshold and half-life numbers are unaffected by the correction; what changes is only the confidence with which the residence phase, once reached, is described as durable. This paper does not attempt a residence-survival calculation folding the bimodal figures into a multiplicative correction on p_s or on the deferred payoff; that calculation is Paper E's appendix-dependent territory (episodic TDE hazard, engineered response) and is not carried here. The dominant recurrent hazard, tidal-disruption-event loading with recurrence $\gtrsim 10^7$ yr, is survivable by engineered response and enters the deferred payoff only as a small amortization correction on that horizon (Swanson, 2026a). The logarithmic entry of p_s just exhibited is what keeps the threshold stable against transit-side revisions; it says nothing about the bimodal residence question just raised.

Figure 2 maps ρ^* against destination distance for the beamed-sail speed range, marking named Galactic targets. Because $\tau \propto d$, the threshold scales as $1/d$: closer acceptable destinations relax the patience requirement proportionally, which is one economic reading of Paper A's selection pressure toward the nearest adequate system rather than the best one. The ω Cen marker sits at $\rho^* = 2.24 \times 10^{-4}$ yr^{-1}, slightly above the fiducial 2.0×10^{-4} yr^{-1} quoted above, because the figure uses the bare transit time at $v = 0.2c$ ($\tau = d/v \approx 8.95 \times 10^4$ yr) while the fiducial rounds τ up to 10^5 yr to include construction.

4.2 Meaning of the threshold

Equation (11) converts the inward-family debate into a single empirical-psychological question: do long-lived technological lineages discount the far future at more or less than a few parts in 10^4 per year? The literature's positions map onto the two sides. Bostrom (2003) argues that agents with astronomical stakes should approach $\rho = 0$; on his side of the line, migration dominates by many orders of magnitude. Steep discounters (lineages whose institutions cannot sustain plans beyond centuries) stay home and densify, and for them the inward family predicts nothing. The threshold also prices the goal-stability problem: Eq. (11) says the plan need only survive attrition at the few-per-10^4 annual rate during the transit window, not for the 10^8-yr exploitation phases that follow (see Eq. 7), and post-arrival value drift merely redistributes the payoff among successor agents while leaving the dynamical residues of Paper A in place. Under the symmetric valuation of Section 3 this transit-window reading holds whether λ prices lineage death or plan abandonment; the residual asymmetry is that a deliberating lineage in transit is exposed to institutional-drift channels a non-deliberating payload is not, one more consideration favouring seeds over whole-lineage relocation.




Figure 2: The patience threshold $\rho^* = \ln(Gp_s)/\tau$ (Eq. 11) versus destination distance, for the beamed-sail speed range and fiducial $Gp_s = 5 \times 10^8$. A lineage migrates when its combined discount-plus-hazard rate falls below the curve. Named points assume the marker's cluster is the nearest adequate destination for the deciding lineage; distances from Baumgardt and Vasiliev (2021) (47 Tuc, M54) and Häberle et al. (2025) (ω Cen, matching the Table 1 fiducial). The logarithmic dependence on Gp_s keeps the curves' movement modest: one decade in Gp_s moves ρ^* by ∼12 per cent, and the full adopted G bracket (10^3–6.6×10^{15}, ADIOS floor to Paper-A-compliant realized ceiling) moves it by a factor of ∼5.7. Both curves use the bare cruise time $\tau = d/v$; deceleration multiplying τ by 2–3 lowers them by the same factor (Table 2).

5 The hybrid result

5.1 Dominance

Comparing Eq. (6) with Eqs. (4) and (5): under additive utility and a full-fidelity seed ($p_d \geq p_d^{\mathrm{M}}$, $p_s^{\mathrm{seed}} = p_s$; Section 3), $U_{\mathrm{H}} > U_{\mathrm{M}}$ in one line (the stay component is pure addition), and $U_{\mathrm{H}} > U_{\mathrm{S}}$ whenever

$$f \;<\; G\, p_s\, e^{-(\rho+\lambda)\tau}, \quad \text{under additive utility and } p_d \geq p_d^{\mathrm{M}}. \tag{13}$$

The right-hand side at fiducials is $\sim 5 \times 10^8\, e^{-20\,(\rho+\lambda)/\rho^*}$, and the exponential is unforgiving: at the migration threshold itself the tolerable seed cost is $5 \times 10^8 e^{-20} \sim 1$, but at $\rho + \lambda = 2\rho^*$ it has fallen to $\sim 2 \times 10^{-9}$, and at $10\rho^*$ to $\sim 10^{-78}$. The physical seed cost, gram-to-kilogram payloads plus a launch array (Lubin, 2016; Freitas, 1980), is tiny against a Dyson-scale economy: Armstrong and Sandberg (2013) estimate that seeding every galaxy in the reachable universe costs hours of one star's output, which supports $f \lesssim 10^{-6}$ a fortiori for a single Galactic destination. Setting $f = 10^{-6}$ in Eq. (13) gives breakeven at $(\rho + \lambda)\tau = \ln(5 \times 10^{14}) \approx 34$, i.e. $\rho + \lambda \approx 1.7\,\rho^*$. The hybrid therefore dominates both pure strategies in a band extending modestly beyond the migration threshold, roughly 1.7× the threshold at $f = 10^{-6}$ (and only logarithmically further for still-cheaper seeds), because the exponential discount on the deferred payoff crushes the seed's expected value within a factor of a few of ρ^*. Outside that band the utility calculus reverts to pure S; inside it, and everywhere migration itself clears, H is strictly best barring non-additive utilities (Section 8). Figure 3 maps the three regimes in the $(G,\ \rho + \lambda)$ plane.

The comparison with M is one line only under full seed fidelity and $p_s^{\mathrm{seed}} = p_s$. In that case Eq. (10) reduces to $(1 - f)\, U_{\mathrm{S}} > (1 - p_d)\, U_{\mathrm{M}}$ at $p_d^{\mathrm{M}} = 1$ (a deliberating migrant credited at full fidelity); deep in migration-favourable territory ($U_{\mathrm{M}}/U_{\mathrm{S}} \approx 2.2 \times 10^4$ at $\rho + \lambda = 0.5\,\rho^*$) that requires $1 - p_d \lesssim 4 \times 10^{-5}$ (more closely 4.47×10^{-5}). In general the tolerance on the seed's fidelity deficit $d = p_d^{\mathrm{M}} - p_d$ is $d < (1 - f)(G p_s)^{x-1}$ with $x = (\rho + \lambda)/\rho^*$, because $U_{\mathrm{M}}/U_{\mathrm{S}} = (G p_s)^{1-x}$: it is 2×10^{-9} at $\rho + \lambda \to 0$, 3×10^{-7} at $x = 0.25$, 4.5×10^{-5} at $x = 0.5$, 6.7×10^{-3} at $x = 0.75$, unity at the threshold, and unconstrained above it, so the 4×10^{-5} is the value at half the threshold, not a bound across the band. The rescuing assumption, stated here because the dominance claim rests on it: a migrating lineage carries its own drift factor p_d^{M} over the same horizon, and a non-deliberating seed drifts no more than a deliberating lineage in transit ($p_d \geq p_d^{\mathrm{M}}$), whence Eq. (10)'s right-hand side is bounded above by zero and $U_{\mathrm{H}} \geq U_{\mathrm{M}}$ term by term. Given that assumption, and given $p_s^{\mathrm{seed}} = p_s$ (Table 1: adopted, not derived, since seed transit hazards are unmodelled separately), the dominance itself is close to a bookkeeping identity; the substantive content of this section is the narrowness of the band beyond the threshold, $1.7\,\rho^*$ at $f = 10^{-6}$, given a nearly free seed. A seed with materially lower survival than the deliberating lineage ($p_s^{\mathrm{seed}} < p_s$) would need a correspondingly larger p_d margin to compensate, per Eq. (10); this paper does not attempt that separate estimate.

5.2 Consequences

The dominance of H restructures the Fermi implications of the family in three ways.

First, compliance is cheap for the patient: the incomplete-compliance objection (Paper B, problem P4) imagines lineages weighing a costly relocation and many declining. Under H there is no relocation to decline: dispatching seeds is compatible with staying, and the marginal cost is negligible against Dyson-scale output. For every lineage inside the dominance band ($\rho + \lambda \lesssim 1.7\,\rho^*$), the question "what fraction migrates?" becomes "what fraction of computation-valuing lineages fails to spend $f \sim 10^{-6}$ of output on a large expected return?" The compliance question is thereby displaced, not dissolved: it




Figure 3: The phase space of the migration decision in the $(G,\ \rho + \lambda)$ plane, at $p_s = 0.5$ and $\tau = 10^5$ yr (Table 1). Green: migration clears, $\rho + \lambda < \rho^* = \ln(G p_s)/\tau$ (Eq. 11); within it the hybrid weakly dominates pure M under the seed-fidelity assumption. Orange: the hybrid band, where the hybrid beats S while pure M does not, extending to $\rho + \lambda = \ln(G p_s/f)/\tau = 1.7\,\rho^*$ at the fiducial seed cost $f = 10^{-6}$ (Eq. 13). Blue: S dominates; neither migration nor seeding pays. The star marks the fiducial point ($G = 10^9$, $\rho + \lambda = 2 \times 10^{-4}$ yr^{-1}), which sits on the migration threshold. The plotted G axis does not extend to the Paper-A-compliant realized-gain ceiling ($G \approx 6.6 \times 10^{15}$, Section 3); that ceiling lies further right along the same green region and moves ρ^* by the factor of ~1.8 quoted in Section 4.

becomes the question of what fraction of maximizers sit below the modestly enlarged threshold, which is a claim about the distribution of $\rho + \lambda$ rather than about willingness to pay.

Second, destinations saturate early: if seeds are cheap, the nearest adequate IMBH systems receive seeds from every computation-valuing lineage in their neighbourhood, and first arrival plausibly matters. This turns the uncontested-claim criterion of Paper A (criterion 3) into a race dynamic of the kind Hanson (1998) analysed for interstellar colonization generally, in which selection favours the fastest and least encumbered colonizers of contested oases, and it implies that the residues of Paper A's P4 should be present in suitable systems even if no lineage ever "migrates" in the whole-population sense. One scope note: the threshold of Eq. (11) is a single-agent calculation, and competition for contested oases only lowers it, since preemption adds option value to early arrival; Eq. (11) is therefore an upper bound on the patience required.

The apparent conflict with Paper A's uncontested-claim criterion resolves by economic era rather than by contradiction: the destination is uncontested by fusion-era economics, for which low-mass cluster stars are poor assets, and maximally contested among computation-maximizing lineages, for which they are the prize; this is the form in which Paper A now states criterion 3, and it is the resolution both papers adopt. Congestion also enters the threshold only logarithmically. If k lineages arrive and share the destination, the per-lineage gain is G/k and the threshold shifts by $\ln k$ inside Eq. (11); if the destination is winner-take-all, the survival factor becomes p_s/k, with the same logarithmic entry. For $k \lesssim 10^2$ the shift is smaller than the uncertainties already carried in G, so the single-agent threshold stands.

Third, the loud residue does not vanish. Throughout this paper, loud denotes an expansion-front signature in the grabby-aliens sense (Section 6); localized waste heat from a stationary Matrioshka-class collector is a distinct residue class, treated on its own terms in Section 7. Under pure M, migrated lineages leave planetary space entirely and the sky is silent. Under dominant H, the stay component continues local densification, whose Matrioshka waste heat is the classic detectable technosignature (Dyson, 1960; Wright et al., 2014). The hybrid therefore predicts both residue classes in linked proportion, the basis of the new population statistic in Section 7.

6 Population consequences

Let a fraction x of long-lived technological lineages be expansionists, the class supplying the loud signals whose absence constitutes the paradox (Hanson et al., 2021; Olson, 2015); the remainder are satisficers and other quiet types that contribute nothing to N_{loud} regardless of the inward gradient. Expansionism and computation maximization are not exclusive: a lineage can expand and also value integrated computation, and it is that overlap the inward family acts on. Within the expansionist class, let q be the fraction susceptible to the inward gradient, that is, the conditional probability that an expansionist also holds the P1 optimization goal, so that the gradient bears on its behaviour at all. Within that susceptible subset, let m be the fraction whose parameters fall below the (hybrid-enlarged) threshold of Section 5, and $w \in [0, 1]$ the capture weight: the probability that susceptibility plus a cleared threshold actually converts loud expansion into quiet inward behaviour before or during the expansion phase.

The inward family's contribution to the silence is the claim, argued qualitatively in Paper A, that the maximizer subset contains the oldest and most capable lineages, so its withdrawal from loud behaviour thins the expected loud population disproportionately. The expected number of loud civilizations in a survey volume containing N lineages is

$$N_{\text{loud}} \;=\; N\big[\, x\,(1 - w\,q\,m)\,\big], \tag{14}$$

with q, m, and w the nested conditional fractions defined above, so that wqm is the capture probability

for a randomly drawn expansionist. Equation (14) identifies only the product wqm: a measured $N_{\rm loud}$ constrains one number, and the (q, w) grid below is a re-parameterization of that single degree of freedom, useful for exhibiting which value combinations reach it, not evidence of two independently constrained parameters.

One large step separates $N_{\rm loud}$ from anything observed. $N_{\rm loud}$ counts loud civilizations that *exist*; the surveys invoked in Section 7 bound the number that would have been *detected*, so the silence likelihood is $P(0\ \text{detected}) = \exp(-N_{\rm loud}\langle p_{\rm det}\rangle)$, where $N_{\rm loud}\langle p_{\rm det}\rangle = 1$ is a 37 per cent chance of the observed silence and the 95 per cent upper limit, the value above which silence is excluded, is $N_{\rm loud}\langle p_{\rm det}\rangle < 3.0$ (the condition $N_{\rm loud}\langle p_{\rm det}\rangle < 1$ used below is therefore conservative), with $\langle p_{\rm det}\rangle$ the completeness of those surveys for the signature in question. For Galactic Matrioshka-class sources that completeness is well below unity and is distance- and temperature-limited in the ways Section 7 describes (Carrigan, 2009; Zackrisson et al., 2018). Table 3 is therefore evaluated at $\langle p_{\rm det}\rangle = 1$ and its bold cells are the most restrictive corner of the condition; any realistic completeness widens the silence-compatible region, in some cases substantially. We tabulate the $\langle p_{\rm det}\rangle = 1$ case because it is the one that does not require a survey-completeness model this paper has no means to supply, and we flag the direction of the bias rather than absorb it. For the potential-expansionist count Nx, the dissolution analysis of Sandberg et al. (2018) shows that reasonable uncertainty over the Drake factors puts substantial probability on Nx of order unity as well as on large values, which is why we tabulate three decades of it. The three rows are scenarios, not equally weighted hypotheses: that analysis produces a posterior on N given the silence, and it places its largest mass on the low branch, where no inward capture is needed to dissolve the paradox at all. We tabulate the high branches because the family's contribution is only interesting there, not because they carry equal weight. Table 3 evaluates the conservative silence condition $N_{\rm loud} < 1$ across the (q, w) grid for three values of the potential-expansionist count Nx. The table's message matches what Papers A and B anticipated, with the dependence on Nx making the thinning claim quantitative rather than binary. At the fiducial $Nx = 10$, inward capture alone dissolves the paradox only in the complete-capture corner ($wq \to 1$); at moderate capture it thins the expectation by factors of a few and the balance must come from conventional rarity. At $Nx = 100$ even complete capture of the maximizer subset leaves the paradox intact unless $q \to 1$, so the family there functions only as a rarity multiplier. At $Nx = 3$ the silence condition opens up: capture products $wq \gtrsim 0.7$ suffice, and several non-degenerate cells satisfy $N_{\rm loud} < 1$ without full compliance. Hybrid dominance no longer drives $m \to 1$ across the board: Section 5 enlarges the acting region only to $\rho + \lambda \lesssim 1.7\,\rho^*$, so m is the fraction of susceptible expansionists below that modestly enlarged threshold and remains a live economic parameter alongside q and w. The residual uncertainty therefore splits between the distribution of patience (m) and the distribution of terminal values (q, w), with the economics of this paper constraining only the first.

7 Observational corollaries: the residue ratio

Epistemic status: This section converts the decision analysis into population-level observables. It inherits all conditionality of the optimization premise.

Hybrid dominance predicts linked residues: for every lineage acting on the gradient, a continuing local densification (Matrioshka-class mid-infrared source) and a seeded quiet-cluster system (the Paper A dynamical residues: anomalous spin, suppressed accretion, core depletion). Pure M predicts quiet clusters

Table 3: Expected loud count $N_{\rm loud}$ (Eq. 14) for three values of the potential-expansionist count Nx per survey volume ($Nx = 10$ fiducial), with q the fraction of expansionists susceptible to the inward gradient, w the capture weight, and $m = 1$ throughout (entries are lower bounds on $N_{\rm loud}$ when part of the population discounts above $\sim 1.7\,\rho^*$). Bold: $N_{\rm loud} < 1$, where silence has probability > 0.37; the 95 per cent upper limit is 3.0. Only the single product wqm enters Eq. (14); the grid re-parameterizes that one degree of freedom to exhibit which combinations reach silence.

		$w = 0.25$	$w = 0.5$	$w = 0.75$	$w = 1$
$Nx = 3$	$q = 0.25$	2.8	2.6	2.4	2.3
	$q = 0.5$	2.6	2.3	1.9	1.5
	$q = 0.75$	2.4	1.9	1.3	**0.8**
	$q = 1$	2.3	1.5	**0.8**	**0.0**
$Nx = 10$	$q = 0.25$	9.4	8.8	8.1	7.5
	$q = 0.5$	8.8	7.5	6.3	5.0
	$q = 0.75$	8.1	6.3	4.4	2.5
	$q = 1$	7.5	5.0	2.5	**0.0**
$Nx = 100$	$q = 0.25$	94	88	81	75
	$q = 0.5$	88	75	63	50
	$q = 0.75$	81	63	44	25
	$q = 1$	75	50	25	**0.0**

without Matrioshka sources; pure S the reverse. The observable is the sky ratio

$$R_{\rm sky} \equiv \frac{N_{\rm Matrioshka}}{N_{\text{quiet-cluster residue}}} = \begin{cases} 0 & \text{pure M} \\ O(1) \times \dfrac{n_{\rm lineage}}{n_{\rm target}} & \text{hybrid H} \\ \infty & \text{pure S,} \end{cases} \tag{15}$$

where $n_{\rm lineage}/n_{\rm target}$ is the ratio of acting lineages to adequate destination systems in the relevant volume (several lineages can seed one cluster; one lineage densifies one home system). The Milky Way contains $\sim$150 globular clusters of which a handful satisfy the Paper A criteria, so under H with even a few Galactic lineages the formation-rate ratio satisfies $R \gtrsim 1$.

The observed ratio is a snapshot, however, and the two residue classes have very different lifetimes. Paper E's cascade analysis finds that an abandoned Matrioshka-class swarm collides itself into debris on 10^2–10^3 yr and (at the destination) drains into the hole, so dead installations vanish quickly and only actively maintained ones radiate; the quiet-cluster dynamical residues (anomalous spin above all, the one fossil that outlasts the Hubble time) persist for $\gtrsim$ Gyr (Swanson, 2026a). The observable sky ratio is therefore the formation-rate ratio multiplied by a lifetime ratio, $R_{\rm obs} = R \times (t_{\rm Matrioshka}/t_{\rm residue})$, and if home installations are maintained only while the lineage remains interested, $t_{\rm Matrioshka}/t_{\rm residue}$ can be $\ll 1$, driving $R_{\rm obs} \ll 1$ even under fully compliant H.

This puts existing null results to work, with the lifetime factor carried explicitly. For extragalactic waste heat, the WISE-based $\hat{G}$ surveys (Wright et al., 2014; Griffith et al., 2015) bound galaxy-scale reprocessed emission in $\sim 10^5$ external galaxies; for Galactic Dyson- and Matrioshka-class sources, the operative bounds are the IRAS search of Carrigan (2009), the Gaia/2MASS-selected candidate analysis of Zackrisson et al. (2018), and the Project Hephaistos series: Suazo et al. (2024) report seven unresolved Galactic candidates surviving their photometric cuts, most or all of which subsequent analyses attribute to hot-dust-obscured background AGN contamination (Ren et al., 2024; Blain, 2024; Ren et al., 2025). Together these bound $N_{\rm Matrioshka}$ in the Galaxy at or near zero over their completeness ranges, pending contamination analysis

of the surviving Hephaistos candidates. Through Eq. (15), those bounds propagate to the hybrid branch only after division by the lifetime ratio, and the propagation is a limit rather than a deduction. With $N_{\text{Matrioshka}}$ observed as zero, the ratio $N_{\text{Matrioshka}}/N_{\text{quiet}}$ carries no information about its own denominator; what the data support is a joint Poisson likelihood on the two counts with R_{obs} as a parameter, from which an upper limit on N_{quiet} follows only through an assumed R_{obs} and its uncertainty: with zero observed Matrioshka sources the 95 per cent one-sided Poisson limit is $N_{\text{quiet}} < 3.0/(R_{\text{obs}}\langle p_{\text{det}}\rangle)$, so the constraint is $N_{\text{quiet}} < 3$ at $R_{\text{obs}} = 1$ and weakens as $1/R_{\text{obs}}$. Stated that way: if $R_{\text{sky,obs}} \gtrsim 1$ (long-maintained home installations), the infrared nulls place a comparable upper limit on $N_{\text{quiet-cluster residue}}$, and the family's Galactic prediction concentrates on the pure-M corner: lineages whose utilities are non-additive across descendants (Section 8) or whose home infrastructure is dismantled at departure. If instead maintenance lifetimes are short, the infrared nulls carry little weight against H and the quiet-cluster channel is left unconstrained by them. The Paper C campaign tests the quiet-cluster channel directly at ω Cen either way. Conversely, a confirmed Paper A residue (a near-extremal spin on a gas-starved IMBH) with a continuing Matrioshka null would be evidence about the utility structure of whatever laid the residue down: it would indicate migration without a maintained home presence, the signature of either whole-lineage relocation or post-departure home abandonment. That a spin measurement could carry information about value structure is an inference this pricing framework licenses; we state it in advance of any data.

The archival anomaly-detection agenda of the Dyson Minds workshop (Curtis et al., 2026) and the cluster-ranking metrics of Huang et al. (2026) both operate on the two source classes Eq. (15) links; a joint analysis (ranking clusters and IR-anomaly candidates in the same volume and testing the correlation Eq. (15) implies) is executable from public archives and is, to our knowledge, unproposed.

8 Limits and objections

Model dependence of the utility. Equation (1) assumes utility linear in computation rate and additive across locations and descendants. Identity-centred utilities break additivity: a lineage that values only its own continuous thread of computation may not credit a forked seed's output, restoring the pure S-versus-M choice of Section 4, which is why the crossover analysis is retained rather than replaced by the hybrid result. Satisficing utilities (compute enough, not most) exit the framework entirely; they belong to the sustainability solution (Haqq-Misra and Baum, 2009), not to the inward family.

Discounting form. Exponential discounting is the tractable choice, and no directional claim survives its relaxation cleanly. Hyperbolic discounters (Laibson, 1997) have low long-run rates, but they are time-inconsistent: the deciding present self applies its steep short-run rate to the whole deferred payoff at the moment of choice, so hyperbolic discounting does not automatically favour migration. The outcome depends on sophistication: a naive hyperbolic agent perpetually postpones a launch it always "intends" to make, while a sophisticated one may commit early (a seed, once launched, is a commitment device immune to later preference reversals, one more argument for H over M). A genuinely zero-discount agent (Ramsey, 1928; Bostrom, 2003) is handled by the finite-horizon kernel of Section 3: the infinite-horizon utilities diverge at $\rho + \lambda = 0$, but the ratio of Eq. (8) remains defined and migration wins by the capped factor $p_s G\, T_{\text{M}}/T_{\text{S}}$: $\sim 2.3 \times 10^7$–2.3×10^8 against the unlifted, solar-analogue T_{S}, or $\sim 1.15 \times 10^5$–1.15×10^7 against the star-lifted horizon for a fully convective low-mass host (Section 3).

Uncertainty about the discount rate pushes the same direction, and more forcefully. Under gamma (declining) discounting (Weitzman, 1998, 2001), an agent uncertain over its own ρ values the far future by

the expectation of the discount factor, giving a certainty-equivalent rate

$$\rho_{\text{eff}}(t) \;=\; -\frac{1}{t}\,\ln \mathbb{E}\big[e^{-\rho t}\big]\,, \tag{16}$$

which declines with horizon toward the minimum plausible rate: the patient scenarios dominate the expectation at long t. An agent placing equal weight on $\rho = 10^{-3}$ and $\rho = 10^{-6}$ has $\rho_{\text{eff}}(\tau) = -\ln[\tfrac{1}{2}(e^{-100} + e^{-0.1})]/\tau \approx 7.9 \times 10^{-6}$ yr^{-1} at $\tau = 10^5$ yr (the certainty-equivalent rate approaches the 10^{-6} floor only at $t \gtrsim 10^7$ yr), still a factor of 25 below the threshold of Eq. (11). Migration therefore requires only uncertainty about patience, not patience itself, which strengthens the case beyond the exponential baseline. For calibration in the other direction, empirical institutional discount rates cluster at 10^{-2}–10^{-3} yr^{-1} (Weitzman's survey of professional economists' rate judgements spans this range and declines toward its lower end for far-future horizons; Weitzman 2001), failing the threshold by up to two orders of magnitude: the framework describes old lineages, not present-day institutions.

Sail speed and the waiting alternative. Heller (2017) showed that a civilization expecting propulsion to improve faces an incentive trap: launching now can be dominated by waiting for a faster vehicle that overtakes the earlier one. In this framework the trap does not bind. Waiting costs discounted utility at the rate $\rho + \lambda$ on the entire deferred payoff, while speed improvements enter only through $\tau \propto 1/v$ inside the same energy economics that already caps beamed sails at 0.1–0.2c; a bounded reduction in τ cannot repay an open-ended delay, and the seed option makes the launch cost too small for the saving to matter.

Real options. The migration decision is the canonical problem of irreversible investment under uncertainty (McDonald and Siegel, 1986; Dixit and Pindyck, 1994), and the wait-for-sails argument above is an informal option-value comparison. In the deterministic setup of this paper the option to defer is worthless: with v, G, and τ known, waiting buys no information, only discounting, which is the conclusion of the preceding paragraph restated in that literature's terms. The seed strategy then has a standard name: it buys the option cheaply rather than exercising it, holding the deferred claim at cost f while the state of the world reveals itself. A full treatment requires stochastic dynamics for v or G (a Poisson-arrival improvement process is the natural minimal model) and is deferred.

The hazard rate and goal stability. Folding P2 into λ prices the problem without solving it; the price is the problem's size: under either reading of λ given the symmetric valuation of Section 3, coherence is required at the few-per-10^4-per-year level during transit only (Section 4), a far weaker demand than the 10^8-yr coherence the exploitation architecture assumes, and one that seed automation (a payload that does not deliberate en route) plausibly meets by construction; the residual seed-side drift is carried continuously by the valuation factor p_d of Section 3 rather than as an all-or-nothing loss. The deep version of P2 (whether the destination civilization retains the values that sent it) remains open and remains, as Paper B noted, not an astronomy problem.

Parameter unmeasurability. q, w, ρ, λ are distributions over agents we have never observed. The defensible claims are structural: the logarithmic insensitivity of the threshold (Eq. 11), the dominance region of the hybrid (Eq. 13), and the sign of the residue-ratio prediction (Eq. 15). Point estimates in this paper are fiducials for orientation, not forecasts.

Grabby-aliens integration. The Hanson et al. (2021) machinery conditions on our early arrival to bound the density of loud expanders. Equation (14) plugs into that frame by modifying the loud fraction while leaving the arrival-time statistics unchanged: inward capture reduces the volume-conversion rate of loud lineages without changing when civilizations arise. A full integration (re-deriving the grabby-aliens posterior with a capture term) is beyond this paper's scope and is the natural next step for whoever takes up Paper B's problem P1; Eq. (14) supplies the term to insert.

Outward counter-positions. Ivliev (2026) (preprint, submitted to *Acta Astronautica*) argues from a comparable automation premise to the opposite directional conclusion: once autonomous AI industry crosses a capability threshold, quiet outward expansion through low-mass probes and seed factories becomes too cheap for any civilization to refuse, and the silence reflects expansion signatures too weak to detect rather than migration. The two frameworks agree that seeds are nearly free (his threshold argument and our Eq. 13 are the same observation) and disagree about the objective: under our kernels the same cheap automation buys more computation per unit resource by seeding inward than by saturating stellar systems, so the disagreement reduces to whether coverage or computation is the maximand, i.e. to the distribution of terminal values already carried by q in Section 6.

Two further outward positions deserve engagement. Hooper (2018) argues that a civilization anticipating accelerating cosmic expansion should harvest stars from surrounding volumes and bring them home before the horizon removes them, the strongest quantitative argument for outward action in the literature. The mechanism concedes this paper's core premise, mass relocation across kiloparsec scales in the service of future computation, and disputes only the direction; and its clock is the dark-energy horizon at $\sim 10^{11}$ yr, which never binds the $\tau = 10^5$ yr decision analysed here: a lineage can seed inward now and harvest outward later without conflict between the two programmes. Ord (2021) maps the affectable universe and reaches a structurally similar conclusion, that the resources within outward reach are bounded and shrinking, which if anything sharpens the case for optimizing what a lineage can already touch.

Anthropic caveats. As with Paper A, our own existence as a young loud lineage neither confirms nor embarrasses the analysis; the framework describes the asymptotic behaviour of old lineages, and we are not one. Self-indication corrections to the population frame of Section 6 are absorbed into the grabby-aliens integration deferred above.

9 Conclusion

Paper B posed the question this paper answers: under what conditions does relocation to a thermodynamically privileged destination beat staying home and building? The answer has three parts, each built from stipulated inputs (destination multiplier G, transit survival p_s, seed cost f) rather than measured ones; what the analysis derives is their combination, not their values. The crossover between migration and densification is a patience threshold, $\rho + \lambda < \ln(Gp_s)/\tau$ (valid for $Gp_s > 1$), whose logarithmic form makes it robust to the large uncertainties in the destination multiplier and transit risk; at fiducial, stipulated parameters it asks only for combined discount-plus-hazard half-lives beyond a few thousand years. Given a nearly free, stipulated seed cost and near-unity seed fidelity, the seed-and-stay hybrid dominates both pure strategies throughout the migration-favourable region; the result this paper derives, rather than stipulates, is the band's narrowness, extending only to roughly $1.7\times$ the threshold at $f = 10^{-6}$ before the exponential discount on the deferred payoff closes the window. Within that band the incomplete-compliance problem reduces to the joint distribution of patience and the optimization goal. And the hybrid's linked residues yield a population-level prediction, a formation-rate ratio of Matrioshka sources to engineered clusters of

order unity, for which existing infrared nulls supply a joint-likelihood limit only after weighting by the maintenance lifetime of home installations, concentrating the family's surviving parameter space on the pure-migration and short-maintenance corners that the ω Cen campaign of Paper C tests directly.

The broader point is methodological. The inward family has argued from thermodynamic gradients (Paper A), been organized and graded as a literature (Paper B), and acquired a costed observational program (Paper C). What it lacked was a demonstration that the gradient is economically actionable by agents with finite patience and fallible institutions, the step where favourable gradients often fail. This paper supplies that demonstration in closed form, with its assumptions priced and its observational consequences stated in advance. Whether any agent has ever faced the decision is a question for the instruments.

Acknowledgements and disclosure

The author thanks the maintainers of the NASA Astrophysics Data System and arXiv, on which the citation verification for this work relied. The author's conflict of interest as proponent of the hypothesis whose economics this paper analyses is noted; the mitigation is the same as in the companion papers: closed-form results, conservative fiducials, and pre-stated observational consequences. **AI assistance disclosure:** drafting, citation verification, derivation checking, and figure preparation for this manuscript were performed with substantial assistance from a large language model (Claude, Anthropic), under the author's direction; the author reviewed and takes full responsibility for all claims, derivations, and references. Interactive calculators implementing the quantitative material in Sections 3–7 are available at https://omegacentauri.me.

Data availability

No new observational data were generated for this work. All quantitative inputs derive from the cited literature.

Revision record

v2.7 (2026-10-24). Fact fixes from the R10 hostile-referee round, adjudicated by the farm (D300, D305) and independently checked. (1) The seed-fidelity condition of the hybrid is stated with its tolerance, $(1-f)(Gp_s)^{x-1}$ at $x = (\rho+\lambda)/\rho^*$: 4.5×10^{-5} at $x = 0.5$ (the earlier 4×10^{-5}), unity at the threshold, 2×10^{-9} as $\rho+\lambda$ goes to zero. The abstract's "near-unity" is replaced by this statement. (2) The silence condition of Section 6 is a likelihood, $P(0) = \exp(-N_{\rm loud}\langle p_{\rm det}\rangle)$: $N_{\rm loud}\langle p_{\rm det}\rangle < 1$ corresponds to a 37 per cent chance of the observed silence, and silence excludes $N_{\rm loud}\langle p_{\rm det}\rangle$ above 3.0 at 95 per cent ($e^{-3} = 0.050$). The farm's earlier statement that 95 per cent compatibility needs $N_{\rm loud} < 0.05$ inverted this inference and is not carried. The bold cells of the table keep the conservative threshold of 1. (3) The archival-null limit of Section 7 carries its Poisson form, $N_{\rm quiet} < 3.0/(R_{\rm obs}\langle p_{\rm det}\rangle)$, and the abstract says the nulls supply the frame for a conditional limit. (4) The seed-cost entry states that f is a fraction of the discounted payoff, a permanent tax. (5) The transit time is 10^5–10^6 years at the two places that gave 10^4–10^6 (the bare cruise at $0.15c$ is 1.19×10^5 yr); the $\tau = 10^5$ row is labelled the rounded fiducial. (6) The $p_s = 0.5$ entry is labelled a placeholder, the destination definition names criterion 3, and the first-pricing claims read "we are not aware of a prior pricing". (7) The Paper A gain condition quoted in the sink discussion is the vacuum state ($N_{\rm eff} \ll 1$), with the thermal case a factor 20 deficit (Paper A v2.14); the v2.5 entry below states it in the superseded wording.

v2.6 (2026-10-23). Fact fixes from the farm audit D269. The ADIOS-floor threshold of Section 3 is stated with its inputs: $G \approx 2.6 \times 10^3$ gives $\rho^* = \ln(1.3 \times 10^3)/10^5 = 0.72 \times 10^{-4}$ yr^{-1}, and the printed 0.6×10^{-4} belongs to the rounded $G = 10^3$ ($\ln 500/10^5 = 0.62 \times 10^{-4}$). The accessible reservoir of 10^5–10^6 $M_\odot$ sits 1.9–19× below Paper A's 1.9×10^6 $M_\odot$ within 10 pc (was 4–40×, which divided by the 4.3×10^6 stars rather than the mass).

v2.5 (2026-10-22). The realized entropy-disposal gain quoted from Paper A (§4.2) is corrected from 10^6–10^9 to about 10^3 (fed flow) and 10^7 (ambient medium), conditional on carrier modes colder than the 2.7 K sky and of order unity otherwise, following Paper A v2.12; the realized band of §3 is correspondingly 6.6×10^{11}–6.6×10^{15} under that condition. The remaining derived values are recomputed from $\rho^* = \ln(G p_s)/\tau$ at $p_s = 0.5$, $\tau = 10^5$ yr: at the ceiling $G = 6.6 \times 10^{15}$ the threshold is 3.6×10^{-4} yr^{-1} ($t_{1/2} \approx 1{,}900$ yr, was 4.0×10^{-4} and 1,700 at 6.6×10^{17}), the factor over the fiducial is ≤ 1.8 (was ≤ 2.0), and the factor across the full adopted bracket in the caption of Figure 2 is ~5.7 (was ~6.5). The fiducial $G = 10^9$ of Strategy M is the Eddington power ratio and is unchanged.

v2.4 (2026-10-22). Citation audit. The Bennett et al. (2019) argument is restated as an erasure-cost result (waiting does not lower erasure cost), in the abstract and in the gap paragraph. The transit-speed range now cites Hoang et al. (2017) and notes that Lubin (2016) give $> c/4$ for gram-scale probes. The 16th-percentile bound-cusp floor is 4.3×10^7 yr (was 2.2×10^7, the superseded fixed-number convention), the unbound envelope minimum is 7.6×10^8 yr (was 3×10^8), the ambient Bondi ceiling is $\sim 8 \times 10^5$ $L_\odot$ (was $\sim 10^6$), and the Hooper (2018) horizon clock is $\sim 10^{11}$ yr (was 10^9–10^{10}). The realized gain is described as a per-erasure cost ratio rather than a rate ratio. The Laibson (1997) page range is 443–478.

References

Armstrong, S. and Sandberg, A. (2013). Eternity in six hours: Intergalactic spreading of intelligent life and sharpening the Fermi paradox. *Acta Astronautica*, 89:1–13.

Badescu, V. and Cathcart, R. B. (2000). Stellar engines for Kardashev's type II civilisations. *Journal of the British Interplanetary Society*, 53:297–306.

Baumgardt, H. and Vasiliev, E. (2021). Accurate distances to Galactic globular clusters through a combination of Gaia EDR3, HST, and literature data. *Monthly Notices of the Royal Astronomical Society*, 505:5957–5977.

Bennett, C. H. (1973). Logical reversibility of computation. *IBM Journal of Research and Development*, 17:525–532.

Bennett, C. H., Hanson, R., and Riedel, C. J. (2019). Comment on 'The aestivation hypothesis for resolving Fermi's paradox'. *Foundations of Physics*, 49:820–829.

Blain, A. W. (2024). Did WISE detect Dyson spheres/structures around Gaia-2MASS-selected stars? Submitted to MNRAS.

Bostrom, N. (2003). Astronomical waste: the opportunity cost of delayed technological development. *Utilitas*, 15:308–314.

Bradbury, R. J. (1999). Matrioshka brains. Unpublished working manuscript, revised through 2001. Archived at https://gwern.net/doc/ai/scaling/hardware/1999-bradbury-matrioshkabrains.pdf; mirrored at https://web.archive.org/web/2024/https://gwern.net/doc/ai/scaling/hardware/1999-bradbury-matrioshkabrains.pdf.

Carrigan, Jr., R. A. (2009). IRAS-based whole-sky upper limit on Dyson spheres. *The Astrophysical Journal*, 698:2075–2086.

Curtis, O., Adams, V. H., Angerhausen, D., Bates, J., Berea, A., Dick, S. J., Elvis, M., Khatri, S. P., Linares, R., Muco, M., Seager, S., and Wright, J. T. (2026). The Dyson Minds 2025 workshop: SETI around black holes. *Publications of the Astronomical Society of the Pacific*, 138:046001.

Dixit, A. K. and Pindyck, R. S. (1994). *Investment under Uncertainty*. Princeton University Press, Princeton.

Dyson, F. J. (1960). Search for artificial stellar sources of infrared radiation. *Science*, 131(3414):1667–1668.

Freitas, Jr., R. A. (1980). A self-reproducing interstellar probe. *Journal of the British Interplanetary Society*, 33:251–264.

Garrett, M. (2026). Slysh haloes: the waste heat of cold computing as a submillimetre technosignature.

Griffith, R. L., Wright, J. T., Maldonado, J., Povich, M. S., Sigurdsson, S., and Mullan, B. (2015). The $\hat{\mathrm{g}}$ infrared search for extraterrestrial civilizations with large energy supplies. III. The reddest extended sources in WISE. *The Astrophysical Journal Supplement Series*, 217(2):25.

Häberle, M., Neumayer, N., Clontz, C., Seth, A. C., Bellini, A., et al. (2025). oMEGACat. VI. Analysis of the overall kinematics of Omega Centauri in 3D: Velocity dispersion, kinematic distance, anisotropy, and energy equipartition. *The Astrophysical Journal*, 983:95.

Hanson, R. (1998). Burning the cosmic commons: Evolutionary strategies for interstellar colonization. Working paper, Department of Economics, George Mason University. July 1, 1998; https://mason.gmu.edu/~rhanson/filluniv.pdf.

Hanson, R., Martin, D., McCarter, C., and Paulson, J. (2021). If loud aliens explain human earliness, quiet aliens are also rare. *The Astrophysical Journal*, 922(2):182.

Haqq-Misra, J. D. and Baum, S. D. (2009). The sustainability solution to the Fermi paradox. *Journal of the British Interplanetary Society*, 62:47–51.

Heller, R. (2017). Relativistic generalization of the incentive trap of interstellar travel with application to Breakthrough Starshot. *Monthly Notices of the Royal Astronomical Society*, 470:3664–3671.

Heller, R. and Hippke, M. (2017). Deceleration of high-velocity interstellar photon sails into bound orbits at α Centauri. *The Astrophysical Journal Letters*, 835:L32.

Hoang, T., Lazarian, A., Burkhart, B., and Loeb, A. (2017). The interaction of relativistic spacecrafts with the interstellar medium. *The Astrophysical Journal*, 837(1):5.

Hooper, D. (2018). Life versus dark energy: How an advanced civilization could resist the accelerating expansion of the universe. arXiv:1806.05203; unpublished as of 2026-08-16.

Hosek, W. R. (2007). Economics and the Fermi paradox. *Journal of the British Interplanetary Society*, 60:137–141.

Huang, B.-L., Tao, Z.-Z., and Zhang, T.-J. (2026). Phase-space crystallization in Galactic globular clusters: A Gaia-based metric and implications for technosignature searches. *The Astrophysical Journal*, 1005:102.

Ivliev, S. (2026). Autonomous AI-cosmoindustry and the quiet expansion filter: A threshold-based resolution of the Fermi paradox. arXiv:2606.13914; preprint.

Laibson, D. (1997). Golden eggs and hyperbolic discounting. *The Quarterly Journal of Economics*, 112(2):443–478.

Landauer, R. (1961). Irreversibility and heat generation in the computing process. *IBM Journal of Research and Development*, 5:183–191.

Lubin, P. (2016). A roadmap to interstellar flight. *Journal of the British Interplanetary Society*, 69:40–72.

McDonald, R. and Siegel, D. (1986). The value of waiting to invest. *The Quarterly Journal of Economics*, 101:707–727.

Olson, S. J. (2015). Homogeneous cosmology with aggressively expanding civilizations. *Classical and Quantum Gravity*, 32:215025.

Ord, T. (2021). The edges of our universe. arXiv:2104.01191; unpublished as of 2026-08-16.

Perakis, N. and Hein, A. M. (2016). Combining magnetic and electric sails for interstellar deceleration. *Acta Astronautica*, 128:13–20.

Ramsey, F. P. (1928). A mathematical theory of saving. *The Economic Journal*, 38(152):543–559.

Ren, T., Garrett, M. A., and Siemion, A. P. V. (2024). Background contamination of the Project Hephaistos Dyson spheres candidates. *Research Notes of the AAS*, 8(5):145.

Ren, T., Garrett, M. A., Siemion, A. P. V., et al. (2025). High-resolution imaging of the radio source associated with Project Hephaistos Dyson sphere candidate G. *Monthly Notices of the Royal Astronomical Society: Letters*, 538:L56–L61.

Sandberg, A., Armstrong, S., and Ćirković, M. M. (2016). That is not dead which can eternal lie: the aestivation hypothesis for resolving Fermi's paradox. *Journal of the British Interplanetary Society*, 69(11):406–415.

Sandberg, A., Drexler, E., and Ord, T. (2018). Dissolving the Fermi paradox.

Scoggins, M. T. and Kipping, D. (2023). Lazarus stars: numerical investigations of stellar evolution with star-lifting as a life extension strategy. *Monthly Notices of the Royal Astronomical Society*, 523:3251–3257.

Suazo, M., Zackrisson, E., Mahto, P. K., Lundell, F., Nettelblad, C., Korn, A. J., Wright, J. T., and Majumdar, S. (2024). Project Hephaistos – II. Dyson sphere candidates from Gaia DR3, 2MASS, and WISE. *Monthly Notices of the Royal Astronomical Society*, 531:695–707.

Swanson, T. (2026a). Engineered intermediate-mass black hole systems: Infrastructure constraints, observable residue, and a multi-messenger adjudication framework. Preprint; omegacentauri.me.

Swanson, T. (2026b). Inward resolutions of the Fermi paradox: A critical review of migration down thermodynamic gradients. Preprint; omegacentauri.me/paper.html.

Swanson, T. (2026c). The macro transcension hypothesis: Spinning black holes in dense stellar clusters as thermodynamic attractors for advanced civilizations, with Omega Centauri as an observational test bed. Preprint; omegacentauri.me/paper.html.

Swanson, T. (2026d). A multi-messenger technosignature and anomaly-detection campaign for Omega Centauri. Preprint; omegacentauri.me/campaign.html.

Weitzman, M. L. (1998). Why the far-distant future should be discounted at its lowest possible rate. *Journal of Environmental Economics and Management*, 36(3):201–208.

Weitzman, M. L. (2001). Gamma discounting. *American Economic Review*, 91(1):260–271.

Wright, J. T. (2023). Application of the thermodynamics of radiation to Dyson spheres as work extractors and computational engines, and their observational consequences. *The Astrophysical Journal*, 956:34.

Wright, J. T., Mullan, B., Sigurdsson, S., and Povich, M. S. (2014). The ĝ infrared search for extraterrestrial civilizations with large energy supplies. I. Background and justification. *The Astrophysical Journal*, 792(1):26.

Zackrisson, E., Korn, A. J., Wehrhahn, A., and Reiter, J. (2018). SETI with Gaia: the observational signatures of nearly complete Dyson spheres. *The Astrophysical Journal*, 862:21.

Zubrin, R. M. and Andrews, D. G. (1991). Magnetic sails and interplanetary travel. *Journal of Spacecraft and Rockets*, 28(2):197–203.