

An erratum to the jerk noise-floor constant in Prager et al. (2017) and Abbate et al. (2019)

TIM SWANSON¹

¹*The Omega Centauri Society*

ABSTRACT

Pulsar spin-frequency second derivatives, jerks, probe cluster gravitational potentials and central masses against a stochastic floor of nearest-neighbour stellar encounters, whose characteristic amplitude is $\dot{a}_0 = (2\pi\xi/3) G\langle m\rangle\sigma n$, with n the stellar number density, σ the velocity dispersion, and $\langle m\rangle$ the mean stellar mass. Prager et al. (2017) quote the dimensionless constant $\xi \simeq 3.04$, and Abbate et al. (2019) adopt it. The double integral that defines ξ evaluates in closed form to $\sqrt{2\pi}[1 + \ln(2 + \sqrt{3})/(2\sqrt{3})] = 3.4595806610$: five evaluations of that one integral (a closed-form reduction, adaptive quadrature, symbolic evaluation, antiderivative differentiation, and high-precision quadrature) plus two independently coded Monte Carlo simulations of the underlying stellar field that never evaluate the integral, at 15.8σ and 14.6σ respectively, both reject 3.04; Section 3 states the common-mode caveat on the first five. Because $\dot{a}_0$ is linear in ξ , the printed constant understates the jerk floor by 12.1 per cent, inflating the apparent significance of jerk-based central-mass inferences.

Keywords: pulsars: general – globular clusters: general – methods: analytical – methods: statistical

1. INTRODUCTION

Long-term timing of millisecond pulsars in globular clusters measures spin-frequency second derivatives, jerks, which probe the cluster gravitational potential and any intermediate-mass black hole it hosts (Prager et al. 2017; Abbate et al. 2019). Jerks from passing cluster stars follow the statistics of random backgrounds long studied as source confusion in radio astronomy (Condon 1974): nearest neighbours dominate, the sum has no finite variance, and its scale is set by a characteristic amplitude,

$$\dot{a}_0 = \frac{2\pi\xi}{3} G\langle m\rangle\sigma n, \quad (1)$$

rather than by an rms. Here n is the stellar number density, σ the one-dimensional velocity dispersion, $\langle m\rangle$ the mean stellar mass, and ξ a dimensionless constant collecting all of the geometry (Prager et al. 2017). Both papers quote $\xi \simeq 3.04$.

While implementing this formalism for a study of NGC 5139 (ω Cen) now in preparation, I found that the quoted value disagrees with the integral that defines it, which has a closed-form evaluation (Section 2). Section 3 reports the verification, five evaluations of the integral plus two independently coded Monte Carlo simulations of the physical field; Section 4 traces how the error entered the published literature; Section 5 states the consequence for jerk-based inference.

2. THE INTEGRAL AND ITS CLOSED FORM

Prager et al. (2017) define ξ by the double integral of their arXiv v1 Equation B7,

$$\xi = \int_0^\infty dx \int_{-1}^1 d\mu \left[1 - \exp\left(-\frac{1+3\mu^2}{2x^2}\right) \right]. \quad (2)$$

For fixed μ the bracket depends on x only through a/x^2 with $a \equiv 1 + 3\mu^2$, and the inner integral evaluates to $\sqrt{\pi/2}(1+3\mu^2)^{1/2}$. The remaining integral is elementary, $\int_{-1}^1 (1+3\mu^2)^{1/2} d\mu = 2 + \operatorname{asinh} \sqrt{3}/\sqrt{3}$, and $\operatorname{asinh} \sqrt{3} = \ln(2 + \sqrt{3})$, so

$$\xi = \sqrt{2\pi} \left[1 + \frac{\ln(2 + \sqrt{3})}{2\sqrt{3}} \right] = 3.4595806610, \quad (3)$$

or, to fifty digits, 3.45958066104621561909285714056923304379183879500167. The closed form exceeds the printed value by 13.8 per cent.

3. VERIFICATION

I verified Equation (3) with five evaluations of Equation (2) itself, (i) (ii) (iv) (v) (vi) below, and two independently coded Monte Carlo simulations of the physical system, (iii), that never evaluate the formula. Because routes (i), (ii), (iv), (v), and (vi) all evaluate or manipulate the same double integral, they share whatever algebraic or limiting error the integral itself might

carry; only the two Monte Carlo passes are independent of Equation (2), and the rejection of 3.04 rests on those. (i) The closed form itself. (ii) Adaptive quadrature of Equation (2), agreeing with the closed form to 2.1×10^{-13} relative (7.3×10^{-13} absolute). (iii) Two Monte Carlo passes of the physical system, described next. (iv) End-to-end symbolic evaluation of both integrals, returning the inner result above and $2 + \operatorname{asinh} \sqrt{3}/\sqrt{3}$ for the outer one. (v) Differentiation of the closed-form antiderivative used in the reduction, which simplifies identically to zero against the integrand. (vi) Arbitrary-precision quadrature of Equation (2) at 55 decimal digits (mpmath), substituting $x = \tan \theta$ to remove the $1/x^2$ tail rather than truncating it, agreeing with the closed form to 3.1×10^{-35} relative (fH_verify_xi_mpmath_quad.py, deposited with the others; a 70-digit attempt did not converge in reasonable time). This route is a true numerical quadrature of the double integral, not arithmetic on the closed form.

Route (iii), first pass, never evaluates the formula. Each of 4×10^4 realizations draws Poisson(nV) stars uniformly in a sphere, assigns Maxwellian velocities, sums the line-of-sight single-star jerk of Prager et al.’s Equation B1 directly, and estimates the Cauchy scale from the interquartile range, equal to twice the scale for a Cauchy law and insensitive to the heavy tail that invalidates the sample mean. The result is $\xi_{\text{MC}} = 3.4886 \pm 0.0283$: consistent with Equation (3) at 1.0σ and excluding 3.04 at 15.8σ (Figure 1). The same sample reproduces the published line-of-sight law in shape, with a Kolmogorov–Smirnov $p = 0.26$ against the Cauchy form, and in tail index, with a Hill estimate of 1.025 against the 1-stable value of 1.

A second, separately coded Monte Carlo (fH_verify_xi_second_mc.py, deposited with the others) built directly on the projected line-of-sight jerk expression with an independent seed, drew 4×10^4 realizations, and estimated the Lorentzian scale from the interquartile range with a bootstrap standard error over 2000 resamples: $\xi_{\text{MC},2} = 3.4403 \pm 0.0275$, agreeing with Equation (3) at -0.7σ and excluding 3.04 at 14.6σ . The two Monte Carlo passes sit on opposite sides of the closed form, at $+1.0\sigma$ and -0.7σ , consistent with sampling scatter between independent implementations; a check of the interquartile-range Cauchy-scale estimator itself, 3000 synthetic draws of 4×10^4 standard-Cauchy variates each, finds it unbiased to within 0.02 per cent, so the scatter is not an estimator artifact. The rejection of 3.04 rests on the two Monte Carlo passes; the 13.8 per cent correction rests on the closed form.

I also re-derived Equations B4–B10 through the characteristic function, reproducing every step including the




Figure 1. Route (iii). Line-of-sight jerk sums from 4×10^4 realizations of a Poisson stellar field with Maxwellian velocities, in the scaled units $G = \langle m \rangle = \sigma = n = 1$, against the Cauchy law of Prager et al.’s Equation B10 at the corrected scale ($\xi = 3.4596$, solid) and the printed scale ($\xi \approx 3.04$, dashed). The interquartile estimate for this sample is $\xi_{\text{MC}} = 3.4886 \pm 0.0283$, excluding 3.04 at 15.8σ .

angular factor $s^2(1 + 3\mu^2)$. The algebra is sound; the discrepancy is confined to the numerical evaluation of Equation (2). The verification scripts and the Monte Carlo drivers are prepared for a Zenodo deposit (deposit pending; DOI to be added before submission).

4. PROVENANCE IN THE PUBLISHED SOURCES

The derivation survives in full only in version 1 of the arXiv posting of Prager et al. (2017) (1612.04395v1), where Appendix B carries Equations B1–B10 and Equation B7 defines ξ as the double integral. In the v1 source the defining equation reads, for $s \geq 0^2$

$$C(s) = \frac{2\pi G s}{3} \int dm P(m) m \sigma \int_0^\infty dx \int_{-1}^1 d\mu \times \left(1 - e^{-(1+3\mu^2)/2x^2}\right) \quad (4)$$

$$\equiv \frac{2\pi \xi}{3} G \langle m \rangle \sigma s,$$

² As printed, this equation is dimensionally inconsistent with Equation (1): both sides carry no factor of the stellar number density n , so $C(s)$ has dimensions of volume rather than being dimensionless. Reducing the right-hand side with $x \equiv r^3/(sGm\sigma)$, so that $d^3r = 2\pi(sGm\sigma/3) dx d\mu$, gives $C(|s|) = (2\pi/3) n G \langle m \rangle \sigma |s| \xi$; the reduction also pulls σ outside $\int dm P(m)$, which holds only if the velocity dispersion is assumed independent of stellar mass, an assumption Prager et al. leave implicit. I reproduce the source equation as printed and note the omission here rather than silently inserting n into a quotation.

where $C(s)$ is the cumulant of the line-of-sight jerk’s characteristic function at conjugate variable $s \geq 0$ (the full symmetric-Cauchy cumulant is linear in $|s|$, not s), followed, in the same paragraph, by the sentence “where $\langle m \rangle = \int dm m P(m)$ and $\xi \simeq 3.04$ is the numerical value of the integral.” Integral and number share a sentence, so no alternative convention can reconcile them: the text offers 3.04 as the value of this integral, and the integral equals 3.4596.

The published article (ApJ, 845, 148) truncates Appendix B after Equation B3 and so removes the definition of ξ , while main-text Equation 8 retains $\xi \simeq 3.04$ and points to Appendix B for its origin; a reader checking the constant against the journal version alone cannot reproduce it from what is printed. Equation 6 of [Abbate et al. \(2019\)](#) carries the same constant, citing [Prager et al. \(2017\)](#). A citation screen of the works citing [Prager et al. \(2017\)](#) or [Abbate et al. \(2019\)](#) (OpenAlex, 2026 October 3; 102 works; per-title dispositions deposited with the verification scripts) finds no published correction to the constant, and exactly two works adopt the printed value in an inference: Equation 6 of [Abbate et al. \(2019\)](#), and [Corongiu et al. \(2024\)](#), who print “where $\xi \simeq 3.04$

is a numerical constant” in the characteristic-jerk relation and use it in a published core-mass inference for NGC 6752.

5. CONSEQUENCE FOR JERK-BASED INFERENCE

By Equation (1) the floor is linear in ξ , so the printed constant understates the stochastic jerk floor by 12.1 per cent. The direction of the error is the unfavourable one for inference: a floor assumed too low inflates the apparent significance of any jerk-based central-mass constraint, including the Bayes factors that [Abbate et al. \(2019\)](#) tabulate, which inherit the constant. I have not recomputed those values; a 12 per cent rescaling of the neighbour term is unlikely to move qualitative conclusions, but quantitative claims near threshold deserve re-examination by the groups concerned. Any analysis using the formalism should state which value of ξ it adopts.

Software: NumPy, SciPy, SymPy, mpmath, Matplotlib

REFERENCES

- Abbate, F., Spera, M., & Colpi, M. 2019, MNRAS, 487, 769 (arXiv:1905.05209)
- Corongiu, A., Ridolfi, A., Abbate, F., et al. 2024, ApJ, 972, 198 (arXiv:2407.03271)
- Condon, J. J. 1974, ApJ, 188, 279
- Prager, B. J., Ransom, S. M., Freire, P. C. C., Hessels, J. W. T., Stairs, I. H., Arras, P., & Cadelano, M. 2017, ApJ, 845, 148 (arXiv:1612.04395)