

A Multi-Messenger Technosignature and Anomaly-Detection Campaign for Omega Centauri

Tim Swanson

The Omega Centauri Society / Post Oak Labs

tim@postoaklabs.com

July 2026

Draft v1.1 (last revised 2026-07-18) — third paper of the set; companion to *The Macro Transcension Hypothesis* (Paper A), the inward-migration review (Paper B), the migration economics (Paper D), and the engineered-IMBH systems paper (Paper E); prepared for omegacentauri.me; intended for submission to arXiv

Abstract

Omega Centauri (NGC 5139), the most massive Galactic globular cluster and the probable stripped nucleus of an accreted dwarf galaxy, presents a unique conjunction of observational circumstances: the strongest current candidate for an intermediate-mass black hole (IMBH) in the Galaxy, anchored by seven stars moving above the local escape velocity (Häberle et al., 2024b); a formally unresolved factor-of-several tension between kinematic lower bounds ($\geq 8,200 M_{\odot}$) and a pulsar-timing upper bound ($\lesssim 6,000 M_{\odot}$; Bañares-Hernández et al. 2025); complete electromagnetic silence to the deepest radio and infrared limits ever placed on a globular cluster core (Mahida et al., 2026; Chen et al., 2025a); a southern declination (-47.5°) optimal for the newest southern-hemisphere facilities; and a 12-Gyr-old stellar population. To our knowledge, no dedicated technosignature search of ω Cen has ever been conducted at any wavelength: the first dedicated globular-cluster SETI survey (Huang et al., 2026b) could not observe it. We present a coordinated, hypothesis-agnostic, multi-messenger observational campaign that addresses both conventional astrophysical questions (IMBH reality, mass, and spin; cluster dynamics) and technosignature/anomaly hypotheses (including the inward-migration class surveyed in the companion review; Swanson 2026b) with the same data. The campaign comprises eight instrument-matched programs: (1) JWST infrared accretion and waste-heat limits; (2) deep radio imaging, narrowband SETI, and millisecond-pulsar timing with MeerKAT, transitioning to SKA-Mid; (3) astrometric monitoring with HST, Gaia DR4, the Nancy Grace Roman Space Telescope (launch scheduled for late 2026, with the agency commitment extending to 2027), and ELT/MICADO; (4) a LISA extreme-mass-ratio-inspiral forecast; (5) a KM3NeT/ARCA neutrino burst pipeline exploiting ω Cen's up-going geometry from the Mediterranean; (6) Fermi-LAT archival analysis and CTAO-South follow-up; (7) optical/infrared time-domain and archival anomaly searches with Rubin/LSST and oMEGACat; and (8) supplementary archival channels (LIGO–Virgo–KAGRA continuous waves; Pierre Auger ultra-high-energy cosmic rays). For each program we state quantitative sensitivity estimates, time requests, decision thresholds, and explicit falsification criteria, including negative results: we show that direct astrometric acceleration detection of the fast stars is below 1σ at nominal parameters before ~ 2040 , and route the astrometric science through photocentric-wander and reference-frame measurements instead. Total program cost is $\lesssim$ US\$7M over 2026–2035, most of it archival analysis and piggyback observing on already-funded facilities; the decisive measurements (IMBH mass and spin via LISA) arrive as free by-products of planned mission science. The campaign is constructed so that every null result constrains conventional astrophysics, and every anomaly is adjudicated by at least two independent messengers.

Keywords: Omega Centauri; NGC 5139; intermediate-mass black holes; technosignatures; SETI; multi-messenger astronomy; millisecond pulsars; gravitational waves; neutrino astronomy

Contents

1	Introduction	5
1.1	Why Omega Centauri, and why now	5
1.2	The technosignature rationale	5
1.3	Design principles	6
1.4	Structure of this paper	7
2	The target and the constraint landscape	7
2.1	Adopted parameters	7
2.2	The mass tension	7
2.3	Three hypotheses, fixed in advance	9
3	Program 1: JWST infrared accretion and waste-heat limits	9
3.1	Science case	9
3.2	Implementation	9
3.3	Decision criteria	10
4	Program 2: Radio — deep imaging, narrowband SETI, and pulsar timing	10
4.1	Science case	10
4.2	Implementation: MeerKAT era (2026–2030)	11
4.3	Implementation: SKA era (2029–)	11
4.4	Decision criteria	12
5	Program 3: Astrometry — HST, Gaia DR4, Roman, and ELT/MICADO	12
5.1	Sensitivity of the direct acceleration test	12
5.2	Three astrometric measurements that do work	12
5.3	Decision criteria	14
6	Program 4: The LISA forecast	14
6.1	Science case	14
6.2	Occurrence rates	14
6.3	The spin measurement as the H_2 adjudicator	15
6.4	Deliverables before launch	15
7	Program 5: KM3NeT/ARCA neutrino monitoring	15
7.1	Science case and geometry	15
7.2	Implementation	15
7.3	Decision criteria	16
8	Program 6: Gamma rays — Fermi-LAT archival and CTAO-South	16
8.1	Science case	16
8.2	Implementation	17
8.3	Decision criteria	17

9	Program 7: Optical/IR time domain — Rubin/LSST and archival anomalies	17
9.1	Science case	17
9.2	Implementation	18
9.3	Decision criteria	18
10	Program 8: Supplementary archival channels	18
11	Coordination, joint statistics, and data policy	19
11.1	The two-messenger rule, operationalized	19
11.2	Alert latency and standing authorizations	19
11.3	Data policy	19
11.4	Program summary	20
12	Cost and timeline	20
12.1	Costing	20
12.2	Timeline	20
13	The decision structure through 2040	22
14	Conclusion	22

1 Introduction

1.1 Why Omega Centauri, and why now

Three independent developments between 2024 and 2026 have transformed Omega Centauri (ω Centauri, NGC 5139) from one interesting globular cluster among many into arguably the single best-posed multi-messenger target in the Galaxy outside the Galactic Centre.

First, the IMBH candidacy became concrete. Häberle et al. (2024b), using a proper-motion catalogue of 1.4 million stars built from over 500 HST epochs (Häberle et al., 2024a), identified seven stars within 3 arcsec of the cluster centre moving faster than the local escape velocity: stars that require a bound central mass of $\geq 8,200 M_{\odot}$ from their velocities alone, and $\geq 21,100 M_{\odot}$ at 99 per cent confidence when acceleration limits are included. Independent N -body modelling finds a best-fitting in-situ-grown IMBH of $\sim 4.7\text{--}5.1 \times 10^4 M_{\odot}$ (González Prieto et al., 2025). Against this, a joint analysis of stellar kinematics and millisecond-pulsar (MSP) timing places a 3σ upper limit of $\sim 6 \times 10^3 M_{\odot}$ on any central point mass, favouring instead an extended $\approx 2\text{--}3 \times 10^5 M_{\odot}$ component of stellar remnants (Bañares-Hernández et al., 2025). The constraints are formally inconsistent under their stated assumptions. A factor-of-several disagreement between two mature measurement techniques, centred on the nearest IMBH candidate in the sky, warrants a focused observational campaign to resolve it.

Second, the electromagnetic silence became quantitative. The deepest radio observation ever made of a globular cluster (~ 170 h with ATCA, reaching $1.1 \mu\text{Jy beam}^{-1}$ rms) detects nothing at any proposed cluster centre, bounding the accretion efficiency of a putative IMBH at $\lesssim 4 \times 10^{-3}$ of the Bondi rate (Mahida et al., 2026), extending two decades of radio non-detections of globular-cluster IMBHs (Strader et al., 2012; Tremou et al., 2018). JWST NIRCам and MIRI photometry of the central field likewise finds no source with the spectral energy distribution of an accreting IMBH (Chen et al., 2025a). Whatever sits at the centre of ω Cen is dark to a depth that is a scientific result in its own right.

Third, the instrument landscape shifted decisively in the campaign’s favour. Within an eighteen-month window centred on this writing: the Vera C. Rubin Observatory began the Legacy Survey of Space and Time (LSST) in early 2026 (Ivezić et al., 2019); the Nancy Grace Roman Space Telescope completed construction and is scheduled for launch in late 2026, with the agency commitment extending to 2027 (WFIRST Astrometry Working Group et al., 2019); Gaia Data Release 4, with a 5.5-year astrometric baseline, is scheduled for December 2026 (Gaia Collaboration et al., 2016); KM3NeT/ARCA reached 51 of 230 planned detection units (as of early 2026) with real-time alerts entering commissioning (Adrián-Martínez et al., 2016; The KM3NeT Collaboration, 2025); the first telescopes of CTAO-South are, as of early 2026, arriving at Paranal (Cherenkov Telescope Array Consortium et al., 2019); SKA-Mid is assembling toward 2029 science verification (Braun et al., 2019); ELT first light is scheduled for 2029 (Davies et al., 2021); and LISA, formally adopted by ESA in January 2024, is in hardware development for a 2035 launch (Colpi et al., 2024). Nearly every facility on this list is southern-hemisphere or all-sky; ω Cen, at declination -47.5° , sits in the sweet spot of all of them. A campaign organized now can ride this entire wave at marginal cost.

1.2 The technosignature rationale

This paper is the observational companion to a speculative hypothesis paper (Swanson, 2026c, hereafter Paper A), which argues that rapidly spinning massive black holes in dense old stellar systems are thermodynamic attractors for hypothetical computation-optimizing civilizations, and that ω Cen is the most accessible system satisfying the resulting selection criteria. We neither assume nor argue for that hypothesis here. The campaign is designed to be *hypothesis-agnostic* in the specific sense that

every observation in it is independently justified by conventional astrophysics: IMBH demographics (Greene et al., 2020), cluster dynamics, pulsar timing, accretion physics at the lowest Eddington ratios, and EMRI astrophysics (Amaro-Seoane, 2018). The technosignature dimension adds analysis channels (narrowband drift searches, burst-coincidence triggers, anomaly statistics) to data that would be taken anyway, in the cost-effective tradition of commensal SETI (Tarter, 2001; Wright et al., 2022; Lacki and DiKerby, 2025), consistent with the archival anomaly-detection strategy recommended by the Dyson Minds workshop for black-hole-hosted intelligence (Curtis et al., 2026), and aligned with recent calls for integrated, multi-channel technosignature-search strategies as the efficient way to constrain Fermi-paradox hypotheses (Crawford, 2026).

Three considerations nevertheless justify making the technosignature dimension explicit rather than incidental. First, globular clusters are almost entirely unexplored SETI territory: the first dedicated globular-cluster survey, a FAST pilot of five northern clusters, was published only this year and could not observe ω Cen from FAST’s latitude (Huang et al., 2026b); Gaia-based dynamical metrics for prioritizing clusters as technosignature targets have followed (Huang et al., 2026a). Second, ω Cen is a privileged target on entirely generic arguments: 10^7 stars of age 12.08 ± 0.01 Gyr (Clontz et al., 2024) in a beam a few arcminutes across, a possible IMBH, and (as the probable nucleus of an accreted dwarf; Hilker and Richtler 2000; Ibata et al. 2019) an independent galactic chemical-evolution history, so that any ETI prior that weights stellar age and number density at all concentrates sharply here. Third, the high-energy technosignature channels (neutrino bursts from hypothetical black-hole-based computation; Dvali and Osmanov 2023) happen to be testable at ω Cen essentially for free, because the cluster’s declination makes it an up-going source for the Mediterranean neutrino telescopes. Where Paper A’s specific predictions sharpen a test, we say so explicitly and label the dependence; readers uninterested in that hypothesis may strike those sentences without loss to the campaign.

Epistemic status: The observational programs of Sections 3–10 stand on conventional astrophysics and are stated in the empirical register. The technosignature interpretation of any anomaly they return belongs to the speculative register, is conditional on the optimization premise of Papers A and B, and is adjudicated by the quantitative framework of Paper E rather than by this campaign.

1.3 Design principles

Five principles govern the campaign design, and we state them up front because they discipline everything that follows.

(i) Dual-use data. Every observing program must produce publishable conventional astrophysics on its own; the technosignature analysis is a parallel pipeline on the same photons (or neutrinos, or strain data), never a dedicated expenditure that a null result would waste.

(ii) Honest sensitivity accounting. Where a measurement cannot work at nominal parameters, we say so quantitatively. Section 5 demonstrates this standard: the direct acceleration test on the fast stars, superficially the most natural follow-up to Häberle et al. (2024b), is below 1σ before ~ 2040 for plausible masses, and we restructure the astrometric program accordingly.

(iii) Two-messenger adjudication. No anomaly claim survives on one channel. Every positive trigger (a neutrino multiplet, a narrowband line, an infrared transient) must specify in advance which independent channel confirms or kills it, with combined significance assessed by pre-registered methods (Section 11).

(iv) Pre-registered thresholds. Detection and falsification thresholds are stated numerically before the data arrive (Tables 4 and 3), following standard pre-registration practice.

(v) Parasitism on funded science. The decisive measurements (IMBH reality, mass, and spin) will be made by LISA as part of its core science program whether or not anyone organizes a campaign (Colpi et al.,

Table 1: Adopted parameters for ω Centauri (NGC 5139).

Quantity	Value	Source
Distance	5.49 ± 0.06 kpc	Häberle et al. (2025)
Cluster mass	$\approx 4 \times 10^6 M_\odot$	Harris (1996) ; Baumgardt and Vasiliev (2021)
Stellar count	$\sim 10^7$	Harris (1996)
Mean stellar age	12.08 ± 0.01 Gyr (spread 0.75 Gyr)	Clontz et al. (2024)
Origin	stripped dwarf-galaxy nucleus	Hilker and Richtler (2000) ; Ibata et al. (2019) ; Souza et al. (2024)
Position (J2000)	RA $13^h 26^m 47^s$, Dec $-47^\circ 28' 46''$	Harris (1996)
Known millisecond pulsars	18	Dai et al. (2020) ; Chen et al. (2023) ; Colomo i Bernadich et al. (2023)
Central γ -ray source	4FGL J1326.7–4729 (MSP ensemble)	Dai et al. (2020, 2023)
Angular scale	$1'' = 0.0266$ pc	—

[2024; Babak et al., 2017](#)). The campaign’s job is to ensure that when those numbers arrive, the contextual data (timing, astrometry, electromagnetic limits, high-energy monitoring) already exist to interpret them.

1.4 Structure of this paper

Section 2 reviews the target and the present constraint landscape. Sections 3–10 present the eight programs, each with science case, technical implementation, sensitivity estimates, and decision criteria. Section 11 describes cross-program coordination, alert protocols, and joint statistics. Section 12 presents costing and timeline. Section 13 assembles the decision tree through 2040. Section 14 concludes.

2 The target and the constraint landscape

2.1 Adopted parameters

Table 1 collects the parameters adopted throughout. We use the oMEGACat kinematic distance of 5.49 ± 0.06 kpc ([Häberle et al., 2025](#)), consistent within systematics with the Gaia EDR3 parallax distance ([Soltis et al., 2021](#)) and the combined-catalogue value ([Baumgardt and Vasiliev, 2021](#)). At this distance $1''$ subtends 0.0266 pc, so the entire fast-star region ($r < 3''$) fits within 0.08 pc, and the cluster core ($r_c \approx 2.4'$) within ~ 4 pc.

2.2 The mass tension

Table 2 and Figure 1 summarize the central-mass constraints. Two features matter for campaign design. First, the disagreement is not marginal: the velocity-only kinematic lower bound ($8,200 M_\odot$) exceeds the pulsar-timing point-mass upper bound ($6,000 M_\odot$, 3σ) outright, and the acceleration-informed lower bound ($21,100 M_\odot$) exceeds it by a factor of 3.5. At least one analysis is wrong, or incomplete, in an instructive way: for instance through the radial distribution assumed for the MSPs, the foreground/membership treatment of the fast stars, or the possibility that the central mass is genuinely extended ([Bañares-Hernández et al., 2025](#); [Zocchi et al., 2019](#); [Breen and Hogg, 2013](#)). Second, the tension is resolvable on a known schedule: more pulsars and longer timing baselines tighten the upper bound as roughly $1/\sqrt{N_{\text{psr}} T^2}$ (Section 4); Roman and Gaia DR4 sharpen the astrometric frame within two years (Section 5); and LISA, if a compact-object inspiral is caught, measures the mass to ~ 0.1 per cent ([Babak et al., 2017](#)).

Table 2: Current constraints on the central dark mass of ω Cen (cf. Paper A, Table 2).

Constraint	Value	Method / source
Lower bound (velocities only)	$\geq 8,200 M_{\odot}$	7 fast stars, HST proper motions (Häberle et al., 2024b)
Lower bound (with accelerations)	$\geq 21,100 M_{\odot}$ (99%)	same stars, acceleration limits (Häberle et al., 2024b)
N -body growth models	$\sim 4.7\text{--}5.1 \times 10^4 M_{\odot}$	cluster-evolution simulations (González Prieto et al., 2025)
Upper bound (point mass)	$< 6 \times 10^3 M_{\odot}$ (3σ)	stellar kinematics + MSP timing (Bañares-Hernández et al., 2025)
Favoured alternative	extended $2\text{--}3 \times 10^5 M_{\odot}$	same analysis (Bañares-Hernández et al., 2025)
Timing (joint MeerKAT–Parkes)	insensitive at $10^3\text{--}10^4 M_{\odot}$; $< 10^5 M_{\odot}$ (90%)	MSP timing (Colomo & Bernadich et al., 2026)
Confirmed stellar-mass BH	$4.46^{+1.22}_{-1.01} M_{\odot}$ (94-yr binary, $a = 31$ AU)	HST/JWST astrometry (Whitaker et al., 2026)
Historical upper limits	$\lesssim 10^4 M_{\odot}$ (model-dep.)	HST proper motions (van der Marel and Anderson, 2010); see also Noyola et al. (2008); Zocchi et al. (2019)




Figure 1: The central-mass constraint landscape (logarithmic mass axis). The pulsar-timing point-mass upper limit and the fast-star kinematic lower bounds leave no allowed point mass: at least one constraint must yield. The campaign is structured to determine which, on a known schedule, using instruments that share no dominant systematics.

2.3 Three hypotheses, fixed in advance

Following Paper A we carry three competing hypotheses through every program:

H_0 (**gas-starved IMBH or near-IMBH**). A genuine central point mass in the $\sim 10^4 M_\odot$ range exists and is silent for the ordinary reason: a relaxed, gas-poor, 12-Gyr-old cluster supplies essentially nothing to accrete. Default and most parsimonious if the kinematic bounds hold.

H_1 (**extended remnant component**). No IMBH; the central mass is a spatially extended cluster of stellar remnants (Bañares-Hernández et al., 2025; Breen and Hoggie, 2013; Zocchi et al., 2019). Default and most parsimonious if the timing bound holds.

H_2 (**anomalous/managed system**). A central IMBH whose observational properties are shaped by technology, per Paper A. Carries a low prior; becomes interesting only through conjunctions that H_0 and H_1 jointly fail to explain (e.g. near-extremal spin on a gas-starved hole; repeated coincident multi-messenger bursts; statistically anomalous depletion of the loosely bound core population).

The campaign’s purpose is to measure the system well enough that H_0 and H_1 are decided on their merits, with H_2 ’s distinctive residues either appearing in the data or being retired by them, rather than to hunt for H_2 . These labels map one-to-one onto the hypothesis set of the companion adjudication framework (Swanson, 2026a, hereafter Paper E): H_0 corresponds to Paper E’s H_q (quiescent astrophysical IMBH), H_1 to H_{sub} (sub-IMBH remnant alternative), and H_2 to H_{eng} (engineered system).

3 Program 1: JWST infrared accretion and waste-heat limits

3.1 Science case

For any central mass $M_\bullet$, the Eddington luminosity is $L_{\text{Edd}} \simeq 1.26 \times 10^{31} (M_\bullet/M_\odot) \text{ W}$: $1.0 \times 10^{35} \text{ W}$ at $8,200 M_\odot$, $2.5 \times 10^{35} \text{ W}$ at $2 \times 10^4 M_\odot$, $6.2 \times 10^{35} \text{ W}$ at $4.9 \times 10^4 M_\odot$. The existing non-detections already establish extreme quiescence: the radio limit bounds accretion at $\lesssim 4 \times 10^{-3}$ of the Bondi rate (Mahida et al., 2026), the archival Chandra limit bounds the X-ray output at $L_X \lesssim 1.6 \times 10^{30} \text{ erg s}^{-1}$ (Haggard et al., 2013), and the JWST photometry bounds the radiative output at $\lesssim 10^{-9}$ of Eddington (Chen et al., 2025a), among the most extreme quiescence levels measured for any black hole. The published JWST analysis, however, is an archival by-product, not an optimized deep program. A purpose-designed campaign improves the limits by an order of magnitude and adds the two dimensions the archival data lack: variability (accretion at these rates is expected to flicker; a managed or artificial suppression of accretion, the H_2 residue, is not) and waste heat (a mid-infrared excess over the stellar population model is the classic Dyson-type technosignature; Dyson 1960; Wright et al. 2014; Hsiao et al. 2021, and at ω Cen the relevant solid angle is a few arcseconds, not a galaxy).

3.2 Implementation

Deep imaging (new observations). NIRCam F200W/F356W/F444W at 20 ks per filter plus MIRI F770W/F1000W at 15 ks per filter, in two epochs separated by ~ 6 months: $\sim 65 \text{ h}$ total. In the crowded core the limits are confusion- rather than photon-limited; with empirical-PSF subtraction anchored to the oMEGACat astrometric catalogue (Häberle et al., 2024a; Nitschai et al., 2023), simulated point-source recovery reaches $\sim 5\text{--}10 \text{ nJy}$ (3σ) at F444W and $\sim 30\text{--}70 \text{ nJy}$ at F770W/F1000W after a $\sim 3\times$ crowding penalty. At 5.49 kpc, 10 nJy at F444W corresponds to a monochromatic $\nu L_\nu \approx 2.5 \times 10^{28} \text{ erg s}^{-1}$; for

plausible quiescent-accretion SEDs the bolometric limit is of order 10^{29} – 10^{30} erg s^{−1}, i.e. Eddington ratios of 10^{-13} – 10^{-12} at $2 \times 10^4 M_\odot$ — probing a regime in which even Sgr A* would be conspicuous (Event Horizon Telescope Collaboration, 2022).

Two-epoch variability. Differential photometry between epochs is robust to the static crowding systematics that dominate the absolute limits; variability at the ≥ 20 –30 per cent level for any source above ~ 30 nJy flags an accretion candidate and triggers the coordination protocol of Section 11. Two-epoch photometry should not be mistaken for a test of Paper E’s regulation statistic R , the variability *deficit* in the MAD flux-eruption band (10^{-3} – 10^{-1} Hz for a $2 \times 10^4 M_\odot$ hole; cycle times of 10–100 s), which requires fast-cadence radio or X-ray monitoring (Swanson, 2026a). Accordingly, if accretion is ever detected in any band, a fast-cadence contingency activates: sub-second-binned Program-2 continuum monitoring plus the archival X-ray timing of Program 8, targeted at the eruption band.

Waste-heat analysis (archival + new). The 10–30 μ m field photometry is fitted against the synthetic stellar-population SED constructed from the oMEGACat spectroscopic catalogue (Nitschai et al., 2023); the technosignature statistic is a spatially coherent excess in the central 0.1 pc exceeding 3σ over the population model. The deliverable is quantitative through Paper E’s waste-heat floor, $L_{\text{waste}} \geq 10^{-4} P_{\text{comp}}$: the current JWST-derived limit converts to a ceiling on sustained computational power of $P_{\text{comp}} \lesssim 10^3$ – $10^4 L_\odot$, and the ceiling tightens linearly with mid-infrared depth (Swanson, 2026a); the 3σ spatial-coherence excess remains the anomaly trigger. The same fit yields conventional science: the most precise mid-IR census of the post-main-sequence population in any globular cluster.

Spectroscopic environment test (Cycle 6+, contingent). If astrometry or timing strengthens the IMBH case, a NIRSspec IFU mosaic of the inner 0.5 pc (~ 125 h) maps abundance anomalies (Li enhancement, refractory depletion) that would discriminate between ordinary tidal-disruption debris chemistry (Hills, 1988; Gezari, 2021) and the engineered-feeding residue of H_2 . We do not request this time until the trigger condition is met.

3.3 Decision criteria

Null result (expected under H_0 , H_1 , and mature- H_2 alike): bolometric limit $\lesssim 10^{30}$ erg s^{−1} recorded (radiative output $\lesssim 10^{-12}$ of Eddington across the contested mass range), complementing the radio Bondi-efficiency bound and jointly constraining any radiatively inefficient flow via the fundamental plane (Merloni et al., 2003; Mahida et al., 2026). A static detection consistent with a quiescent accretion SED supports H_0 and immediately sharpens every dynamical program (the source position becomes the kinematic centre). A variable detection triggers multi-messenger follow-up. A $> 3\sigma$ extended mid-IR excess with no stellar counterpart is the single strongest electromagnetic anomaly the campaign can produce and would be adjudicated under Section 11 rules.

4 Program 2: Radio — deep imaging, narrowband SETI, and pulsar timing

4.1 Science case

The radio program carries three loads at once. (i) *Continuum*: a deeper interferometric limit (or first detection) on the central compact source, extending Mahida et al. (2026) and the globular-cluster IMBH radio campaigns (Strader et al., 2012; Tremou et al., 2018). (ii) *Narrowband SETI*: to our knowledge the first dedicated technosignature search of any kind at ω Cen, closing the gap left by the FAST pilot survey’s latitude limit (Huang et al., 2026b). (iii) *Pulsar timing*: the binding constraint on the central mass comes

from the cluster’s MSP population (five discovered with Parkes, Dai et al. 2020, timed to $\dot{\nu}$ within 3.5 yr, Dai et al. 2023; then thirteen more with MeerKAT/TRAPUM, Chen et al. 2023; Stappers and Kramer 2016, for eighteen total). The joint MeerKAT–Parkes timing analysis now extends the census and the constraints: its measurements are insensitive to an IMBH of 10^3 – $10^4 M_\odot$ and place a 90 per cent upper limit of $< 10^5 M_\odot$ on the central mass (Colomé Bernadich et al., 2026). The path to resolving the mass tension runs directly through more pulsars and longer baselines (Bañares-Hernández et al., 2025; Chen et al., 2025b).

4.2 Implementation: MeerKAT era (2026–2030)

Timing. Bi-weekly L-band sessions (~ 26 per year, ~ 3 – 5 h each) on all eighteen MSPs with MeerKAT (Jonas and MeerKAT Team, 2016), producing times of arrival at $\sim 1 \mu\text{s}$ precision for the brightest objects. Line-of-sight acceleration precision from timing scales steeply with baseline ($\sigma_a \propto T^{-5/2}$ for white noise); by analogy with the mature 47 Tuc and Terzan 5 programs (Freire et al., 2017; Prager et al., 2017), five years of MeerKAT data yield per-pulsar acceleration uncertainties of order 10^{-11} – $10^{-10} \text{ m s}^{-2}$ for the best-timed objects. For scale, a $4 \times 10^4 M_\odot$ point mass imposes $a \approx 6 \times 10^{-8} \text{ m s}^{-2}$ at $r = 0.3 \text{ pc}$: the discriminating signal is not the detection of acceleration (already achieved; Dai et al. 2023; Bañares-Hernández et al. 2025) but the radial profile of accelerations and jerks across the MSP population, which distinguishes a point mass ($a \propto r^{-2}$) from an extended remnant component. Commensal search observations are expected to add 5–10 MSPs (the luminosity function is far from exhausted; Chen et al. 2023), and the population-level point-mass upper bound tightens approximately as $1/\sqrt{N_{\text{psr}}} (T_0/T)^{\geq 1}$; this is an ensemble, bound-level scaling, distinct from the per-pulsar white-noise acceleration scaling $\sigma_a \propto T^{-5/2}$ quoted above, because the ensemble constraint is limited by the pulsars’ unknown line-of-sight positions and intrinsic spin-down as well as by timing noise. Ten well-timed pulsars with ten-year baselines push a $6,000 M_\odot$ bound to the $3,000$ – $4,000 M_\odot$ level if no point-mass signal emerges, or find one.

Narrowband SETI. Commensally with every timing session, a 1-Hz-resolution spectrometer backend records the full primary beam ($\sim 1^\circ$, comfortably containing the whole cluster). Processing follows standard drift-search practice (Enriquez et al., 2017): Doppler drifts $\pm 4 \text{ Hz s}^{-1}$, SNR > 15 candidate threshold, on–off cadence against reference pointings, and re-detection in an independent session required before any candidate advances. Seventy hours over two years reach a minimum detectable EIRP of $\sim (3\text{--}6) \times 10^{17} \text{ W}$ at 5.49 kpc. This is roughly an order of magnitude above the FAST pilot’s $\sim 10^{16} \text{ W}$ thresholds at its closer, northern targets (Huang et al., 2026b), but it is the first limit of any kind for this cluster, and it is sufficient to detect any transmitter exceeding $\sim 10^4$ times the EIRP of the Arecibo planetary radar (a modest output for an energy-rich civilization) from any of $\sim 10^7$ stars in the beam simultaneously.

Continuum. The summed continuum visibilities from the timing campaign provide a ~ 100 + h synthesis image at L/S band; combined with the ATCA 7.25-GHz limit (Mahida et al., 2026) this constrains both the flat-spectrum (jet) and steep-spectrum (pulsar) interpretations of any future central source candidate.

4.3 Implementation: SKA era (2029–)

SKA-Mid science verification is expected in 2029 and early operational cycles in the early 2030s (Braun et al., 2019). The ω Cen program transfers wholesale: timing precision improves by the sensitivity ratio (factor ~ 4 – 5 over MeerKAT for these declinations), the MSP census plausibly doubles, and the narrowband EIRP threshold drops toward $\sim 10^{16} \text{ W}$, reaching FAST-class sensitivity on a southern target

FAST cannot see. By LISA launch, the pulsar acceleration map will be the best non-GW constraint on the central potential in any globular cluster.

4.4 Decision criteria

The timing program is the campaign’s primary near-term discriminator between H_0 and H_1 : a smooth r^{-2} acceleration profile centred on the kinematic centre at $\geq 10^4 M_\odot$ retires H_1 ; a persistently tightening upper bound below the kinematic lower bounds forces revision of the fast-star analysis (membership, foreground, or binarity systematics) and shifts the campaign’s centre of gravity to H_1 . For SETI: any re-detected narrowband candidate enters the two-messenger protocol; a null is published as the first ω Cen technosignature limit. Paper A dependence (labelled): under H_2 , P3 of that paper predicts no leakage radiation, so SETI nulls carry no evidential weight against H_2 ; they constrain only conventional beacon scenarios. Consistent with this, Paper E’s Bayes-factor analysis finds that additional radio depth beyond current limits moves $\ln K$ only weakly for every hypothesis pairing (Swanson, 2026a); the case for the deeper continuum limit therefore rests on the H_0/H_1 accretion physics, not on technosignature discrimination.

5 Program 3: Astrometry — HST, Gaia DR4, Roman, and ELT/MICADO

5.1 Sensitivity of the direct acceleration test

The intuitively obvious astrometric test (watch the seven fast stars curve) does not work on a useful timescale, which we now demonstrate quantitatively. The proper-motion acceleration induced by a central mass $M_\bullet$ at projected radius r is

$$a = \frac{GM_\bullet}{r^2} \approx 4.4 \times 10^{-7} \left(\frac{M_\bullet}{2 \times 10^4 M_\odot} \right) \left(\frac{r}{0.08 \text{ pc}} \right)^{-2} \text{ m s}^{-2}, \quad (1)$$

which at $d = 5.49$ kpc corresponds to $\approx 5.3 \times 10^{-4}$ mas yr $^{-2}$ (mean projection factors of order unity absorbed). Against this, the oMEGACat astrometry (~ 0.02 mas single-epoch precision over a ~ 15 -year HST baseline) delivers acceleration uncertainties of $\sigma_a \approx 1 \times 10^{-3}$ mas yr $^{-2}$ per star (Häberle et al., 2024b,a): a 0.5σ measurement. Extending HST monitoring to 2028 (a 26-year baseline) improves this only to $\sim 0.7\sigma$; a four-year ELT/MICADO campaign at ~ 50 – $100 \mu\text{as}$ per epoch (Davies et al., 2021) reaches $\sigma_a \approx 2 \times 10^{-3}$ mas yr $^{-2}$: worse, because acceleration precision scales as T^{-2} and four years is short (Figure 2). Direct 5σ curvature detection at nominal parameters requires either $M_\bullet \gtrsim 10^5 M_\odot$, a star at $r < 0.015$ pc, or $\gtrsim 30$ -year baselines on ELT-class astrometry. We therefore retain the fast-star monitoring (cheap; protects against the lucky cases, since González Prieto et al. 2025-mass holes and undiscovered inner stars are both live possibilities) but route the decision-grade astrometry through three measurements that do work, below.

5.2 Three astrometric measurements that do work

(i) **Photocentric wander (Roman).** An IMBH of mass $M_\bullet$ in dynamical equilibrium with stars of mean mass $\bar{m}$ executes Brownian motion with velocity $\sigma_\bullet \sim \sigma_* \sqrt{\bar{m}/M_\bullet}$ relative to the cluster centre of mass. The lighter the hole, the larger its wander: the predicted amplitude differs by ~ 17 per cent between $6,000$ and $8,200 M_\odot$ (the square root of the mass ratio), and by a factor ~ 2.4 between $8,200$ and $4.9 \times 10^4 M_\odot$. Resolving it requires measuring the reflex motion of the innermost stellar distribution at the few- μas yr $^{-1}$ level against the bulk cluster frame: the regime of Roman’s wide-field astrometry,




Figure 2: Why direct acceleration detection on the known fast stars is marginal: 1σ acceleration uncertainty versus baseline for HST-class and ELT-class astrometric campaigns (curves, $\sigma_a \propto \sigma_{\text{pos}} T^{-2}$, anchored to the measured oMEGACat performance and nominal MICADO performance), against the expected signals (horizontal lines, Eq. 1). Curves assume uniform observing cadence over the full baseline and are therefore optimistic for sparse epoch-limited extensions (the realistic 2028 HST extension yields only $\sim 0.7\sigma$). Even the heavy González Prieto et al. (2025) mass crosses 5σ only for baselines approaching two decades. The astrometric program is therefore built around wander, reference-frame, and discovery measurements instead (Section 5.2).

which delivers $\sim 10 \mu\text{as}$ -class differential astrometry over fields vastly larger than HST’s, tying the inner arcseconds to thousands of cluster reference stars at once (WFIRST Astrometry Working Group et al., 2019). Roman’s launch is scheduled for late 2026, with the agency commitment extending to 2027; a five-year cadenced guest-observer program (ω Cen lies outside the core community surveys, so epochs must be requested, but the per-epoch cost is small) delivers, as its headline result, the factor-2.4 discrimination between the light-IMBH and heavy-IMBH wander regimes, independently of individual-star orbits. In the measurement-noise-limited case the program would do better: distinguishing the 17 per cent light-IMBH amplitude difference requires the wander amplitude itself measured to $\lesssim 6$ per cent, which is a reference-frame rather than photon-noise problem, and with $\gtrsim 10^3$ cluster reference stars per field tying the inner-arcsecond frame at each of $\gtrsim 10$ epochs the per-epoch frame error averages well below the expected few- $\mu\text{as yr}^{-1}$ wander signal, in principle supporting $3\text{--}5\sigma$ discrimination within the light-IMBH regime. We do not headline that figure, because the discrimination is limited by the N -body modelling of the stellar background, a systematic that remains unquantified; the factor-2.4 light-versus-heavy separation is what survives a several-fold error inflation, and it is the deliverable we pre-register.

The astrometric channel itself now carries a proof of capability: Whitaker et al. (2026) used the same HST astrometric archive (2002–2023), extended with JWST imaging, to detect the reflex wobble of a main-sequence star orbiting an unseen $4.46^{+1.22}_{-1.01} M_{\odot}$ companion, the first stellar-mass black hole confirmed in ω Cen and the first astrometric black-hole detection in any globular cluster. A dynamically measured dark remnant in the same field, with a 94-yr period and 31-AU separation recovered from curvature over a two-decade baseline, demonstrates end-to-end that the catalogue’s astrometry supports orbit-grade inference at the precision this program assumes.

(ii) The absolute reference frame (Gaia DR4, 2026 December). The dominant systematic in all current inner-field astrometry is the tie between HST’s relative frame and the absolute (ICRS) frame. Gaia DR4’s 5.5-year solutions for the $r > 10''$ cluster members (Gaia Collaboration et al., 2016) improve proper motions by a factor ~ 4.5 over DR2-era ties, propagating directly into the fast-star velocity vectors — and

thus into the $8,200 M_{\odot}$ lower bound itself, which rests on those velocities exceeding the escape speed. This is the cheapest decisive measurement in the entire campaign: a re-derivation of the Häberle et al. (2024b) bound on the DR4 frame either firms the tension into a $> 5\sigma$ contradiction with the timing bound or dissolves it. Deliverable in 2027 from archival data alone.

(iii) **Inner-star discovery (ELT/MICADO, 2029+).** MICADO’s ~ 10 mas resolution and 39-m aperture (Davies et al., 2021) penetrate the central arcsecond at magnitudes HST cannot reach in crowding, with the realistic prize being new fast stars at $r < 0.02$ pc. A single star at 0.01 pc raises the acceleration signal of Eq. (1) by a factor of 64, converting Figure 2’s verdict from “decades” to “a few epochs” — this is how the Galactic-Centre program succeeded (GRAVITY Collaboration et al., 2018), and ω Cen is its natural second act at $10^2\times$ lower mass. Two epochs per year from 2029; first acceleration-grade results plausible by the early 2030s.

5.3 Decision criteria

DR4 re-derivation (2027): if the fast-star velocity excess survives at $> 5\sigma$, H_1 requires the MSP analysis to be systematically biased — a specific, checkable claim (radial distribution priors; Bañares-Hernández et al. 2025). Roman wander (2031): light/heavy discrimination feeds the LISA prior. MICADO inner stars (2030s): any star with measured Keplerian curvature yields $M_{\bullet}$ to tens of per cent, independent of statistical modelling, before LISA flies.

6 Program 4: The LISA forecast

6.1 Science case

LISA is the campaign’s endgame: the only instrument that can measure the central object’s mass and spin to high precision regardless of its electromagnetic state (Colpi et al., 2024; Amaro-Seoane et al., 2017). For a compact object inspiralling into a 10^4 -class IMBH, matched-filter parameter estimation delivers fractional precisions of order 10^{-3} – 10^{-4} on mass and 10^{-3} on spin (Babak et al., 2017; Amaro-Seoane, 2018). At 5.49 kpc (versus the Gpc distances over which LISA expects to detect its EMRI population), any ω Cen inspiral in band during the mission would be loud (signal-to-noise in the hundreds to thousands), making occurrence probability, not sensitivity, the limiting factor.

6.2 Occurrence rates

Event-rate estimates in the LISA literature imply IMRI formation rates per cluster of order 10^{-9} – 10^{-6} yr $^{-1}$ depending on IMBH mass, binary fraction, and core density (Mandel et al., 2008; Amaro-Seoane, 2018; Fragione et al., 2018); over a 4–10-year mission this yields an in-band inspiral probability for ω Cen of $\lesssim 10^{-5}$, possibly enhanced by the cluster’s unusually dense remnant population, but not by the four to five orders of magnitude required to make it likely. The forecast must therefore be stated in three tiers. **Tier 1 (probable):** no ω Cen source; LISA still constrains the stochastic foreground from the core remnant population, directly testing H_1 ’s extended-component density profile. **Tier 2 (plausible):** a quasi-monochromatic source (a compact object orbiting the IMBH well outside plunge) detectable over the mission as a continuous-wave source; this measures $M_{\bullet}$ through the orbital frequency and its drift, to precision intermediate between timing and a full inspiral. **Tier 3 (lucky):** a genuine inspiral; per-mille mass and spin. Even in Tier 1, a spin constraint for the class arrives statistically: LISA’s detected population of comparable IMBH systems at cosmological distances calibrates the natural spin distribution against which any eventual ω Cen measurement is judged. The campaign’s structure ensures that even

Tier 1 is decisive when combined with the pulsar acceleration map (Section 4): a stochastic-foreground non-detection plus a point-mass timing profile isolates H_0 .

6.3 The spin measurement as the H_2 adjudicator

Paper A dependence (labelled): under H_2 the single sharpest residue is near-extremal spin ($a_\star \gtrsim 0.9$) on a hole that has demonstrably accreted nothing for Gyr; under H_0 natural formation channels for cluster IMBHs span low-to-moderate spin. A Tier 3 spin measurement therefore adjudicates H_2 at the same instant it completes the H_0/H_1 question — the entire reason this campaign’s electromagnetic and dynamical groundwork must precede LISA rather than follow it. The full decision tree, including this branch, is assembled in Section 13.

6.4 Deliverables before launch

Between now and the mid-2030s the program’s work is preparatory but concrete: (i) maintain the dynamical model of the inner parsec (timing + astrometry) so that any LISA source has an immediate host context; (ii) publish the ω Cen-specific EMRI/IMRI rate forecast with the post-2026 mass constraints folded in; (iii) ensure the cluster’s barycentric ephemeris and distance (± 0.06 kpc; Häberle et al. 2025) are maintained at the precision LISA parameter estimation will assume.

7 Program 5: KM3NeT/ARCA neutrino monitoring

7.1 Science case and geometry

ω Cen at declination -47.5° is an up-going source for the Mediterranean: KM3NeT/ARCA observes it through the Earth, with atmospheric muons filtered out and sub- 0.2° angular resolution for track events at $\gtrsim 10$ TeV (Adrián-Martínez et al., 2016). For IceCube the same source is down-going (degraded but usable above ~ 100 TeV; Aartsen et al. 2020); the combined ANTARES+IceCube southern-sky analysis (Albert et al., 2020) defines the archival baseline. ARCA stood at 51 of 230 detection units as of early 2026, with real-time alert distribution entering commissioning; sensitivity grows roughly linearly with instrumented volume through the decade. The partially built array has already delivered a discovery-class result: the ~ 220 -PeV track event KM3-230213A, the most energetic neutrino yet observed (The KM3NeT Collaboration, 2025), demonstrates that ARCA detects and reconstructs the rare high-energy events on which this program’s burst criterion rests. The conventional science driver is a deep point-source limit on a dense old stellar system; the technosignature driver (Paper A dependence, labelled) is the burst-mode prediction of black-hole-based computation models (Dvali and Osmanov, 2023; Lacki and DiKerby, 2025), for which a cluster like ω Cen is the natural test bed and which the existing point-source pipelines do not target: their time-integrated statistics dilute rare short multiplets.

7.2 Implementation

Steady-state search (archival + ongoing). Standard unbinned-likelihood point-source analysis (Braun et al., 2008) at the ω Cen coordinates in (i) the ANTARES+IceCube combined dataset, (ii) the growing ARCA exposure. Full-array ARCA reaches $E^2\Phi \sim \text{few} \times 10^{-12} - 10^{-11}$ TeV cm $^{-2}$ s $^{-1}$ at this declination over multi-year integrations — the deepest neutrino limit ever placed on a globular cluster.

Burst pipeline (new, the campaign’s contribution). A pre-registered transient search in sliding windows of 10^2 , 10^3 , and 10^4 s within 1° of the cluster centre, $E \gtrsim 10$ TeV, on ARCA data plus IceCube




Figure 3: Design of the neutrino burst search. Expected atmospheric-neutrino background within 1° of ω Cen versus window length, for the working $E > 10$ TeV threshold (solid) and a softer 1 TeV threshold (dashed). At 10^3 s the expectation is $\sim 10^{-4}$ events, so the pre-registered triplet criterion is intrinsically secure against background even after a decade-scale trials budget; rate normalizations are order-of-magnitude, with exact values set by the detector Monte Carlo at analysis time.

alerts. The detection criterion (≥ 3 tracks in one window, $\geq 5\sigma$ post-trials, no plausible astrophysical counterpart) is calibrated by the background expectation: the atmospheric-neutrino rate above 10 TeV within a 1° cone is taken as 10^{-7} s^{-1} ($\sim 3 \text{ events yr}^{-1}$ in the cone), a deliberately conservative envelope that likely overstates the measured up-going rate for a cone this size by one to two orders of magnitude; even so, a 10^3 -s window expects only $\sim 10^{-4}$ events and a triplet is intrinsically far beyond 5σ *before* trials; the design problem is the trials budget over a decade of monitoring ($\sim 3 \times 10^6$ windows), which the post-trials threshold absorbs (Figure 3). Any KM3NeT or IceCube real-time alert within 1° triggers the electromagnetic protocol of Section 11 — the alert system’s first ω Cen-relevant cycle begins in late 2026.

7.3 Decision criteria

A decade-scale null at full-array sensitivity closes the burst-mode channel of Dvali and Osmanov (2023) for this system at its predicted strength — a publishable falsification regardless of H_2 (the underlying micro-black-hole physics is itself contested; Álvarez-Domínguez et al. 2024; Loeb 2024). One confirmed multiplet with no counterpart, repeated at the same location, would be among the most significant anomalies in high-energy astrophysics independent of any ETI interpretation; the protocol deliberately treats it as an astrophysical discovery first.

8 Program 6: Gamma rays — Fermi-LAT archival and CTAO-South

8.1 Science case

ω Cen is a GeV source: 4FGL J1326.7–4729 coincides with the core, and its emission is attributed to the MSP ensemble after deep searches found no individual pulsations (Dai et al., 2020, 2023). This is both foreground and opportunity: the steady MSP emission is a calibration source against which *transient* or *spectrally anomalous* excesses can be sought with high contrast. The technosignature channel shares the burst phenomenology of Program 5 (the Dvali and Osmanov 2023 spectrum is “democratic” across

species, implying γ -ray counterparts to any neutrino burst); the conventional channel is a TeV detection or limit on the cluster, of independent interest for MSP magnetospheric and inverse-Compton physics.

8.2 Implementation

Fermi-LAT (archival, immediate). Re-analysis of the now >17-year LAT exposure at the cluster position: (i) refit of the steady source with current diffuse models; (ii) a sliding-window transient search ($\Delta t = 10^2\text{--}10^4$ s, matched to Program 5’s windows) over the full mission, with the steady MSP flux as null model; (iii) spectral-cutoff analysis: MSP ensembles cut off at a few GeV, so any significant emission above ~ 30 GeV flags a non-MSP component. All three are pure archival analyses.

CTAO-South (2027+). The first Paranal telescopes are, on early-2026 construction schedules, arriving through late 2026 (Cherenkov Telescope Array Consortium et al., 2019); intermediate arrays operate while construction proceeds. We propose (i) a 50-h deep exposure reaching $\sim 10^{-13}$ erg cm $^{-2}$ s $^{-1}$ at 1 TeV for the completed Alpha-configuration array; the 2027-era partial array delivers a factor 2–4 less, and the depth quoted here is recovered as construction completes. Either figure substantially better the archival VHE limits placed on globular clusters by the H.E.S.S. cluster survey, making this the first CTAO-depth TeV limit on any globular cluster core. We also propose (ii) a standing target-of-opportunity trigger: any Program-5 alert within 1° pre-authorizes prompt CTAO follow-up, the latency being minutes for a source that transits Paranal $\sim 23^\circ$ from zenith.

8.3 Decision criteria

A LAT transient coincident with a neutrino window elevates any Program-5 candidate to the campaign’s highest tier. A TeV detection of steady emission is conventional astrophysics: it would make ω Cen the second globular cluster with detected VHE emission, after H.E.S.S.’s detection coincident with Terzan 5 (Abramowski et al., 2011), and the first with a resolved connection to a candidate IMBH environment. Its absence at CTAO depth tightens the inverse-Compton budget of the MSP population well beyond the existing H.E.S.S.-era cluster limits. Spectral excess above 30 GeV without transient behaviour prompts a dark-matter-versus-exotica analysis before any H_2 language is entertained.

9 Program 7: Optical/IR time domain — Rubin/LSST and archival anomalies

9.1 Science case

Two distinct searches share this program. First, transients: tidal disruption of a star by an IMBH is the one natural event that would announce the central object unambiguously (Hills, 1988; Gezari, 2021); rates are low ($\sim 10^{-7}\text{--}10^{-6}$ yr $^{-1}$ per cluster for full disruptions, higher for partial-disruption flares of the dense remnant environment) but the cost of monitoring is now effectively zero, because LSST began surveying in early 2026 and ω Cen sits in the survey footprint (Ivezić et al., 2019). Second, archival photometric anomalies: the oMEGACat HST archive (500+ epochs over 20+ years for 10^6 stars; Häberle et al. 2024a; Nitschai et al. 2023) is an essentially unmined time-domain dataset in which periodic dimmings, secular fades, or statistically anomalous variability classes (the photometric residues conjectured for megastructures; Wright et al. 2014; Hsiao et al. 2021; Socas-Navarro et al. 2021) can be sought with two decades of baseline for free.

9.2 Implementation

LSST stream and bright-transient coverage. At the cluster’s distance modulus of $\mu = 13.7$, a full TDE peaking at $M_V \approx -14$ to -17 would appear at $V \approx 0$ to -3 : hopelessly saturated for LSST difference imaging, and indeed for any survey-class imager. The bright-event trigger therefore belongs to ASAS-SN/Evryscope-class all-sky photometry, which monitors this magnitude range continuously; the campaign’s role is to register the ω Cen position with those pipelines. LSST covers the complementary faint regime: a standing broker filter on the field flags (i) any nuclear transient in the unsaturated range $r \gtrsim 16$, which captures partial disruptions and other sub-luminous accretion flares, and (ii) anomalous variables among $\sim 10^5$ measurable members outside the saturated core. Difference-imaging in the crowded annulus is handled with the oMEGACat reference catalogue.

Archival mining. A systematic variability census of the oMEGACat photometric archive: period search to ~ 0.05 mag depth across 10^6 stars, secular-trend extraction, and an explicitly pre-registered anomaly statistic (population-level excess of non-physical light-curve classes) with human vetting of survivors. Conventional yield: the deepest variable-star census of any globular cluster, eclipsing-binary distance cross-checks, and stellar-rotation demographics — publishable regardless of anomalies.

Core-depletion statistic (joint with Program 3). Paper A dependence (labelled): H_2 ’s P4c predicts secular depletion of the loosely bound core population beyond N -body expectations. The measurement (star counts by binding-energy quantile against matched N -body models; [González Prieto et al. 2025](#)) is identical to standard mass-segregation analysis and is conventional dynamics on its face. We track the depletion direction as an exploratory statistic rather than a pre-registered one: at present it has a predicted sign but no quantified effect size or H_0/H_1 envelope definition, and elevating it to the pre-registered tier must await the matched N -body envelope work.

9.3 Decision criteria

A nuclear TDE settles the IMBH question electromagnetically (and its decay light curve constrains $M_\bullet$; [Gezari 2021](#)); the campaign’s role is to guarantee the multi-wavelength response is pre-arranged. Archival anomalies are population statistics: no single weird star means anything (the false-positive history of photometric SETI is long); only a $\geq 3\sigma$ class-level excess, surviving vetting, enters the coordination protocol.

10 Program 8: Supplementary archival channels

Three channels are cheap enough to run as graduate-student archival projects and complete the messenger coverage.

Archival X-rays (Chandra, XMM-Newton, eROSITA). The X-ray band is the standard probe of quiescent black-hole accretion, and the campaign as presented so far would otherwise lack it. The benchmark limit is the ~ 291 ks Chandra exposure of [Haggard et al. \(2013\)](#), which detects no central point source and bounds the unabsorbed luminosity at $L_X(0.5\text{--}7.0 \text{ keV}) \lesssim 1.6 \times 10^{30} \text{ erg s}^{-1}$, an Eddington ratio of $\lesssim 10^{-12}$ for a $10^4 M_\odot$ hole. The archival program re-derives this limit at the current oMEGACat kinematic centre (the 2013 analysis predates the fast-star centre determination), stacks the subsequent Chandra and XMM-Newton exposures of the field, and folds in the eROSITA all-sky survey epochs as a variability screen; a joint X-ray–radio treatment through the fundamental plane ([Merloni et al., 2003](#)) then constrains radiatively inefficient flow models alongside the Program 1 and Program 2 limits. Any X-ray point source at the kinematic centre, at any flux, would be a major result; the expected null sharpens the deepest multi-band quiescence measurement for any black hole candidate.

Continuous gravitational waves (LIGO–Virgo–KAGRA). Narrowband searches at twice the spin frequencies of the eighteen known MSPs, using public O4/O5 data and the MeerKAT ephemerides, following the known-pulsar methodology of [Abbott et al. \(2022\)](#). Expected upper limits $h_0 \lesssim 10^{-26}$ translate to ellipticity limits $\varepsilon \lesssim \text{few} \times 10^{-8}$ at 5.49 kpc for the faster spinners — conventional neutron-star physics in an environment (a dense old cluster) where the recycled-pulsar population is dynamically distinctive. The technosignature reading (structured masses in MSP orbits) is not separately funded; it shares the identical data product.

Ultra-high-energy cosmic rays (Pierre Auger). ω Cen sits well inside Auger’s southern acceptance. A stacked directional analysis around the cluster position in the public $E > 8$ EeV dataset ([Pierre Auger Collaboration et al., 2017](#)), with 1° and 3° apertures bracketing magnetic deflection, either contributes an upper limit on hadronic acceleration in the core (of conventional interest given the MSP wind population) or flags an anisotropy for which the Galactic-Centre region (18° away) is the obvious confounder to exclude first.

11 Coordination, joint statistics, and data policy

11.1 The two-messenger rule, operationalized

Table 3 pre-registers the campaign’s trigger and confirmation matrix. Each row names a primary trigger, the channels that can confirm it, and the joint-significance requirement. Combined significances across independent channels use Fisher’s method on the per-channel p -values, with the global claim threshold set at $p < 5 \times 10^{-7}$ post-trials — deliberately at discovery convention, because the base rate of true technosignatures is unknown and plausibly zero, so the loss function is asymmetric: a false positive damages the field’s credibility far more than a delayed true positive damages the discovery ([Wright et al., 2022](#)).

This matrix is one of two pre-registrations governing the campaign, and the two occupy different layers. The Fisher-combined $p < 5 \times 10^{-7}$ matrix of Table 3 is the *trigger layer*: it decides when an event or excess is promoted from a single-channel candidate to a campaign-level anomaly requiring adjudication. Paper E’s hierarchical Bayes-factor framework, with Kass–Raftery $\ln K$ action bands whose “candidate” band itself requires evidence in at least two messengers, is the *inference layer*: it takes any promoted trigger and adjudicates it against the full $H_q/H_{\text{sub}}/H_{\text{eng}}$ hypothesis set ([Swanson, 2026a](#)). The frequentist matrix therefore controls the false-alarm budget of what enters adjudication, while the Bayesian framework determines what the adjudicated evidence supports; a trigger that clears Table 3 but lands in a weak $\ln K$ band is reported as an unexplained astrophysical candidate, not an anomaly claim.

11.2 Alert latency and standing authorizations

The latency-critical path is neutrino $\rightarrow$ gamma/optical: KM3NeT real-time alerts (commissioning through 2026) distribute within seconds–minutes; the CTAO ToO and LSST-broker hooks are standing authorizations requiring no human decision for the first response hour. Slow-path coordination (JWST DDT requests, radio re-pointing) is pre-drafted as template proposals with trigger criteria attached, cutting submission latency to < 24 h.

11.3 Data policy

All pipelines, thresholds, and analysis code are public from the start (the pre-registration is itself the methods paper); all derived catalogues are released with their papers; raw data inherit host-facility policies.

Table 3: Pre-registered trigger and confirmation matrix. “Joint” = Fisher-combined post-trials significance required for any anomaly claim; single-channel events, however significant, are reported as astrophysical candidates only.

Primary trigger	Channel threshold	Confirming channels	Disposition
Neutrino multiplet (P5)	≥ 3 tracks / 1° / 10^{2-3} s / ≥ 10 TeV, 5σ post-trials	Fermi-LAT window (P6); CTAO ToO (P6); JWST/ground IR (P1)	repeat at same position required for anomaly claim; single event published as astrophysical transient
Narrowband radio line (P2)	SNR > 15 , re-detected in independent session	off-source cadence; second telescope (e.g. ATCA)	interference exclusion protocol precedes any claim
IR variable/transient (P1, P7)	$\geq 20\%$ amplitude or new source $> 5\sigma$	radio continuum (P2); X-ray archival; LSST stream (P7)	classified as accretion/TDE candidate first
Photometric anomaly class (P7)	$\geq 3\sigma$ population-level excess	independent archive (Gaia epochs); spectroscopy	population statistic only; no single-star claims
Astrometric anomaly (P3)	wander outside H_0/H_1 envelopes at 3σ (core-depletion statistic exploratory only; Section 9)	timing acceleration map (P2)	feeds LISA prior; no standalone claim
>30 GeV spectral excess (P6)	$\geq 5\sigma$ non-MSP component in LAT refit	CTAO follow-up (P6); radio/X-ray positional coincidence (P2, P8)	dark-matter-versus-exotica analysis precedes any anomaly language
CW detection at MSP frequency (P8)	$\geq 5\sigma$ in known-pulsar narrowband search	EM timing solution and position (P2)	classified as neutron-star physics first
UHECR anisotropy (P8)	$\geq 3\sigma$ post-trials stacked excess, 1° – 3° apertures	Fermi-LAT/CTAO coincidence (P6); Galactic-Centre confounder excluded	directional statistic only; no source claim without γ -ray support
X-ray point source at centre (P8)	any source $> 3\sigma$ at kinematic centre	radio continuum (P2); JWST IR counterpart (P1)	classified as accretion candidate; feeds fundamental-plane fit
LISA spin/mass result (P4)	per mission parameter estimation	all of the above as context	adjudicates $H_0/H_1/H_2$ per Section 13

Negative results are published on a fixed, pre-announced cadence, so that the decision to publish never depends on what the data showed.

11.4 Program summary

Table 4 assembles the eight programs with sensitivities, requests, and timelines.

12 Cost and timeline

12.1 Costing

Table 5 presents full-cost estimates (salaries, computing, travel, publication; US academic rates) for each program. Facility time is excluded (it is competitively allocated and carries no cash cost to the campaign), as is LISA, which is mission science. The total, $\lesssim$ US\$7M over a decade, is calibrated against the per-program budgets developed in the precursor proposal studies for this campaign and is small by every relevant comparison: a single mid-scale NASA Explorer instrument, one JWST cycle’s archival-funding pool, or the marginal cost of a few nights of ELT operations.

12.2 Timeline

Figure 4 lays the programs against the facility schedule. The structure is deliberate: the 2026–2028 window is dominated by archival work and standing-pipeline construction (cheap, immediate); 2029–2034

Table 4: Campaign summary. “Type”: A = archival/commensal, N = new observations, F = free by-product of funded survey. Costs are rough full-cost estimates (Section 12); facility time is listed separately since it is allocated, not purchased.

#	Program	Type	Key sensitivity	Facility request	Timeline	Decides
P1	JWST IR limits	N	~ 10 nJy (F444W); $L_{\text{bol}} \lesssim 10^{30}$ erg s $^{-1}$	~ 65 h, 2 epochs	2027–2029	accretion state; waste heat
P2	Radio: timing + SETI + continuum	N	$\sigma_a \sim 10^{-11} - 10^{-10}$ m s $^{-2}$ (5 yr); EIRP (3–6) $\times 10^{17}$ W	~ 130 h yr $^{-1}$ MeerKAT, 5 yr	2026–2031+	H_0 vs H_1 (profile); first SETI limit
P3	Astrometry: DR4 + Roman + ELT	A/F/N	frame to DR4; wander factor-2.4 (light vs heavy); inner-star discovery	archival; Roman survey; ~ 28 h ELT	2026–2035	mass tension; wander mass
P4	LISA forecast	F	$\delta M/M \sim 10^{-3-4}$, $\delta a_\star \sim 10^{-3}$ (if inspiral)	none (mission science)	2035+	everything (Tier 3)
P5	Neutrino monitoring	A/N	$E^2 \Phi \sim \text{few} \times 10^{-12}$ TeV cm $^{-2}$ s $^{-1}$; burst triplets	pipeline on ARCA stream	2026–2040	Dvali–Osmanov channel
P6	Gamma rays	A/N	LAT transient windows; 10^{-13} erg cm $^{-2}$ s $^{-1}$ at 1 TeV (50 h)	50 h CTAO + ToO	2026–2032	burst counterparts; first CTAO-depth TeV limit
P7	Time domain	A/F	TDE flares unmissable; 10^6 -star archival census	LSST stream; archival	2026–2036	TDE; anomaly classes
P8	X-ray + CW + UHECR archival	A	$L_X \lesssim 10^{30}$ erg s $^{-1}$; $h_0 \lesssim 10^{-26}$; stacked UHECR limit	public data	2026–2029	completeness

Table 5: Indicative full-cost budget (US\$ k , 2026 dollars). Personnel figures assume postdoc-led programs with PI/Co-I fractions and graduate students where noted. Estimates for the Roman component of P3 and for P5, P6, P7 are new to this paper; others follow the precursor proposal studies.

Program	Cost	Years	Dominant element
P1 JWST deep imaging + waste heat	1,310	3	postdoc + grad + HPC
P1b JWST spectroscopic phase (contingent)	257	2	postdoc (triggered only)
P2 MeerKAT timing + SETI + continuum	2,900	5	2 postdocs + grad + computing
P3 HST/Gaia frame + Roman wander	425	4	postdoc + HPC
P3b ELT/MICADO epochs	490	4	postdoc share + travel
P4 LISA preparatory modelling	150	3	PI/Co-I fractions
P5 Neutrino burst pipeline	250	3	postdoc + compute
P6 Fermi archival + CTAO ToO	200	3	postdoc share
P7 LSST broker + archival mining	300	3	postdoc + broker engineering
P8 X-ray + CW + UHECR archival	239	3	graduate RA + HPC
Coordination, data releases, workshops	150	10	part-time coordinator
Total (P1b excluded)	6,414	—	—




Figure 4: Campaign timeline against the facility schedule, 2026–2040. Decision points: D1 (2027): Gaia-DR4 re-derivation of the fast-star bound. D2 (2030): five-year pulsar acceleration profile; Roman wander first results. D3 (2033): pre-LISA synthesis — H_0 versus H_1 called at available significance. D4 (2037): LISA verdict, if a source is caught.

by the new-facility harvest (ELT, SKA-Mid, CTAO completion, Roman mid-mission); 2035+ by LISA. Decision points D1–D4 are marked and defined in Section 13.

13 The decision structure through 2040

Figure 5 assembles the campaign into a single decision tree, extending the gravitational-wave branch of Paper A’s falsification framework backward into the electromagnetic-and-timing era. Rough branch weights are stated where the inputs exist to estimate them; they are planning priors, not forecasts, and we expect D1/D2 to revise them substantially.

14 Conclusion

Omega Centauri in 2026 presents a configuration that observational astronomy rarely supplies: a contested discovery (the Galaxy’s best IMBH candidate) whose two strongest constraints formally contradict each other; an electromagnetic silence deep enough to be a result in itself; an essentially untouched technosignature parameter space; and a fleet of new southern and all-sky instruments (Rubin, Roman, Gaia DR4, KM3NeT, CTAO, SKA-Mid, ELT, and ultimately LISA) arriving on the timescale needed to resolve all of it. The campaign presented here is designed to use that conjunction fully.

Its architecture reflects three commitments. First, *dual use*: every program stands as conventional astrophysics (the mass tension, the quiescent-accretion frontier, the MSP dynamical laboratory, the first deep TeV and neutrino limits on a globular cluster), so that no null result is a loss. Second, *conservative sensitivity accounting*: where the obvious measurement fails (direct fast-star acceleration before ~2040), we demonstrate the failure quantitatively and redesign around it, because a program that oversells its tests invites the credibility collapse that has repeatedly damaged this field. Third, *pre-registration*: hypotheses $H_0/H_1/H_2$, thresholds, confirmation rules, and decision points are fixed in advance, so that whatever the




Figure 5: Decision structure through 2040. Red boxes retire hypotheses; the single green branch is the only path on which the technosignature interpretation gains material support, and even there the claim is “anomaly requiring explanation.” Percentages are planning priors only. Every terminal node is a publishable scientific result about a first-rank astrophysical object.

sky delivers (a remnant swarm, a quiet heavy hole, or a genuine anomaly), the inference chain was written down before anyone knew the answer.

The cost is under US\$7M across a decade, most of it salaries for archival analysis riding on facilities that are funded regardless. The payoff structure is asymmetric in the campaign’s favour: the probable outcomes (a resolved mass tension, a characterized central object, the first technosignature limits on 10^7 old stars) are solid science at modest cost, and the improbable outcome (the green branch of Figure 5) would justify the program many thousands of times over. We commit, symmetrically, to the red branches: if the timing profile confirms an extended remnant component, or LISA delivers a light or slowly spinning hole, the anomaly hypotheses retire here without special pleading, and what remains is what was always underneath: the most interesting stellar system in the southern sky, finally measured properly.

Acknowledgements and disclosure

The author thanks the maintainers of the NASA Astrophysics Data System and arXiv, on which the citation verification for this work relied. **AI assistance disclosure:** drafting, citation verification, derivation checking, and figure preparation for this manuscript were performed with substantial assistance from a large language model (Claude, Anthropic), under the author’s direction; the author reviewed and takes full responsibility for all claims, derivations, and references. Interactive calculators implementing the quantitative material in this paper, together with the underlying per-instrument proposal studies, are available at <https://omegacentauri.me>.

Data availability

No new observational data were generated for this work. All quantitative claims derive from the cited literature; analysis conventions for the proposed programs will be released with the respective pipeline papers.

References

- Aartsen, M. G. et al. (2020). Time-integrated neutrino source searches with 10 years of IceCube data. *Physical Review Letters*, 124:051103.
- Abbott, R., Abe, H., Acernese, F., Ackley, K., Adhikari, N., Adhikari, R. X., et al. (2022). Searches for gravitational waves from known pulsars at two harmonics in the second and third LIGO-Virgo observing runs. *The Astrophysical Journal*, 935:1.
- Abramowski, A. et al. (2011). Very-high-energy gamma-ray emission from the direction of the galactic globular cluster terzan 5. *Astronomy & Astrophysics*, 531:L18.
- Adrián-Martínez, S. et al. (2016). Letter of intent for KM3NeT 2.0. *Journal of Physics G: Nuclear and Particle Physics*, 43(8):084001.
- Albert, A., André, M., Anghinolfi, M., Ardid, M., Aubert, J.-J., et al. (2020). ANTARES and IceCube combined search for neutrino point-like and extended sources in the Southern Sky. *The Astrophysical Journal*, 892:92.
- Álvarez-Domínguez, Á., Garay, L. J., Martín-Martínez, E., and Polo-Gómez, J. (2024). No black holes from light. *Physical Review Letters*, 133:041401.

- Amaro-Seoane, P. (2018). Relativistic dynamics and extreme mass ratio inspirals. *Living Reviews in Relativity*, 21:4.
- Amaro-Seoane, P. et al. (2017). Laser interferometer space antenna. LISA L3 mission proposal submitted to ESA.
- Babak, S., Gair, J., Sesana, A., Barausse, E., Sopuerta, C. F., Berry, C. P. L., Berti, E., Amaro-Seoane, P., Petiteau, A., and Klein, A. (2017). Science with the space-based interferometer LISA. V. Extreme mass-ratio inspirals. *Physical Review D*, 95(10):103012.
- Bañares-Hernández, A., Calore, F., Martín Camalich, J., and Read, J. I. (2025). New constraints on the central mass contents of Omega Centauri from combined stellar kinematics and pulsar timing. *Astronomy & Astrophysics*, 693:A104.
- Baumgardt, H. and Vasiliev, E. (2021). Accurate distances to Galactic globular clusters through a combination of Gaia EDR3, HST, and literature data. *Monthly Notices of the Royal Astronomical Society*, 505:5957–5977.
- Braun, J., Dumm, J., De Palma, F., Finley, C., Karle, A., and Montaruli, T. (2008). Methods for point source analysis in high energy neutrino telescopes. *Astroparticle Physics*, 29:299–305.
- Braun, R., Bonaldi, A., Bourke, T., Keane, E., and Wagg, J. (2019). Anticipated performance of the Square Kilometre Array – Phase 1 (SKA1). *arXiv e-prints*.
- Breen, P. G. and Heggie, D. C. (2013). Dynamical evolution of black hole subsystems in idealized star clusters. *Monthly Notices of the Royal Astronomical Society*, 432:2779–2797.
- Chen, S., Hare, J., Kargaltsev, O., Yang, H., Cioffi, D., Häberle, M., and Seth, A. (2025a). The intermediate mass black hole in Omega Centauri: Constraints on accretion from JWST. Submitted to ApJ.
- Chen, W., Freire, P. C. C., Ridolfi, A., Barr, E. D., Stappers, B., Kramer, M., et al. (2023). MeerKAT discovery of 13 new pulsars in Omega Centauri. *Monthly Notices of the Royal Astronomical Society*, 520:3847–3856.
- Chen, X., Vázquez-Aceves, V., Chen, S., Lee, K., Guo, Y., and Liu, K. (2025b). Detecting intermediate-mass black holes using miniature pulsar timing arrays in globular clusters. *arXiv e-prints*, page arXiv:2507.08201.
- Cherenkov Telescope Array Consortium, Acharya, B. S., Agudo, I., Al Samarai, I., et al. (2019). *Science with the Cherenkov Telescope Array*. World Scientific, Singapore.
- Clontz, C., Seth, A. C., Dotter, A., Häberle, M., Neumayer, N., et al. (2024). oMEGACat. IV. Constraining the ages of Omega Centauri subgiant branch stars with HST and MUSE. *The Astrophysical Journal*, 977:14.
- Colomo Bernadich, M., Dai, S., Abbate, F., Kerr, M., Bachetti, M., Buchner, S., Johnston, S., Burgay, M., Possenti, A., Ridolfi, A., Freire, P. C. C., Kramer, M., Stappers, B., et al. (2026). A joint MeerKAT and Parkes view of Omega Centauri: New TRAPUM searches and pulsar timing. *arXiv e-prints*, page arXiv:2603.21845. Under review.
- Colpi, M. et al. (2024). LISA definition study report. ESA-SCI-DIR-RP-002.

- Crawford, I. A. (2026). Some thoughts on the future of technosignature searches: Constraining the Fermi paradox. To appear in Proceedings of IAU Symposium 404.
- Curtis, O., Adams, V. H., Angerhausen, D., Bates, J., Berea, A., Dick, S. J., Elvis, M., Khatri, S. P., Linares, R., Muco, M., Seager, S., and Wright, J. T. (2026). The Dyson Minds 2025 workshop: SETI around black holes. *Publications of the Astronomical Society of the Pacific*, 138:046001.
- Dai, S. et al. (2020). Discovery of millisecond pulsars in the globular cluster Omega Centauri. *The Astrophysical Journal Letters*, 888:L18.
- Dai, S. et al. (2023). Timing of pulsars in the globular cluster Omega Centauri. *Monthly Notices of the Royal Astronomical Society*, 521:2616–2622.
- Davies, R. et al. (2021). MICADO: The multi-adaptive optics camera for deep observations. *The Messenger*, 182:17–21.
- Dvali, G. and Osmanov, Z. N. (2023). Black holes as tools for quantum computing by advanced extraterrestrial civilizations. *International Journal of Astrobiology*, 22(6):617–640.
- Dyson, F. J. (1960). Search for artificial stellar sources of infrared radiation. *Science*, 131(3414):1667–1668.
- Enriquez, J. E., Siemion, A., Foster, G., Gajjar, V., Hellbourg, G., Hickish, J., Isaacson, H., Price, D. C., Croft, S., DeBoer, D., et al. (2017). The Breakthrough Listen search for intelligent life: 1.1–1.9 GHz observations of 692 nearby stars. *The Astrophysical Journal*, 849:104.
- Event Horizon Telescope Collaboration (2022). First Sagittarius A* Event Horizon Telescope results. I. The shadow of the supermassive black hole in the center of the Milky Way. *The Astrophysical Journal Letters*, 930:L12.
- Fragione, G., Leigh, N. W. C., Ginsburg, I., and Kocsis, B. (2018). Tidal disruption events and gravitational waves from intermediate-mass black holes in evolving globular clusters across space and time. *The Astrophysical Journal*, 867:119.
- Freire, P. C. C., Ridolfi, A., Kramer, M., Jordan, C., Manchester, R. N., Torne, P., et al. (2017). Long-term observations of the pulsars in 47 Tucanae – II. Proper motions, accelerations and jerks. *Monthly Notices of the Royal Astronomical Society*, 471:857–876.
- Gaia Collaboration, Prusti, T., de Bruijne, J. H. J., Brown, A. G. A., Vallenari, A., Babusiaux, C., et al. (2016). The Gaia mission. *Astronomy & Astrophysics*, 595:A1.
- Gezari, S. (2021). Tidal disruption events. *Annual Review of Astronomy and Astrophysics*, 59:21–58.
- González Prieto, E., Rodríguez, C. L., and Cabrera, T. (2025). Growing the intermediate-mass black hole in Omega Centauri. *The Astrophysical Journal Letters*, 990:L69.
- GRAVITY Collaboration, Abuter, R., Amorim, A., Anugu, N., Bauböck, M., Benisty, M., et al. (2018). Detection of the gravitational redshift in the orbit of the star S2 near the Galactic centre massive black hole. *Astronomy & Astrophysics*, 615:L15.
- Greene, J. E., Strader, J., and Ho, L. C. (2020). Intermediate-mass black holes. *Annual Review of Astronomy and Astrophysics*, 58:257–312.

- Häberle, M., Neumayer, N., Bellini, A., Libralato, M., Clontz, C., Seth, A. C., et al. (2024a). oMEGACat. II. Photometry and proper motions for 1.4 million stars in Omega Centauri and its rotation in the plane of the sky. *The Astrophysical Journal*, 970:192.
- Häberle, M., Neumayer, N., Clontz, C., Seth, A. C., Bellini, A., et al. (2025). oMEGACat. VI. Analysis of the overall kinematics of Omega Centauri in 3D: Velocity dispersion, kinematic distance, anisotropy, and energy equipartition. *The Astrophysical Journal*, 983:95.
- Häberle, M., Neumayer, N., Seth, A., Bellini, A., Libralato, M., Anderson, J., Whitaker, M., van de Ven, G., Watkins, L. L., Clontz, C., et al. (2024b). Fast-moving stars around an intermediate-mass black hole in ω Centauri. *Nature*, 631(8020):285–289.
- Haggard, D., Cool, A. M., Heinke, C. O., van der Marel, R., Cohn, H. N., Lugger, P. M., and Anderson, J. (2013). A deep Chandra X-ray limit on the putative IMBH in Omega Centauri. *The Astrophysical Journal Letters*, 773:L31.
- Harris, W. E. (1996). A catalog of parameters for globular clusters in the Milky Way. *The Astronomical Journal*, 112:1487. 2010 edition: arXiv:1012.3224.
- Hilker, M. and Richtler, T. (2000). ω Centauri – a former nucleus of a dissolved dwarf galaxy? New evidence from Strömgen photometry. *Astronomy & Astrophysics*, 362:895–909.
- Hills, J. G. (1988). Hyper-velocity and tidal stars from binaries disrupted by a massive Galactic black hole. *Nature*, 331:687–689.
- Hsiao, T. Y.-Y., Goto, T., Hashimoto, T., Santos, D. J. D., On, A. Y. L., Kilerci-Eser, E., Wong, Y. H. V., Kim, S. J., Wu, C. K.-W., Ho, S. C.-C., and Lu, T.-Y. (2021). A dyson sphere around a black hole. *Monthly Notices of the Royal Astronomical Society*, 506(2):1723–1732.
- Huang, B.-L., Tao, Z.-Z., and Zhang, T.-J. (2026a). Phase-space crystallization in Galactic globular clusters: A Gaia-based metric and implications for technosignature searches. *The Astrophysical Journal*. Accepted.
- Huang, B.-L., Tao, Z.-Z., Zhang, T.-J., and Gajjar, V. (2026b). The FAST-SETI Milky Way globular cluster survey. I. A pilot multibeam on-the-fly search of five globular clusters at the L band. *The Astronomical Journal*, 171(1):51.
- Ibata, R. A., Bellazzini, M., Malhan, K., Martin, N., and Bianchini, P. (2019). Identification of the long stellar stream of the prototypical massive globular cluster ω Centauri. *Nature Astronomy*, 3:667–672.
- Ivezić, Ž., Kahn, S. M., Tyson, J. A., Abel, B., Acosta, E., Allsman, R., et al. (2019). LSST: From science drivers to reference design and anticipated data products. *The Astrophysical Journal*, 873:111.
- Jonas, J. L. and MeerKAT Team (2016). The MeerKAT radio telescope. In *Proceedings of MeerKAT Science: On the Pathway to the SKA*, volume MeerKAT2016 of *Proceedings of Science*, page 001.
- Lacki, B. C. and DiKerby, S. (2025). Possibilities for SETI at high energy. White paper submitted to the 2025 NASA DARES RFI.
- Loeb, A. (2024). Comment on “No black holes from light”. Preprint; reply by original authors at arXiv:2408.11097.

- Mahida, A. D. et al. (2026). No evidence for accretion around the intermediate-mass black hole in Omega Centauri. *The Astrophysical Journal*, 996:122.
- Mandel, I., Brown, D. A., Gair, J. R., and Miller, M. C. (2008). Rates and characteristics of intermediate mass ratio inspirals detectable by Advanced LIGO. *The Astrophysical Journal*, 681:1431–1447.
- Merloni, A., Heinz, S., and di Matteo, T. (2003). A fundamental plane of black hole activity. *Monthly Notices of the Royal Astronomical Society*, 345:1057–1076.
- Nitschai, M. S., Neumayer, N., Clontz, C., Häberle, M., Seth, A. C., Husser, T.-O., et al. (2023). oMEGACat. I. MUSE spectroscopy of 300,000 stars within the half-light radius of ω Centauri. *The Astrophysical Journal*, 958:8.
- Noyola, E., Gebhardt, K., and Bergmann, M. (2008). Gemini and Hubble Space Telescope evidence for an intermediate-mass black hole in ω Centauri. *The Astrophysical Journal*, 676:1008–1015.
- Pierre Auger Collaboration, Aab, A., Abreu, P., Aglietta, M., et al. (2017). Observation of a large-scale anisotropy in the arrival directions of cosmic rays above 8×10^{18} eV. *Science*, 357:1266–1270.
- Prager, B. J., Ransom, S. M., Freire, P. C. C., Hessels, J. W. T., Stairs, I. H., et al. (2017). Using long-term millisecond pulsar timing to obtain physical characteristics of the bulge globular cluster Terzan 5. *The Astrophysical Journal*, 845:148.
- Socas-Navarro, H., Haqq-Misra, J., Wright, J. T., Kopparapu, R., Benford, J., Davis, R., and TechnoClimes 2020 workshop participants (2021). Concepts for future missions to search for technosignatures. *Acta Astronautica*, 182:446–453.
- Soltis, J., Casertano, S., and Riess, A. G. (2021). The parallax of ω Centauri measured from Gaia EDR3 and a direct, geometric calibration of the tip of the red giant branch and the Hubble constant. *The Astrophysical Journal Letters*, 908:L5.
- Souza, S., Neumayer, N., Seth, A. C., Wang, Z., Clontz, C., Häberle, M., Nitschai, M. S., Smith, P. J., Matsuno, T., et al. (2026). oMEGACat. X. shedding light on the disrupted dwarf galaxy of Omega Centauri. *arXiv e-prints*, page arXiv:2603.23589. Submitted to ApJ.
- Stappers, B. and Kramer, M. (2016). An update on TRAPUM. In *Proceedings of MeerKAT Science: On the Pathway to the SKA*, volume MeerKAT2016 of *Proceedings of Science*, page 009.
- Strader, J., Chomiuk, L., Maccarone, T. J., Miller-Jones, J. C. A., Seth, A. C., Heinke, C. O., and Sivakoff, G. R. (2012). No evidence for intermediate-mass black holes in globular clusters: Strong constraints from the JVL. *The Astrophysical Journal Letters*, 750:L27.
- Swanson, T. (2026a). Engineered intermediate-mass black hole systems: Infrastructure constraints, observable residue, and a multi-messenger adjudication framework. Preprint; omegacentauri.me.
- Swanson, T. (2026b). Inward resolutions of the Fermi paradox: A critical review of migration down thermodynamic gradients. Preprint; omegacentauri.me/paper.html.
- Swanson, T. (2026c). The macro transcension hypothesis: Spinning black holes in dense stellar clusters as thermodynamic attractors for advanced civilizations, with Omega Centauri as an observational test bed. Preprint; omegacentauri.me/paper.html.

- Tarter, J. (2001). The search for extraterrestrial intelligence (SETI). *Annual Review of Astronomy and Astrophysics*, 39:511–548.
- The KM3NeT Collaboration (2025). Observation of an ultra-high-energy cosmic neutrino with KM3NeT. *Nature*, 638(8050):376–382.
- Tremou, E., Strader, J., Chomiuk, L., Shishkovsky, L., Maccarone, T. J., Miller-Jones, J. C. A., et al. (2018). The MAVERIC survey: Still no evidence for accreting intermediate-mass black holes in globular clusters. *The Astrophysical Journal*, 862:16.
- van der Marel, R. P. and Anderson, J. (2010). New limits on an intermediate-mass black hole in Omega Centauri. II. Dynamical models. *The Astrophysical Journal*, 710:1063–1088.
- WFIRST Astrometry Working Group, Sanderson, R. E., Bellini, A., Casertano, S., Lu, J. R., Melchior, P., et al. (2019). Astrometry with the Wide-Field Infrared Space Telescope. *Journal of Astronomical Telescopes, Instruments, and Systems*, 5:044005.
- Whitaker, M., Kerr, E., Seth, A., Häberle, M., Strader, J., Anderson, J., Bellini, A., Clontz, C., Freeman, Z., Griggio, M., Kamann, S., Libralato, M., Neumayer, N., González Prieto, E., Rodriguez, C. L., Saracino, S., Smith, P., van de Ven, G., and Wang, Z. (2026). A long period stellar-mass black hole binary in ω Centauri. *The Astrophysical Journal Letters*.
- Wright, J. T., Haqq-Misra, J., Frank, A., Kopparapu, R., Lingam, M., and Sheikh, S. Z. (2022). The case for technosignatures: Why they may be abundant, long-lived, highly detectable, and unambiguous. *The Astrophysical Journal Letters*, 927(2):L30.
- Wright, J. T., Mullan, B., Sigurdsson, S., and Povich, M. S. (2014). The $\hat{g}$ infrared search for extraterrestrial civilizations with large energy supplies. I. Background and justification. *The Astrophysical Journal*, 792(1):26.
- Zocchi, A., Gieles, M., and Hénault-Brunet, V. (2019). The effect of stellar-mass black holes on the central kinematics of ω Cen: a cautionary tale for IMBH interpretations. *Monthly Notices of the Royal Astronomical Society*, 482:4713–4725.